

**TUKOL MAX ACTION SEVERE CONGESTION AND COUGH- dextromethorphan  
hbr, guaifenesin, phenylephrine hcl liquid  
Genomma Lab USA, Inc**

-----  
**Tukol MAX ACTION Severe Congestion & Cough**

**Drug Facts**

**Active ingredients  
(in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCL 10 mg

**Purpose**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- Helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping an MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are**

taking any other oral nasal decongestant or stimulant.

**Do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache

These could be signs of a serious condition.

**If pregnant or breast-feeding**

ask a professional before use

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- Do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter

---

age	dose
adults and children 12 years of age and older	20 mL every 4 hours
children under 12 years of age	do not use

---

**Other information**

- Each 20 mL dose contains: sodium 10 mg
- store between 15-30°C (59-86 °F)
- do not refrigerate

**Inactive Ingredients**

Anhydrous citric acid, edetate disodium, FD&C Blue # 1, FD&C Red # 40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions or comments?**

1-877-994-3666

Monday to Friday from 8 AM to 6 PM, Central time

**Tukol MAX ACTION Severe Congestion & Cough**

# Tukol<sup>®</sup>

## MAX ACTION

DO NOT USE IF PRINTED SEAL UNDER CAP IS  
TORN OR MISSING / NO SE USE SI EL SELLO IMPRESO  
DEBAJO DE LA TAPA ESTA ROTO O FALTA

# Tukol<sup>®</sup>

## MAX ACTION

### SEVERE CONGESTION & COUGH

### CONGESTIÓN Y TOS SEVERA

Dextromethorphan HBr / Guaifenesin / Phenylephrine HCl  
Dextrometorfano HBr / Guaifenesina / Fenilefrina HCl

Ages / Edades  
12+

MAX  
STRENGTH/  
MÁXIMA  
POTENCIA

Relieves / Alivia:

- COUGH / TOS
- NASAL CONGESTION / CONGESTIÓN NASAL
- CHEST CONGESTION / CONGESTIÓN DE PECHO

6 FL OZ (177 mL)



#### Drug Facts (continued)

##### Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter

age	dose
adults and children 12 years of age and older	20 mL every 4 hours
children under 12 years of age	do not use

##### Other information

- each (20 mL) dose contains: sodium 10 mg
- store between 15-30°C (59-86°F) do not refrigerate.

##### Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-877-994-3666  
Monday to Friday from 8 am to 6 pm, Central Time.

#### Información del Medicamento

##### Ingredientes activos (en cada 20 mL)

Dextrometorfano HBr, 20 mg.....Inhibidor de la tos  
Guaifenesina, 400 mg.....Expectorante  
Fenilefrina HCl, 10 mg.....Descongestionante nasal

##### Propósitos

##### Usos

- ayuda a desprender las flemas (mucosidad) y a adelgazar las secreciones bronquiales para drenar las vías respiratorias
- alivia temporalmente los siguientes síntomas que se presentan con el resfriado:
  - congestión nasal
  - tos debida a irritación leve de garganta y bronquios.

##### Advertencias

No utilizarlo si usted está tomando un inhibidor de la monoamina oxidasa (IMAO) de prescripción (ciertos medicamentos para la depresión, condiciones psiquiátricas o emocionales, o enfermedad de Parkinson), o durante 2 semanas después de suspender el medicamento IMAO. Si usted desconoce si su medicamento de prescripción contiene un IMAO, consulte a su médico o farmacéutico antes de tomar este producto.

##### Consulte a su médico antes de usar, si padece

- enfermedad cardíaca
- hipertensión
- enfermedad de la tiroides
- diabetes
- dificultad al orinar debido a dilatación de la glándula prostática
- tos que se presenta con exceso de flema (mucosidad)
- tos persistente o crónica tal como ocurre con el tabaquismo, asma, bronquitis crónica o enfisema.

#### Información del Medicamento (continuación)

Consulte a su médico o farmacéutico antes de utilizarlo si usted está tomando cualquier otro descongestionante o estimulante nasal oral.

##### No exceda la dosis recomendada.

- Suspenda su uso y consulte a su médico si
  - se siente nervioso, mareado o con insomnio
  - los síntomas no mejoran en los siguientes 7 días o se acompañan de fiebre
  - la tos persiste por más de 7 días, regresa, o se presenta acompañada por fiebre, erupción o dolor de cabeza persistente. Estos podrían ser signos de una condición grave.

En caso de embarazo o lactancia, consulte a un profesional médico antes de su uso.

Manténgase fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o contacte al Centro de Control de Envenenamiento de manera inmediata.

##### Indicaciones

- no tomar más de 6 dosis en un período de 24 horas
- este producto de concentración para adultos no está previsto para su uso en niños menores de 12 años de edad
- medir solo con la copa dosificadora incluida
- mantenga la copa dosificadora con este producto
- mL = mililitro

edad	dosis
adultos y niños de 12 años de edad y mayores	20 mL cada 4 horas
niños menores de 12 años	no usar

##### Otra información

- cada (20 mL) contiene: sodio 10 mg
- almacenar entre 15 y 30°C (59 a 86°F)
- no refrigerar.

##### Ingredientes inactivos

ácido cítrico anhidro, edetato disódico, FD&C azul #1, FD&C rojo #40, sabores, glicerina, propilenglicol, galato de propilo, agua purificada, benzoato sódico, sorbitol, sucralosa, goma xantana.

##### ¿Preguntas o comentarios?

1-877-994-3666 Lunes a Viernes de 8 am a 6 pm, Hora del Centro.

#### Drug Facts

##### Active ingredients Purposes (in each 20 mL)

Dextromethorphan HBr, 20 mg.....Cough suppressant  
Guaifenesin, 400 mg.....Expectorant  
Phenylephrine HCl, 10 mg.....Nasal decongestant

##### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation.

##### Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

##### Do not exceed recommended dosage.

##### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Distributed by / Distribuido por:  
Genomma Lab USA Inc.,  
Houston, TX 77098  
BX-064

2000004535 V-3



LOT No. / LOTE  
EXP.

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING**

**DEXTROMETHORPHAN HBr/GUAIFENESIN/PHENYLEPHRINE HCL**

**Relieves**

- Cough
- Nasal Congestion
- Chest Congestion

Ages 12+

6 FL OZ (177 mL)

Distributed by:  
Genomma Lab USA, Inc.  
Houston, TX 77098

**TUKOL MAX ACTION SEVERE CONGESTION AND COUGH**

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50066-517
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	10 mg in 20 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYL GALLATE</b> (UNII: 8D45N7V92)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-517-25	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/16/2016	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/16/2016	

**Labeler** - Genomma Lab USA, Inc (832323534)

Revised: 1/2024

Genomma Lab USA, Inc