

**COLD, FLU AND SORE THROAT- acetaminophen, dextromethorphan hbr,
guaifenesin, phenylephrine hcl solution
Family Dollar Services Inc**

Family Wellness 44-005

Active ingredients (in each 20 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg
Guaifenesin 400 mg
Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - stuffy nose
 - cough
 - minor aches and pains
 - sore throat
 - sinus congestion and pressure
 - nasal congestion
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- rash
- blisters

- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- liver disease
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- heart disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- **each 20 mL contains:** sodium 10 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

FAMILY Wellness™

COMPARE TO THE
ACTIVE INGREDIENTS IN
MAXIMUM STRENGTH
MUCINEX® FAST-MAX®
COLD, FLU & SORE THROAT*

MAXIMUM STRENGTH
**COLD, FLU &
SORE THROAT**

Acetaminophen

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Pain Reliever/Fever Reducer

Cough Suppressant
Expectorant
Nasal Decongestant

- Relieves body pain, headache, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- Relieves sinus pressure & congestion

AGES
12 YEARS
& OVER

MIXED BERRY
FLAVORED

NDC 55319-705-45

6 FL OZ (177 mL)

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

PARENTS:

**Learn about teen medicine abuse
www.StopMedicineAbuse.org**

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat. 50844 ORG052300545

**DISTRIBUTED BY: MIDWOOD BRANDS LLC, 500 VOLVO PKWY, CHESAPEAKE, VA
23320 USA**

NOT 100% SATISFIED?

Return within 30 days to
the store of purchase for
a refund (with receipt)
or exchange.

FAMILY Wellness™

COMPARE TO THE ACTIVE INGREDIENTS IN MAXIMUM STRENGTH MUCINEX® FAST-MAX® COLD, FLU & SORE THROAT*

MAXIMUM STRENGTH
COLD, FLU & SORE THROAT

AGES 12 YEARS & OVER

Acetaminophen
Dextromethorphan HBr
Guaifenesin
Phenylephrine HCl
Pain Reliever/Fever Reducer
Cough Suppressant
Expectorant
Nasal Decongestant



MIXED BERRY FLAVORED

- Relieves body pain, headache, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- Relieves sinus pressure & congestion

F-005-45
ORG
NDC 55319-705-45

6 FL OZ (177 mL)

PARENTS:
Learn about how medicines abuse
www.StopMedicineAbuse.org

PEEL BACK TAB TO READ COMPLETE DRUG FACTS AND INFORMATION

Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each 20 mL)

Active ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - stuffy nose
 - cough
 - minor aches and pains
 - sore throat
 - sinus congestion and pressure
 - nasal congestion
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat. 50914-ORG0522005-45
DISTRIBUTED BY: MIDWOOD BRANDS LLC, 500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA

NOT 100% SATISFIED? Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

0 32251 92381 4 SKU 903758 B-005-45 ORG

No Print / No Varnish Area
Lot # and Exp. Info

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- rash
- blisters
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- liver disease
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- heart disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur

Drug Facts (continued)

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 10 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments? 1-800-426-9391

Family Wellness 44-005

COLD, FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-705
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-705-45	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/28/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/28/2024	

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(55319-705) , pack(55319-705)

Revised: 6/2024

Family Dollar Services Inc