

CALMICID AC ACID REDUCER- loratadine tablet, film coated
Melaleuca, Inc.

Calmicid AC Content of Labeling

Purpose

Acid reducer

Use

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over three months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Active ingredient (in each tablet)

Famotidine 20 mg

Other information

- store at 20°-25°C (68°-77°F)
- protect from moisture and light

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions or comments?

1-800-282-3000

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Other information

- read the directions and warnings before use
- keep the carton, it contains important information.



194-001 ELA17A-1 **ELA 17A**



Do not use if printed foil under cap is broken or missing

Active ingredient (in each tablet)

Famotidine 20 mg

Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ if you have kidney disease, except under the advice and supervision of a doctor ■ with other acid reducers

Ask a doctor before use if you have

■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with **lightheadedness, sweating, or dizziness** ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness ■ frequent **chest pain** ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ Stomach pain

Stop use and ask a doctor if

■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ adults and children 12 years and over ■ to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew ■ to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours ■ children under 12 years: ask a doctor

Other information

■ read the directions and warnings before use ■ keep the carton. It contains important information ■ store at 20°-25° C (68°-77° F) ■ protect from moisture and light

Questions? 1-800-282-3000

Dist. by Melaleuca, Inc.
Idaho Falls, ID 83402-6003
To order: 1-800-282-3000
www.melaleuca.com
#Z256 1010U

CODE AREA

• no varnish • no color

: 19471 MB F1

MB
Melaleuca
1-800-282-3000

TO ARTIST:
DO NOT use color behind the Part #.
DO NOT change position of part #.

CALMICID AC ACID REDUCER

loratadine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54473-203
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (biconvex)	Size	8mm
Flavor		Imprint Code	L194
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54473-203-50	1 in 1 BOX	01/01/2018	
1	NDC:54473-203-01	50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	01/20/2011	

Labeler - Melaleuca, Inc. (139760102)

Revised: 1/2023

Melaleuca, Inc.