

**CVS CHILDRENS NIGHTTIME MULTI SYMPTOM COLD- acetaminophen ,  
diphenhydramine hydrochloride,phenylephrine hydrochloride liquid  
CVS PHARMACY**

-----  
**CVS Children's Nighttime Multi-Symptoms Cold 4 FL OZ (180 mL)**

**Drug Facts**

---

**Active ingredients (in each 10 mL) Purposes**

<b>Acetaminophen 325 mg</b>	<b>Pain reliever/fever reducer</b>
Diphenhydramine HCl 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

---

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - runny nose
  - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

**Warnings**

**Liver Warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:**Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:**If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- To make child sleepy
- with any other drug containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if the child have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if the child is**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not use more than directed (see Overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 years to 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in 24-hours period
- children under 6 years of age: do not use

### **Other information**

- each 10 mL contains: **sodium 4 mg**
- store at room temperature
- do not refrigerate

### **Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, natural and artificial flavor, potassium citrate, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

### **Questions or comments?**

**1-866-467-2748**

### **PRINCIPAL DISPLAY PANEL**

Compare to the active ingredients in Children's Mucinex<sup>®</sup> Night Time Multi-Symptom Cold

NDC# 51316-733-04

### **Children's**

### **Night Time Multi-Symptom Cold**

- **Acetaminophen** Pain Reliever/Fever Reducer
- Diphenhydramine HCl Antihistamine/Cough Suppressant
- Phenylephrine HCl Nasal Decongestant

Relieves Stuffy Nose

Controls Cough

Relieves Fever & Sore Throat

Relieves Runny Nose & Sneezing

Dosing Cup Included

**For Ages 6 & Over**

**Very Berry Flavor**

Naturally and Artificially Flavored

4 FL OZ (118 mL)

\*This product is not manufactured or distributed by RB Health (US), distributor Children's Mucinex® Nighttime Multi-Symptoms Cold..

**TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.**

Distributed by:

**TAMPER-EVIDENT: DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS BORN OR MISSING.**

**Drug Facts**

**Active Ingredients**  
(in each 10 mL)

Acetaminophen 325 mg..... Pain reliever/Fever reducer  
Diphenhydramine HCl 12.5 mg..... Antihistamine/Cough suppressant  
Phenylephrine HCl 5 mg..... Nasal decongestant

**Uses**

Temporarily relieves these common cold and flu symptoms:

- fever and chills
- headache
- throat pain
- runny nose
- cough
- stuffy nose
- sinus pain
- aches and pains
- sores throat
- congestion
- nasal congestion
- throat irritation
- runny nose
- cough
- stuffy nose
- sinus pain

**Warnings**

**Overdose warning:** This product contains acetaminophen. Severe liver damage may occur if you take it for more than 15 doses in 24 hours, which is the maximum daily amount.

**Do not take:** Do not take other drugs containing acetaminophen.

**Caution:** Ask a doctor before use if you are taking MAO inhibitors (certain drugs for depression, psychostimulants, or anesthetic agents) or have taken MAO inhibitors within 14 days of starting this product.

**Other information:**

- each 10 mL contains sodium 4 mg
- store at room temperature
- do not refrigerate

**Questions or comments?** 1-866-487-4748

©2015 RB Health, Inc. All rights reserved. RB Health, Inc. is not responsible for any damage to your property or health caused by the use of this product.

**CVS Health.**

**Children's Multi-Symptom Nighttime Cold**

**ACETAMINOPHEN 325 mg**  
Pain reliever/Fever reducer

**DIPHENHYDRAMINE HCl 12.5 mg**  
Antihistamine/Cough suppressant

**PHENYLEPHRINE HCl 5 mg**  
Nasal decongestant

**CVS Health.** Compares to the active ingredients in Children's Mucinex® Night Time Multi-Symptom Cold\*\*

**Children's Multi-Symptom Nighttime Cold**

**ACETAMINOPHEN 325 mg**  
Pain reliever/Fever reducer

**DIPHENHYDRAMINE HCl 12.5 mg**  
Antihistamine/Cough suppressant

**PHENYLEPHRINE HCl 5 mg**  
Nasal decongestant

**Relieves:**  
Nasal congestion

Reduces fever  
Soothes cough & sore throat

No added alcohol



**Very Berry Flavor**  
Naturally & Artificially Flavored

**Ages 6 to 11 Years**  
Dosage Cup Included

**4 FL OZ (118 mL)**

**Actual Bottle Size on Side Panel**

**LOT:**

**EXP:**

**FPO 80%  
UPC# 050428561553**

**PARENTS:**  
www.cvs.com/parents

**IMPORTANT:**  
Keep this carton for future reference on full labeling.

**RECYCLE**  
PAPER BOX PLASTIC BOTTLE PLASTIC CUP

**Package Contains One Bottle**

**Actual Size**

acetaminophen , diphenhydramine hydrochloride,phenylephrine hydrochloride liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:51316-733
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>POTASSIUM CITRATE</b> (UNII: EE90ONI6FF)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-733-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/20/2023	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M010	02/20/2023	

**Labeler** - CVS PHARMACY (062312574)

Revised: 8/2024

CVS PHARMACY