

DROSPIRENONE AND ETHINYL ESTRADIOL- drospirenone and ethinyl estradiol Pharmaceuticals International, Inc. (Pii)

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DROSPIRENONE AND ETHINYL ESTRADIOL TABLETS safely and effectively. See full prescribing information for DROSPIRENONE AND ETHINYL ESTRADIOL TABLETS.

DROSPIRENONE and ETHINYL ESTRADIOL tablets, for oral use
Initial U.S. Approval: 2001

WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS

See full prescribing information for complete boxed warning.

- Women over 35 years old who smoke should not use Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg (4).
- Cigarette smoking increases the risk of serious cardiovascular events from combination oral contraceptive (COC) use. (4)

RECENT MAJOR CHANGES

Warnings and Precautions, Hyperkalemia (5.2)

6/2015

INDICATIONS AND USAGE

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are an estrogen/progestin COC, indicated for use by women to:

- Prevent pregnancy. (1.1)
- Treat moderate acne for women at least 14 years old only if the patient desires an oral contraceptive for birth control. (1.3)

DOSAGE AND ADMINISTRATION

- Take one tablet daily by mouth at the same time every day. (2.1)
- Tablets must be taken in the order directed on the blister pack. (2.1)

DOSAGE FORMS AND STRENGTHS

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg consist of 28 film-coated, biconvex tablets in the following order (3):

- 24 pink tablets, each containing 3 mg drospirenone, USP (DRSP) (micronized) and 0.02 mg ethinyl estradiol, USP (EE) (micronized)
- 4 white inert tablets

CONTRAINDICATIONS

- Renal impairment (4)
- Adrenal insufficiency (4)
- A high risk of arterial or venous thrombotic diseases (4)
- Undiagnosed abnormal uterine bleeding (4)
- Breast cancer or other estrogen- or progestin-sensitive cancer (4)
- Liver tumors or liver disease (4)
- Pregnancy (4)

WARNINGS AND PRECAUTIONS

- **Vascular risks:** Stop Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if a thrombotic event occurs. Stop at least 4 weeks before and through 2 weeks after major surgery. Start no earlier than 4 weeks after delivery, in women who are not breastfeeding. (5.1) COCs containing DRSP may be associated with a higher risk of venous thromboembolism (VTE) than COCs containing levonorgestrel or some other progestins. Before initiating Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg in a new COC user or a woman who is switching from a contraceptive that does not contain DRSP, consider the risks and benefits of a DRSP-containing COC in light of her risk of a VTE. (5.1)

- **Hyperkalemia:** DRSP has antimineralocorticoid activity. Do not use in patients predisposed to hyperkalemia. Check serum potassium concentration during the first treatment cycle in women on long-term treatment with medications that may increase serum potassium concentration. (5.2, 7.1, 7.2)
- **Liver disease:** Discontinue Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if jaundice occurs. (5.4)
- **High blood pressure:** Do not prescribe Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg for women with uncontrolled hypertension or hypertension with vascular disease. (5.5)
- **Carbohydrate and lipid metabolic effects:** Monitor prediabetic and diabetic women taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg. Consider an alternative contraceptive method for women with uncontrolled dyslipidemia. (5.7)
- **Headache:** Evaluate significant change in headaches and discontinue Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if indicated. (5.8)
- **Uterine bleeding:** Evaluate irregular bleeding or amenorrhea. (5.9)

----- **ADVERSE REACTIONS** -----

- The most frequent adverse reactions ($\geq 2\%$) in contraception and acne clinical trials were: headache/migraine (6.7%), menstrual irregularities (4.7%), nausea/vomiting (4.2%), breast pain/tenderness (4.0%) and mood changes (2.2%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Pharmaceuticals International, Inc. (Pii) at 1-410-584-0001 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

Drugs or herbal products that induce certain enzymes (for example, CYP3A4) may decrease the effectiveness of COCs or increase breakthrough bleeding. Counsel patients to use a back-up or alternative method of contraception when enzyme inducers are used with COCs. (7.1)

----- **USE IN SPECIFIC POPULATIONS** -----

Nursing Mothers: Not recommended; can decrease milk production. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 3/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS

1 INDICATIONS AND USAGE

- 1.1 Oral Contraceptive
- 1.3 Acne

2 DOSAGE AND ADMINISTRATION

- 2.1 How to Take Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg
- 2.2 How to Start Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg
- 2.3 Advice in case of Gastrointestinal Disturbances

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Thromboembolic Disorders and Other Vascular Problems
- 5.2 Hyperkalemia
- 5.3 Carcinoma of the Breasts and Reproductive Organs
- 5.4 Liver Disease
- 5.5 High Blood Pressure
- 5.6 Gallbladder Disease
- 5.7 Carbohydrate and Lipid Metabolic Effects
- 5.8 Headache
- 5.9 Bleeding Irregularities
- 5.10 COC Use Before or During Early Pregnancy
- 5.11 Depression
- 5.12 Interference with Laboratory Tests

5.13 Monitoring

5.14 Other Conditions

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Effects of Other Drugs on Combined Oral Contraceptives

7.2 Effects of Combined Oral Contraceptives on Other Drugs

7.3 Interference with Laboratory Tests

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Patients with Renal Impairment

8.7 Patients with Hepatic Impairment

8.8 Race

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Oral Contraceptive Clinical Trial

14.3 Acne Clinical Trials

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Storage

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS

Cigarette smoking increases the risk of serious cardiovascular events from combination oral contraceptives (COC) use. This risk increases with age, particularly in women over 35 years of age, and with the number of cigarettes smoked. For this reason, COCs should not be used by women who are over 35 years of age and smoke. [See Contraindications (4)].

1 INDICATIONS AND USAGE

1.1 Oral Contraceptive

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are indicated for use by women to prevent pregnancy.

1.3 Acne

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are indicated for the treatment of moderate acne vulgaris in women at least 14 years of age, who have no known contraindications to oral contraceptive therapy and have achieved menarche. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be used for the treatment of acne only if the patient desires an oral contraceptive for birth control.

2 DOSAGE AND ADMINISTRATION

2.1 How to Take Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg

Take one tablet by mouth at the same time every day. The failure rate may increase when pills are missed or taken incorrectly.

To achieve maximum contraceptive effectiveness, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg must be taken exactly as directed, in the order directed on the blister pack. Single missed pills should be taken as soon as remembered.

2.2 How to Start Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg

Instruct the patient to begin taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg either on the first day of her menstrual period (Day 1 Start) or on the first Sunday after the onset of her menstrual period (Sunday Start).

Day 1 Start

During the first cycle of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg use, instruct the patient to take one pink Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg daily, beginning on Day 1 of her menstrual cycle. (The first day of menstruation is Day 1.) She should take one pink Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg daily for 24 consecutive days, followed by one white inert tablet daily on Days 25 through 28. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be taken in the order directed on the package at the same time each day, preferably after the evening meal or at bedtime with some liquid, as needed. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg can be taken without regard to meals. If Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are first taken later than the first day of the menstrual cycle, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should not be considered effective as a contraceptive until after the first 7 consecutive days of product administration. Instruct the patient to use a non-hormonal contraceptive as back-up during the first 7 days. The possibility of ovulation and conception prior to initiation of medication should be considered.

Sunday Start

During the first cycle of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg use, instruct the patient to take one pink Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg daily, beginning on the first Sunday after the onset of her menstrual period. She should take one pink Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg daily for 24 consecutive days, followed by one white inert tablet daily on Days 25 through 28. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be taken in the order directed on the package at the same time each day, preferably after the evening meal or at bedtime with some liquid, as needed. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg can be taken without regard to meals. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should not be considered effective as a contraceptive until after the first 7 consecutive days of product administration. Instruct the patient to use a non-hormonal contraceptive as back-up during the first 7 days. The possibility of ovulation and conception prior to initiation of medication should be considered.

The patient should begin her next and all subsequent 28-day regimens of Drospirenone and Ethinyl

Estradiol Tablets, USP 3 mg/0.02 mg on the same day of the week that she began her first regimen, following the same schedule. She should begin taking her pink tablets on the next day after ingestion of the last white tablet, regardless of whether or not a menstrual period has occurred or is still in progress. Anytime a subsequent cycle of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg is started later than the day following administration of the last white tablet, the patient should use another method of contraception until she has taken a pink Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg daily for seven consecutive days.

When switching from a different birth control pill

When switching from another birth control pill, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started on the same day that a new pack of the previous oral contraceptive would have been started.

When switching from a method other than a birth control pill

When switching from a transdermal patch or vaginal ring, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started when the next application would have been due. When switching from an injection, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started when the next dose would have been due. When switching from an intrauterine contraceptive or an implant, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started on the day of removal.

Withdrawal bleeding usually occurs within 3 days following the last pink tablet. If spotting or breakthrough bleeding occurs while taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg, instruct the patient to continue taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg by the regimen described above. Counsel her that this type of bleeding is usually transient and without significance; however, advise her that if the bleeding is persistent or prolonged, she should consult her healthcare provider.

Although the occurrence of pregnancy is low if Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are taken according to directions, if withdrawal bleeding does not occur, consider the possibility of pregnancy. If the patient has not adhered to the prescribed dosing schedule (missed one or more active tablets or started taking them on a day later than she should have), consider the possibility of pregnancy at the time of the first missed period and take appropriate diagnostic measures. If the patient has adhered to the prescribed regimen and misses two consecutive periods, rule out pregnancy. Discontinue Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if pregnancy is confirmed.

The risk of pregnancy increases with each active pink tablet missed. For additional patient instructions regarding missed pills, see the "**WHAT TO DO IF YOU MISS PILLS**" section in the **FDA Approved Patient Labeling**. If breakthrough bleeding occurs following missed tablets, it will usually be transient and of no consequence. If the patient misses one or more white tablets, she should still be protected against pregnancy provided she begins taking a new cycle of pink tablets on the proper day.

For postpartum women who do not breastfeed or after a second trimester abortion, start Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg no earlier than 4 weeks postpartum due to the increased risk of thromboembolism. If the patient starts on Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg postpartum and has not yet had a period, evaluate for possible pregnancy, and instruct her to use an additional method of contraception until she has taken Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg for 7 consecutive days.

2.3 Advice in case of Gastrointestinal Disturbances

In case of severe vomiting or diarrhea, absorption may not be complete and additional contraceptive measures should be taken. If vomiting occurs within 3-4 hours after tablet-taking, this can be regarded as a missed tablet.

3 DOSAGE FORMS AND STRENGTHS

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are available in blister packs.

Each blister pack (28 film-coated tablets) contains in the following order:

- 24 pink tablets each containing 3 mg drospirenone, USP (DRSP) (micronized) and 0.02 mg ethinyl estradiol, USP (EE) (micronized)
- 4 white inert tablets

4 CONTRAINDICATIONS

Do not prescribe Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg to women who are known to have the following:

- Renal impairment
- Adrenal insufficiency
- A high risk of arterial or venous thrombotic diseases. Examples include women who are known to:
 - Smoke, if over age 35 [see *Boxed Warning and Warnings and Precautions (5.1)*]
 - Have deep vein thrombosis or pulmonary embolism, now or in the past [see *Warnings and Precautions (5.1)*]
 - Have cerebrovascular disease [see *Warnings and Precautions (5.1)*]
 - Have coronary artery disease [see *Warnings and Precautions (5.1)*]
 - Have thrombogenic valvular or thrombogenic rhythm diseases of the heart (for example, subacute bacterial endocarditis with valvular disease, or atrial fibrillation) [see *Warnings and Precautions (5.1)*]
 - Have inherited or acquired hypercoagulopathies [see *Warnings and Precautions (5.1)*]
 - Have uncontrolled hypertension [see *Warnings and Precautions (5.5)*]
 - Have diabetes mellitus with vascular disease [see *Warnings and Precautions (5.7)*]
 - Have headaches with focal neurological symptoms or have migraine headaches with or without aura if over age 35 [see *Warnings and Precautions (5.8)*]
- Undiagnosed abnormal uterine bleeding [see *Warnings and Precautions (5.9)*]
- Breast cancer or other estrogen- or progestin-sensitive cancer, now or in the past [see *Warnings and Precautions (5.3)*]
- Liver tumors, benign or malignant, or liver disease [see *Warnings and Precautions (5.4)* and *Use in Specific Populations (8.7)*]
- Pregnancy, because there is no reason to use COCs during pregnancy [see *Warnings and Precautions (5.10)* and *Use in Specific Populations (8.1)*]

5 WARNINGS AND PRECAUTIONS

5.1 Thromboembolic Disorders and Other Vascular Problems

Stop Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if an arterial or venous thrombotic (VTE) event occurs.

Based on presently available information on DRSP-containing COCs with 0.03 mg ethinyl estradiol (that is, Yasmin), DRSP-containing COCs may be associated with a higher risk of venous thromboembolism (VTE) than COCs containing the progestin levonorgestrel or some other progestins. Epidemiologic studies that compared the risk of VTE reported that the risk ranged from no increase to a three-fold increase. Before initiating use of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg in a new COC user or a woman who is switching from a contraceptive that does not contain DRSP, consider the risks and benefits of a DRSP-containing COC in light of her risk of a VTE. Known risk factors for VTE include smoking, obesity, and family history of VTE, in addition to other factors that contraindicate use of COCs [see *Contraindications (4)*].

A number of studies have compared the risk of VTE for users of Yasmin (which contains 0.03 mg of EE and 3 mg of DRSP) to the risk for users of other COCs, including COCs containing levonorgestrel. Those that were required or sponsored by regulatory agencies are summarized in Table 1.

Table 1: Estimates (Hazard Ratios) of Venous Thromboembolism Risk in Current Users of Yasmin Compared to Users of Oral Contraceptives that Contain Other Progestins

Epidemiologic Study (Author, Year of Publication) Population Studied	Comparator Product (all are low-dose COCs; with ≤ 0.04 mg of EE)	Hazard Ratio (HR) (95% CI)
i3 Ingenix (Seeger 2007) Initiators, including new users*	All COCs available in the US during the conduct of the study [†]	HR: 0.9 (0.5-1.6)
EURAS (Dinger 2007) Initiators, including new users*	All COCs available in Europe during the conduct of the study [‡]	HR: 0.9 (0.6-1.4)
	Levonorgestrel/EE	HR: 1.0 (0.6-1.8)
“FDA-funded study” (2011) New users* All users (i.e., initiation and continuing use of study combination hormonal contraception)	Other COCs available during the course of the study [§]	HR: 1.8 (1.3-2.4)
	Levonorgestrel/0.03 mg EE	HR: 1.6 (1.1-2.2)
	Other COCs available during the course of the study [§]	HR: 1.7 (1.4-2.1)
	Levonorgestrel/0.03 mg EE	HR: 1.5 (1.2-1.8)

* “New users” – no use of combination hormonal contraception for at least the prior 6 months


[†] Includes low-dose COCs containing the following progestins: norgestimate, norethindrone, levonorgestrel, desogestrel, norgestrel, medroxyprogesterone, or ethynodiol diacetate

[‡] Includes low-dose COCs containing the following progestins: levonorgestrel, desogestrel, dienogest, chlormadinone acetate, gestodene, cyproterone acetate, norgestimate, or norethindrone

[§] Includes low-dose COCs containing the following progestins: norgestimate, norethindrone, or levonorgestrel

In addition to these “regulatory studies,” other studies of various designs have been conducted. Overall, there are two prospective cohort studies (see Table 1): the US post-approval safety study Ingenix [Seeger 2007], the European post-approval safety study EURAS (European Active Surveillance Study) [Dinger 2007]. An extension of the EURAS study, the Long-Term Active Surveillance Study (LASS), did not enroll additional subjects, but continued to assess VTE risk. There are three retrospective cohort studies: one study in the US funded by the FDA (see Table 1), and two from Denmark [Lidegaard 2009, Lidegaard 2011]. There are two case-control studies: the Dutch MEGA study analysis [van Hylckama Vlieg 2009] and the German case-control study [Dinger 2010]. There are two nested case-control studies that evaluated the risk of non-fatal idiopathic VTE: the PharMetrics study [Jick 2011] and the GPRD study [Parkin 2011]. The results of all of these studies are presented in Figure 1.

Figure 1: VTE Risk with Yasmin Relative to LNG-Containing COCs (adjusted risk#)



Risk ratios displayed on logarithmic scale; risk ratio < 1 indicates a lower risk of VTE for DRSP, > 1 indicates an increased risk of VTE for DRSP.

*Comparator “Other COCs”, including LNG- containing COCs

† LASS is an extension of the EURAS study

#Some adjustment factors are indicated by superscript letters: a) Current heavy smoking, b) hypertension, c) obesity, d) family history, e) age, f) BMI, g) duration of use, h) VTE history, i) period of inclusion, j) calendar year, k) education, l) length of use, m) parity, n) chronic disease, o) concomitant medication, p) smoking, q) duration of exposure, r) site


(References: Ingenix [Seeger 2007]¹, EURAS (European Active Surveillance Study) [Dinger 2007]², LASS (Long-Term Active Surveillance Study) [Dinger, unpublished document on file], FDA-funded study [Sidney 2011]³, Danish [Lidegaard 2009]⁴, Danish re-analysis [Lidegaard 2011]⁵, MEGA study [van Hylckama Vlieg 2009]⁶, German Case-Control study [Dinger 2010]⁷, PharMetrics [Jick 2011]⁸, GPRD study [Parkin 2011]⁹)

Although the absolute VTE rates are increased for users of hormonal contraceptives compared to non-users, the rates during pregnancy are even greater, especially during the post-partum period (see Figure 2). The risk of VTE in women using COCs has been estimated to be 3 to 9 per 10,000 woman-years. The risk of VTE is highest during the first year of use. Data from a large, prospective cohort safety study of various COCs suggest that this increased risk, as compared to that in non-COC users, is greatest during the first 6 months of COC use. Data from this safety study indicate that the greatest risk of VTE is present after initially starting a COC or restarting (following a 4 week or greater pill-free interval) the same or a different COC.

The risk of thromboembolic disease due to oral contraceptives gradually disappears after COC use is discontinued.

Figure 2 shows the risk of developing a VTE for women who are not pregnant and do not use oral contraceptives, for women who use oral contraceptives, for pregnant women, and for women in the postpartum period. To put the risk of developing a VTE into perspective: If 10,000 women who are not pregnant and do not use oral contraceptives are followed for one year, between 1 and 5 of these women will develop a VTE.

Figure 2: Likelihood of Developing a VTE



* Pregnancy data based on actual duration of pregnancy in the reference studies. Based on a model assumption that pregnancy duration is nine months, the rate is 7 to 27 per 10,000 WY.

If feasible, stop Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg at least 4 weeks before and through 2 weeks after major surgery or other surgeries known to have an elevated risk of thromboembolism.

Start Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg no earlier than 4 weeks after delivery, in women who are not breastfeeding. The risk of postpartum thromboembolism decreases after the third postpartum week, whereas the risk of ovulation increases after the third postpartum week.

Use of COCs also increases the risk of arterial thromboses such as strokes and myocardial infarctions, especially in women with other risk factors for these events.

COCs have been shown to increase both the relative and attributable risks of cerebrovascular events (thrombotic and hemorrhagic strokes), although, in general, the risk is greatest among older (>35 years of age), hypertensive women who also smoke. COCs also increase the risk for stroke in women with other underlying risk factors.

Oral contraceptives must be used with caution in women with cardiovascular disease risk factors.

Stop Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if there is unexplained loss of vision, proptosis, diplopia, papilledema, or retinal vascular lesions. Evaluate for retinal vein thrombosis immediately. [See Adverse Reactions (6).]

5.2 Hyperkalemia

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg contain 3 mg of the progestin DRSP which has antimineralocorticoid activity, including the potential for hyperkalemia in high-risk patients, comparable to a 25 mg dose of spironolactone. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should not be used in patients with conditions that predispose to hyperkalemia (that is, renal impairment, hepatic impairment and adrenal insufficiency). Women receiving daily, long-term treatment for chronic conditions or diseases with medications that may increase serum potassium concentration

should have their serum potassium concentration checked during the first treatment cycle. Medications that may increase serum potassium concentration include ACE inhibitors, angiotensin-II receptor antagonists, potassium-sparing diuretics, potassium supplementation, heparin, aldosterone antagonists, and NSAIDs.

Consider monitoring serum potassium concentration in high-risk patients who take a strong CYP3A4 inhibitor long-term and concomitantly. Strong CYP3A4 inhibitors include azole antifungals (e.g. ketoconazole, itraconazole, voriconazole), HIV/HCV protease inhibitors (e.g., indinavir, boceprevir), and clarithromycin [see *Clinical Pharmacology* (12.3)].

5.3 Carcinoma of the Breasts and Reproductive Organs

Women who currently have or have had breast cancer should not use Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg because breast cancer is a hormonally-sensitive tumor.

There is substantial evidence that COCs do not increase the incidence of breast cancer. Although some past studies have suggested that COCs might increase the incidence of breast cancer, more recent studies have not confirmed such findings.

Some studies suggest that COCs are associated with an increase in the risk of cervical cancer or intraepithelial neoplasia. However, there is controversy about the extent to which these findings may be due to differences in sexual behavior and other factors.

5.4 Liver Disease

Discontinue Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if jaundice develops. Steroid hormones may be poorly metabolized in patients with impaired liver function. Acute or chronic disturbances of liver function may necessitate the discontinuation of COC use until markers of liver function return to normal and COC causation has been excluded.

Hepatic adenomas are associated with COC use. An estimate of the attributable risk is 3.3 cases/100,000 COC users. Rupture of hepatic adenomas may cause death through intra-abdominal hemorrhage.

Studies have shown an increased risk of developing hepatocellular carcinoma in long-term (>8 years) COC users. However, the attributable risk of liver cancers in COC users is less than one case per million users.

Oral contraceptive-related cholestasis may occur in women with a history of pregnancy-related cholestasis. Women with a history of COC-related cholestasis may have the condition recur with subsequent COC use.

5.5 High Blood Pressure

For women with well-controlled hypertension, monitor blood pressure and stop Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if blood pressure rises significantly. Women with uncontrolled hypertension or hypertension with vascular disease should not use COCs.

An increase in blood pressure has been reported in women taking COCs, and this increase is more likely in older women and with extended duration of use. The incidence of hypertension increases with increasing concentration of progestin.

5.6 Gallbladder Disease

Studies suggest a small increased relative risk of developing gallbladder disease among COC users.

5.7 Carbohydrate and Lipid Metabolic Effects

Carefully monitor prediabetic and diabetic women who are taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg. COCs may decrease glucose intolerance in a dose-related fashion.

Consider alternative contraception for women with uncontrolled dyslipidemias. A small proportion of women will have adverse lipid changes while on COC's.

Women with hypertriglyceridemia, or a family history thereof, may be at an increased risk of pancreatitis when using COCs.

5.8 Headache

If a woman taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg develops new headaches that are recurrent, persistent, or severe, evaluate the cause and discontinue Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if indicated.

An increase in frequency or severity of migraine during COC use (which may be prodromal of a cerebrovascular event) may be a reason for immediate discontinuation of the COC.

5.9 Bleeding Irregularities

Unscheduled (breakthrough or intracyclic) bleeding and spotting sometimes occur in patients on COCs, especially during the first three months of use. If bleeding persists or occurs after previously regular cycles, check for causes such as pregnancy or malignancy. If pathology and pregnancy are excluded, bleeding irregularities may resolve over time or with a change to a different COC.

Based on patient diaries from two contraceptive clinical trials of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg, 8 to 25% of women experienced unscheduled bleeding per 28-day cycle. A total of 12 subjects out of 1,056 (1.1%) discontinued due to menstrual disorders including intermenstrual bleeding, menorrhagia, and metrorrhagia.

Women who use Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg may experience absence of withdrawal bleeding, even if they are not pregnant. Based on subject diaries from contraception trials for up to 13 cycles, 6 to 10% of women experienced cycles with no withdrawal bleeding. Some women may encounter post-pill amenorrhea or oligomenorrhea, especially when such a condition was pre-existent.

If withdrawal bleeding does not occur, consider the possibility of pregnancy. If the patient has not adhered to the prescribed dosing schedule (missed one or more active tablets or started taking them on a day later than she should have), consider the possibility of pregnancy at the time of the first missed period and take appropriate diagnostic measures. If the patient has adhered to the prescribed regimen and misses two consecutive periods, rule out pregnancy.

5.10 COC Use Before or During Early Pregnancy

Extensive epidemiological studies have revealed no increased risk of birth defects in women who have used oral contraceptives prior to pregnancy. Studies also do not suggest a teratogenic effect, particularly in so far as cardiac anomalies and limb-reduction defects are concerned, when taken inadvertently during early pregnancy.

The administration of oral contraceptives to induce withdrawal bleeding should not be used as a test for pregnancy [*see Use in Specific Populations (8.1)*].

5.11 Depression

Women with a history of depression should be carefully observed and Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg discontinued if depression recurs to a serious degree.

5.12 Interference with Laboratory Tests

The use of COCs may change the results of some laboratory tests, such as coagulation factors, lipids, glucose tolerance, and binding proteins. Women on thyroid hormone replacement therapy may need increased doses of thyroid hormone because serum concentrations of thyroid-binding globulin increase with use of COCs. [*see Drug Interactions (7.2)*].

DRSP causes an increase in plasma renin activity and plasma aldosterone induced by its mild antimineralocorticoid activity.

5.13 Monitoring

A woman who is taking COCs should have a yearly visit with her healthcare provider for a blood pressure check and for other indicated healthcare.

5.14 Other Conditions

In women with hereditary angioedema, exogenous estrogens may induce or exacerbate symptoms of angioedema. Chloasma may occasionally occur, especially in women with a history of chloasma gravidarum. Women with a tendency to chloasma should avoid exposure to the sun or ultraviolet radiation while taking COCs.

6 ADVERSE REACTIONS

The following serious adverse reactions with the use of COCs are discussed elsewhere in the labeling:

- Serious cardiovascular events and stroke [*see Boxed Warning and Warnings and Precautions (5.1)*]
- Vascular events [*see Warnings and Precautions (5.1)*]
- Liver disease [*see Warnings and Precautions (5.4)*]

Adverse reactions commonly reported by COC users are:

- Irregular uterine bleeding
- Nausea
- Breast tenderness
- Headache

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Contraception and Acne Clinical Trials

The data provided reflect the experience with the use of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg in the adequate and well-controlled studies for contraception (N=1,056) and for moderate acne vulgaris (N=536).

For contraception, a Phase 3, multicenter, multinational, open-label study was conducted to evaluate safety and efficacy up to one year in 1,027 women aged 17 - 36 who took at least one dose of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg. A second Phase 3 study was a single center, open-label, active-controlled study to evaluate the effect of 7 28-day cycles of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg on carbohydrate metabolism, lipids and hemostasis in 29 women aged 18-35. For acne, two multicenter, double-blind, randomized, placebo-controlled studies, in 536 women aged 14-45 with moderate acne vulgaris who took at least one dose of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg, evaluated the safety and efficacy during up to 6 cycles.

The adverse reactions seen across the 2 indications overlapped, and are reported using the frequencies from the pooled dataset. The most common adverse reactions ($\geq 2\%$ of users) were: headache/migraine (6.7%), menstrual irregularities (including vaginal hemorrhage [primarily spotting] and metrorrhagia (4.7%), nausea/vomiting (4.2%), breast pain/tenderness (4%) and mood changes (mood swings, depression, depressed mood and affect lability) (2.2%).

Adverse Reactions (≥1%) Leading to Study Discontinuation:

Contraception Clinical Trials

Of 1,056 women, 6.6% discontinued from the clinical trials due to an adverse reaction; the most frequent adverse reactions leading to discontinuation were headache/migraine (1.6%) and nausea/vomiting (1.0%).

Acne Clinical Trials

Of 536 women, 5.4% discontinued from the clinical trials due to an adverse reaction; the most frequent adverse reaction leading to discontinuation was menstrual irregularities (including menometrorrhagia, menorrhagia, metrorrhagia and vaginal hemorrhage) (2.2%).

Serious Adverse Reactions

Contraception Clinical Trials: migraine and cervical dysplasia

Acne Clinical Trials: none reported in the clinical trials

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Adverse reactions are grouped into System Organ Classes, and ordered by frequency.

Vascular disorders: Venous and arterial thromboembolic events (including pulmonary emboli, deep vein thrombosis, cerebral thrombosis, retinal thrombosis, myocardial infarction and stroke), hypertension (including hypertensive crisis)

Hepatobiliary disorders: Gallbladder disease, liver function disturbances, liver tumors

Immune system disorders: Hypersensitivity (including anaphylactic reaction)

Metabolism and nutrition disorders: Hyperkalemia, hypertriglyceridemia, changes in glucose tolerance or effect on peripheral insulin resistance (including diabetes mellitus)

Skin and subcutaneous tissue disorders: Chloasma, angioedema, erythema nodosum, erythema multiforme

Gastrointestinal disorders: Inflammatory bowel disease

Musculoskeletal and connective tissue disorders: Systemic lupus erythematosus

7 DRUG INTERACTIONS

Consult the labeling of all concurrently-used drugs to obtain further information about interactions with hormonal contraceptives or the potential for enzyme alterations.

7.1 Effects of Other Drugs on Combined Oral Contraceptives

Substances diminishing the efficacy of COCs: Drugs or herbal products that induce certain enzymes, including cytochrome P450 3A4 (CYP3A4), may decrease the effectiveness of COCs or increase breakthrough bleeding. Some drugs or herbal products that may decrease the effectiveness of hormonal contraceptives include phenytoin, barbiturates, carbamazepine, bosentan, felbamate, griseofulvin, oxcarbazepine, rifampicin, topiramate and products containing St. John's wort. Interactions between oral contraceptives and other drugs may lead to breakthrough bleeding and/or contraceptive failure. Counsel women to use an alternative method of contraception or a back-up method when enzyme inducers are used with COCs, and to continue back-up contraception for 28 days after discontinuing the enzyme inducer to ensure contraceptive reliability.

Substances increasing the plasma concentrations of COCs: Co-administration of atorvastatin and certain COCs containing EE increase AUC values for EE by approximately 20%. Ascorbic acid and acetaminophen may increase plasma EE concentrations, possibly by inhibition of conjugation. CYP3A4 inhibitors such as itraconazole or ketoconazole may increase plasma hormone concentrations.

Concomitant administration of moderate or strong CYP3A4 inhibitors such as azole antifungals (e.g., ketoconazole, itraconazole, voriconazole, fluconazole), verapamil, macrolides (e.g., clarithromycin, erythromycin), diltiazem, and grapefruit juice can increase the plasma concentrations of the estrogen or the progestin or both. In a clinical drug-drug interaction study conducted in premenopausal women, once daily co-administration of DRSP 3 mg/EE 0.02 mg containing tablets with strong CYP3A4 inhibitor, ketoconazole 200 mg twice daily for 10 days resulted in a moderate increase of DRSP systemic exposure. The exposure of EE was increased mildly [see *Warnings and Precautions (5.2) and Clinical Pharmacology (12.3)*].

Human immunodeficiency virus (HIV)/ Hepatitis C virus (HCV) protease inhibitors and non-nucleoside reverse transcriptase inhibitors: Significant changes (increase or decrease) in the plasma concentrations of estrogen and progestin have been noted in some cases of co-administration with HIV/HCV protease inhibitors or with non-nucleoside reverse transcriptase inhibitors.

Antibiotics: There have been reports of pregnancy while taking hormonal contraceptives and antibiotics, but clinical pharmacokinetic studies have not shown consistent effects of antibiotics on plasma concentrations of synthetic steroids.

7.2 Effects of Combined Oral Contraceptives on Other Drugs

COCs containing EE may inhibit the metabolism of other compounds. COCs have been shown to significantly decrease plasma concentrations of lamotrigine, likely due to induction of lamotrigine glucuronidation. This may reduce seizure control; therefore, dosage adjustments of lamotrigine may be necessary. Consult the labeling of the concurrently-used drug to obtain further information about interactions with COCs or the potential for enzyme alterations.

COCs Increasing the Plasma Concentrations of CYP450 Enzymes: In clinical studies, administration of a hormonal contraceptive containing EE did not lead to any increase or only to a weak increase in plasma concentrations of CYP3A4 substrates (e.g., midazolam) while plasma concentrations of CYP2C19 substrates (e.g., omeprazole and voriconazole) and CYP1A2 substrates (e.g., theophylline and tizanidine) can have a weak or moderate increase.

Clinical studies did not indicate an inhibitory potential of DRSP towards human CYP enzymes at clinically relevant concentrations [see *Clinical Pharmacology (12.3)*].

Women on thyroid hormone replacement therapy may need increased doses of thyroid hormone because serum concentration of thyroid-binding globulin increases with use of COCs.

Potential to Increase Serum Potassium Concentration: There is a potential for an increase in serum potassium concentration in women taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg with other drugs that may increase serum potassium concentration [see *Warnings and Precautions (5.2) and Clinical Pharmacology (12.3)*].

7.3 Interference with Laboratory Tests

The use of contraceptive steroids may influence the results of certain laboratory tests, such as coagulation factors, lipids, glucose tolerance, and binding proteins. DRSP causes an increase in plasma renin activity and plasma aldosterone induced by its mild anti-mineralocorticoid activity. [See *Warnings and Precautions (5.12) and Drug Interactions (7.2)*]

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

There is little or no increased risk of birth defects in women who inadvertently use COCs during early pregnancy. Epidemiologic studies and meta-analyses have not found an increased risk of genital or non-genital birth defects (including cardiac anomalies and limb-reduction defects) following exposure to low dose COCs prior to conception or during early pregnancy.

The administration of COCs to induce withdrawal bleeding should not be used as a test for pregnancy. COCs should not be used during pregnancy to treat threatened or habitual abortion.

Women who do not breastfeed may start COCs no earlier than four weeks postpartum.

8.3 Nursing Mothers

When possible, advise the nursing mother to use other forms of contraception until she has weaned her child. Estrogen-containing COCs can reduce milk production in breastfeeding mothers. This is less likely to occur once breastfeeding is well-established; however, it can occur at any time in some women. Small amounts of oral contraceptive steroids and/or metabolites are present in breast milk.

After oral administration of 3 mg DRSP/0.03 mg EE (Yasmin) tablets, about 0.02% of the DRSP dose was excreted into the breast milk of postpartum women within 24 hours. This results in a maximal daily dose of about 0.003 mg DRSP in an infant.

8.4 Pediatric Use

Safety and efficacy of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg has been established in women of reproductive age. Efficacy is expected to be the same for postpubertal adolescents under the age of 18 and for users 18 years and older. Use of this product before menarche is not indicated.

8.5 Geriatric Use

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg have not been studied in postmenopausal women and is not indicated in this population.

8.6 Patients with Renal Impairment

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are contraindicated in patients with renal impairment [see *Contraindications (4) and Warnings and Precautions (5.2)*].

In subjects with creatinine clearance (CL_{cr}) of 50–79 mL/min, serum DRSP levels were comparable to those in a control group with CL_{cr} ≥ 80 mL/min. In subjects with serum DRSP concentrations were on average 37% higher than those in the control group. In addition, there is a potential to develop hyperkalemia in subjects with renal impairment whose serum potassium is in the upper reference range, and who are concomitantly using potassium sparing drugs [see *Clinical Pharmacology (12.3)*].

8.7 Patients with Hepatic Impairment

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg is contraindicated in patients with hepatic disease [see *Contraindications (4) and Warnings and Precautions (5.4)*]. The mean exposure to DRSP in women with moderate liver impairment is approximately three times higher than the exposure in women with normal liver function. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg has not been studied in women with severe hepatic impairment.

8.8 Race

No clinically significant difference was observed between the pharmacokinetics of DRSP or EE in Japanese versus Caucasian women [see *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

There have been no reports of serious ill effects from overdose, including ingestion by children. Overdosage may cause withdrawal bleeding in females and nausea.

DRSP is a spironolactone analogue which has anti-mineralocorticoid properties. Serum concentration of potassium and sodium, and evidence of metabolic acidosis, should be monitored in cases of overdose.

11 DESCRIPTION


Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg provide an oral contraceptive regimen consisting of 24 pink active film-coated tablets each containing 3 mg of drospirenone, USP (micronized) and 0.02 mg of ethinyl estradiol, USP (micronized) and 4 white inert film coated tablets.

The inactive ingredients in the pink tablets are lactose monohydrate NF, corn starch NF, magnesium stearate NF, povidone USP, Hypromellose, USP, Titanium Dioxide, USP, polyethylene glycol 400 and Ferric oxide red. The white inert film-coated tablets contain lactose monohydrate NF, corn starch NF, povidone USP, magnesium stearate NF, polyvinyl alcohol, titanium dioxide, USP, talc, USP, and polyethylene glycol 4000 .


Drospirenone (6R,7R,8R,9S,10R,13S,14S,15S,16S,17S)-1,3',4',6,6a,7,8,9,10,11,12,13,14,15,15a,16-hexadecahydro-10,13-dimethylspiro-[17H-dicyclopropa-[6,7:15,16] cyclopenta [a]phenanthrene-17,2'(5H)-furan]-3,5'(2H)-dione) is a synthetic progestational compound and has a molecular weight of 366.5 and a molecular formula of $C_{24}H_{30}O_3$.

Ethinyl estradiol (19-nor-17 α -pregna 1, 3, 5(10)-triene-20-yne-3, 17-diol) is a synthetic estrogenic compound and has a molecular weight of 296.4 and a molecular formula of $C_{20}H_{24}O_2$.

The structural formulas are as follows:



Drospirenone



Ethinyl estradiol

"USP Dissolution Test Pending"

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

COCs lower the risk of becoming pregnant primarily by suppressing ovulation. Other possible mechanisms may include cervical mucus changes that inhibit sperm penetration and the endometrial changes that reduce the likelihood of implantation.

12.2 Pharmacodynamics

Drospirenone is a spironolactone analogue with anti-mineralocorticoid and antiandrogenic activity. The

estrogen in Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg is ethinyl estradiol.

Contraception

Two studies evaluated the effect of 3 mg DRSP / 0.02 mg EE combinations on the suppression of ovarian activity as assessed by measurement of follicle size via transvaginal ultrasound and serum hormone (progesterone and estradiol) analyses during two treatment cycles (21-day active tablet period plus 7-day pill-free period). More than 90% of subjects in these studies demonstrated ovulation inhibition. One study compared the effect of 3 mg DRSP/0.02 mg EE combinations with two different regimens (24-day active tablet period plus 4-day pill-free period vs. 21-day active tablet period plus 7-day pill-free period) on the suppression of ovarian activity during two treatment cycles. During the first treatment cycle, there were no subjects (0/49, 0%) taking the 24-day regimen who ovulated compared to 1 subject (1/50, 2%) using the 21-day regimen. After intentionally introduced dosing errors (3 missed active tablets on Days 1 to 3) during the second treatment cycle, there was 1 subject (1/49, 2%) taking the 24-day regimen who ovulated compared to 4 subjects (4/50, 8%) using the 21-day regimen.

Acne

Acne vulgaris is a skin condition with a multifactorial etiology including androgen stimulation of sebum production. While the combination of EE and DRSP increases sex hormone binding globulin (SHBG) and decreases free testosterone, the relationship between these changes and a decrease in the severity of facial acne in otherwise healthy women with this skin condition has not been established. The impact of the antiandrogenic activity of DRSP on acne is not known.

12.3 Pharmacokinetics

Absorption

The absolute bioavailability of DRSP from a single entity tablet is about 76%. The absolute bioavailability of EE is approximately 40% as a result of presystemic conjugation and first-pass metabolism. The absolute bioavailability of Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg, which are a combination tablet of DRSP and EE stabilized by betadex as a clathrate (molecular inclusion complex), has not been evaluated. The bioavailability of EE is similar when dosed via a betadex clathrate formulation compared to when it is dosed as a free steroid. Serum concentrations of DRSP and EE reached peak levels within 1 to 2 hours after administration of Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg.

The pharmacokinetics of DRSP are dose proportional following single doses ranging from 1 to 10 mg. Following daily dosing of Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg, steady state DRSP concentrations were observed after 8 days. There was about 2 to 3 fold accumulation in serum C_{max} and AUC (0–24h) values of DRSP following multiple dose administration of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg (see Table 2).

For EE, steady-state conditions are reported during the second half of a treatment cycle. Following daily administration of Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg, serum C_{max} and AUC (0–24h) values of EE accumulate by a factor of about 1.5 to 2 (see Table 2).

Table 2: Table of Pharmacokinetic Parameters of Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg (DRSP 3 mg and EE 0.02 mg)

DRSP					
Cycle / Day	No. of Subjects	C_{max}^a (ng/mL)	T_{max}^b (h)	AUC _(0-24h) ^a (ng•h/mL)	$t_{1/2}^a$ (h)
1/1	23	38.4 (25)	1.5 (1 - 2)	268 (19)	NA ^c
1/21	23	70.3 (15)	1.5 (1 - 2)	763 (17)	30.8 (22)
EE					
Cycle /	No. of	C_{max}^a	T_{max}^b	AUC _(0-24h) ^a	$t_{1/2}^a$

Day	Subjects	(pg/mL)	(h)	(pg·h/mL)	(h)
1/1	23	32.8 (45)	1.5 (1 - 2)	108 (52)	NA ^c
1/21	23	45.1 (35)	1.5 (1 - 2)	220 (57)	NA ^c

a) geometric mean (geometric coefficient of variation)

b) median (range)

c) NA = Not available

Food Effect

The rate of absorption of DRSP and EE following single administration of a formulation similar to Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg was slower under fed (high fat meal) conditions with the serum C_{max} being reduced about 40% for both components. The extent of absorption of DRSP, however, remained unchanged. In contrast, the extent of absorption of EE was reduced by about 20% under fed conditions.

Distribution

DRSP and EE serum concentrations decline in two phases. The apparent volume of distribution of DRSP is approximately 4 L/kg and that of EE is reported to be approximately 4-5 L/kg.

DRSP does not bind to SHBG or corticosteroid binding globulin (CBG) but binds about 97% to other serum proteins. Multiple dosing over 3 cycles resulted in no change in the free fraction (as measured at trough concentrations). EE is reported to be highly but non-specifically bound to serum albumin (approximately 98.5 %) and induces an increase in the serum concentrations of both SHBG and CBG. EE induced effects on SHBG and CBG were not affected by variation of the DRSP dosage in the range of 2 to 3 mg.

Metabolism

The two main metabolites of DRSP found in human plasma were identified to be the acid form of DRSP generated by opening of the lactone ring and the 4,5-dihydrodrospirenone-3-sulfate. These metabolites were shown not to be pharmacologically active. In *in vitro* studies with human liver microsomes, DRSP was metabolized only to a minor extent mainly by CYP3A4.

EE has been reported to be subject to presystemic conjugation in both small bowel mucosa and the liver. Metabolism occurs primarily by aromatic hydroxylation but a wide variety of hydroxylated and methylated metabolites are formed. These are present as free metabolites and as conjugates with glucuronide and sulfate. CYP3A4 in the liver is responsible for the 2-hydroxylation which is the major oxidative reaction. The 2-hydroxy metabolite is further transformed by methylation and glucuronidation prior to urinary and fecal excretion.

Excretion

DRSP serum concentrations are characterized by a terminal disposition phase half-life of approximately 30 hours after both single and multiple dose regimens. Excretion of DRSP was nearly complete after ten days and amounts excreted were slightly higher in feces compared to urine. DRSP was extensively metabolized and only trace amounts of unchanged DRSP were excreted in urine and feces. At least 20 different metabolites were observed in urine and feces. About 38–47% of the metabolites in urine were glucuronide and sulfate conjugates. In feces, about 17–20% of the metabolites were excreted as glucuronides and sulfates.

For EE the terminal disposition phase half-life has been reported to be approximately 24 hours. EE is not excreted unchanged. EE is excreted in the urine and feces as glucuronide and sulfate conjugates and undergoes enterohepatic circulation.

Use in Specific Populations

Pediatric Use: Safety and efficacy of Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg has been established in women of reproductive age. Efficacy is expected to be the same for postpubertal

adolescents under the age of 18 and for users 18 years and older. Use of this product before menarche is not indicated.

Geriatric Use: Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg have not been studied in postmenopausal women and is not indicated in this population.

Race: No clinically significant difference was observed between the pharmacokinetics of DRSP or EE in Japanese versus Caucasian women (age 25–35) when 3 mg DRSP/0.02 mg EE was administered daily for 21 days. Other ethnic groups have not been specifically studied.

Renal Impairment: Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg are contraindicated in patients with renal impairment.

The effect of renal impairment on the pharmacokinetics of DRSP (3 mg daily for 14 days) and the effect of DRSP on serum potassium concentrations were investigated in three separate groups of female subjects (n=28, age 30–65). All subjects were on a low potassium diet. During the study, 7 subjects continued the use of potassium-sparing drugs for the treatment of their underlying illness. On the 14th day (steady-state) of DRSP treatment, the serum DRSP concentrations in the group with CLcr of 50–79 mL/min were comparable to those in the control group with CLcr \geq 80 mL/min. The serum DRSP concentrations were on average 37% higher in the group with CLcr of 30–49 mL/min compared to those in the control group. DRSP treatment did not show any clinically significant effect on serum potassium concentration. Although hyperkalemia was not observed in the study, in five of the seven subjects who continued use of potassium-sparing drugs during the study, mean serum potassium concentrations increased by up to 0.33 mEq/L. [See *Contraindications (4) and Warnings and Precautions (5.2).*]

Hepatic Impairment: Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg are contraindicated in patients with hepatic disease.

The mean exposure to DRSP in women with moderate liver impairment is approximately three times higher than the exposure in women with normal liver function. Drospirenone and Ethinyl Estradiol Tablets, USP 3mg/0.02 mg have not been studied in women with severe hepatic impairment. [See *Contraindications (4) and Warnings and Precautions (5.4).*]

Drug Interactions

Consult the labeling of all concurrently used drugs to obtain further information about interactions with oral contraceptives or the potential for enzyme alterations.

Effects of Other Drugs on Combined Oral Contraceptives

Substances diminishing the efficacy of COCs: Drugs or herbal products that induce certain enzymes, including CYP3A4, may decrease the effectiveness of COCs or increase breakthrough bleeding.

Substances increasing the plasma concentrations of COCs: Co-administration of atorvastatin and certain COCs containing EE increase AUC values for EE by approximately 20%. Ascorbic acid and acetaminophen may increase plasma EE concentrations, possibly by inhibition of conjugation. In a clinical drug-drug interaction study conducted in 20 premenopausal women, co-administration of a DRSP (3 mg)/EE (0.02 mg) COC with the strong CYP3A4 inhibitor ketoconazole (200 mg twice daily) for 10 days increased the AUC(0-24h) of DRSP and EE by 2.68-fold (90% CI: 2.44, 2.95) and 1.40-fold (90% CI: 1.31, 1.49), respectively. The increases in C were 1.97-fold (90% CI: 1.79, 2.17) and 1.39-fold (90% CI: 1.28, 1.52) for DRSP and EE, respectively. Although no clinically relevant effects on safety or laboratory parameters including serum potassium were observed, this study only assessed subjects for 10 days. The clinical impact for a patient taking a DRSP-containing COC concomitantly with chronic use of a CYP3A4/5 inhibitor is unknown [see *Warnings and Precautions (5.2)*].

HIV/HCV protease inhibitors and non-nucleoside reverse transcriptase inhibitors: Significant changes (increase or decrease) in the plasma concentrations of estrogen and progestin have been noted in some cases of co-administration with HIV/HCV protease inhibitors or with non-nucleoside reverse

transcriptase inhibitors.

Antibiotics: There have been reports of pregnancy while taking hormonal contraceptives and antibiotics, but clinical pharmacokinetic studies have not shown consistent effects of antibiotics on plasma concentrations of synthetic steroids.

Effects of Combined Oral Contraceptives on Other Drugs

COCs containing ethinyl estradiol may inhibit the metabolism of other compounds. COCs have been shown to significantly decrease plasma concentrations of lamotrigine, likely due to induction of lamotrigine glucuronidation. This may reduce seizure control; therefore, dosage adjustments of lamotrigine may be necessary. Consult the labeling of the concurrently-used drug to obtain further information about interactions with COCs or the potential for enzyme alterations.

In vitro, EE is a reversible inhibitor of CYP2C19, CYP1A1 and CYP1A2 as well as a mechanism-based inhibitor of CYP3A4/5, CYP2C8, and CYP2J2. Metabolism of DRSP and potential effects of DRSP on hepatic CYP enzymes have been investigated in *in vitro* and *in vivo* studies. In *in vitro* studies DRSP did not affect turnover of model substrates of CYP1A2 and CYP2D6, but had an inhibitory influence on the turnover of model substrates of CYP1A1, CYP2C9, CYP2C19, and CYP3A4, with CYP2C19 being the most sensitive enzyme. The potential effect of DRSP on CYP2C19 activity was investigated in a clinical pharmacokinetic study using omeprazole as a marker substrate. In the study with 24 postmenopausal women [including 12 women with homozygous CYP2C19 genotype] the daily oral administration of 3 mg DRSP for 14 days did not affect the oral clearance of omeprazole (40 mg, single oral dose) and the CYP2C19 product 5-hydroxy omeprazole. Furthermore, no significant effect of DRSP on the systemic clearance of the CYP3A4 product omeprazole sulfone was found. These results demonstrate that DRSP did not inhibit CYP2C19 and CYP3A4 *in vivo*.

Two additional clinical drug-drug interaction studies using simvastatin and midazolam as marker substrates for CYP3A4 were each performed in 24 healthy postmenopausal women. The results of these studies demonstrated that pharmacokinetics of the CYP3A4 substrates were not influenced by steady state DRSP concentrations achieved after administration of 3 mg DRSP/day.

Women on thyroid hormone replacement therapy may need increased doses of thyroid hormone because serum concentration of thyroid-binding globulin increases with use of COCs.

Interactions With Drugs That Have the Potential to Increase Serum Potassium Concentration: There is a potential for an increase in serum potassium concentration in women taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg with other drugs that may increase serum potassium concentration [see Warnings and Precautions (5.2)].

A drug-drug interaction study of DRSP 3 mg/estradiol (E2) 1 mg versus placebo was performed in 24 mildly hypertensive postmenopausal women taking enalapril maleate 10 mg twice daily. Potassium concentrations were obtained every other day for a total of 2 weeks in all subjects. Mean serum potassium concentrations in the DRSP/E2 treatment group relative to baseline were 0.22 mEq/L higher than those in the placebo group. Serum potassium concentrations also were measured at multiple time points over 24 hours at baseline and on Day 14. On Day 14, the ratios for serum potassium C_{max} and AUC in the DRSP/E2 group to those in the placebo group were 0.955 (90% CI: 0.914, 0.999) and 1.010 (90% CI: 0.944, 1.08), respectively. No patient in either treatment group developed hyperkalemia (serum potassium concentrations > 5.5 mEq/L).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 24 month oral carcinogenicity study in mice dosed with 10 mg/kg/day DRSP alone or 1 + 0.01, 3 + 0.03 and 10 + 0.1 mg/kg/day of DRSP and EE, 0.1 to 2 times the exposure (AUC of DRSP) of women taking a contraceptive dose, there was an increase in carcinomas of the Harderian gland in the group that

received the high dose of DRSP alone. In a similar study in rats given 10 mg/kg/day DRSP alone or 0.3 + 0.003, 3 + 0.03 and 10 + 0.1 mg/kg/day DRSP and EE, 0.8 to 10 times the exposure of women taking a contraceptive dose, there was an increased incidence of benign and total (benign and malignant) adrenal gland pheochromocytomas in the group receiving the high dose of DRSP. Mutagenesis studies for DRSP were conducted *in vivo* and *in vitro* and no evidence of mutagenic activity was observed.

14 CLINICAL STUDIES

14.1 Oral Contraceptive Clinical Trial

In the primary contraceptive efficacy study of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg (3 mg DRSP/0.02 mg EE) of up to 1 year duration, 1,027 subjects were enrolled and completed 11,480 28-day cycles of use. The age range was 17 to 36 years. The racial demographic was: 87.8% Caucasian, 4.6% Hispanic, 4.3% Black, 1.2% Asian, and 2.1% other. Women with a BMI greater than 35 were excluded from the trial. The pregnancy rate (Pearl Index) was 1.41 (95% CI [0.73, 2.47]) per 100 woman-years of use based on 12 pregnancies that occurred after the onset of treatment and within 14 days after the last dose of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg in women 35 years of age or younger during cycles in which no other form of contraception was used.

14.3 Acne Clinical Trials

In two multicenter, double-blind, randomized, placebo-controlled studies, 889 subjects, ages 14 to 45 years, with moderate acne received Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg or placebo for six 28-day cycles. The primary efficacy endpoints were the percent change in inflammatory lesions, non-inflammatory lesions, total lesions, and the percentage of subjects with a "clear" or "almost clear" rating on the Investigator's Static Global Assessment (ISGA) scale on day 15 of cycle 6, as presented in Table 3:

Table 3: Efficacy Results for Acne Trials*

	Study 1		Study 2	
	Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg N=228	Placebo N=230	Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg N=218	Placebo N=213
ISGA Success Rate	35 (15%)	10 (4%)	46 (21%)	19 (9%)
Inflammatory Lesions				
Mean Baseline Count	33	33	32	32
Mean Absolute (%) Reduction	15 (48%)	11 (32%)	16 (51%)	11 (34%)
Non-inflammatory Lesions				
Mean Baseline Count	47	47	44	44
Mean Absolute (%) Reduction	18 (39%)	10 (18%)	17 (42%)	11 (26%)
Total Lesions				
Mean Baseline Count	80	80	76	76
Mean Absolute (%) Reduction	33 (42%)	21 (25%)	33 (46%)	22 (31%)

* Evaluated at day 15 of cycle 6, last observation carried forward for the Intent to treat population

15 REFERENCES

1. Seeger, J.D., Loughlin, J., Eng, P.M., Clifford, C.R., Cutone, J., and Walker, A.M. (2007). Risk of thromboembolism in women taking ethinylestradiol/drospirenone and other oral contraceptives. *Obstet Gynecol* 110, 587-593.
2. Dinger, J.C., Heinemann, L.A., and Kuhl-Habich, D. (2007). The safety of a drospirenone-containing oral contraceptive: final results from the European Active Surveillance Study on oral contraceptives based on 142,475 women-years of observation. *Contraception* 75, 344-354.
3. Combined hormonal contraceptives (CHCs) and the risk of cardiovascular endpoints. Sidney, S. (primary author) <http://www.fda.gov/downloads/Drugs/DrugSafety/UCM277384.pdf>, accessed Oct 27, 2011.
4. Lidegaard, O., Lokkegaard, E., Svendsen, A.L., and Agger, C. (2009). Hormonal contraception and risk of venous thromboembolism: national follow-up study. *BMJ* 339, b2890.
5. Lidegaard, O., Nielsen, L.H., Skovlund, C.W., Skjeldestad, F.E., and Lokkegaard, E. (2011). Risk of venous thromboembolism from use of oral contraceptives containing different progestogens and oestrogen doses: Danish cohort study, 2001-9. *BMJ* 343, d6423.
6. van Hylckama Vlieg, A., Helmerhorst, F.M., Vandenbroucke, J.P., Doggen, C.J., and Rosendaal, F.R. (2009). The venous thrombotic risk of oral contraceptives, effects of oestrogen dose and progestogen type: results of the MEGA case-control study. *BMJ* 339, b2921.
7. Dinger, J., Assmann, A., Mohner, S., and Minh, T.D. (2010). Risk of venous thromboembolism and the use of dienogest- and drospirenone-containing oral contraceptives: results from a German case-control study. *J Fam Plann Reprod Health Care* 36, 123-129.
8. Jick, S.S., and Hernandez, R.K. (2011). Risk of non-fatal venous thromboembolism in women using oral contraceptives containing drospirenone compared with women using oral contraceptives containing levonorgestrel: case-control study using United States claims data. *BMJ* 342, d2151.
9. Parkin, L., Sharples, K., Hernandez, R.K., and Jick, S.S. (2011). Risk of venous thromboembolism in users of oral contraceptives containing drospirenone or levonorgestrel: nested case-control study based on UK General Practice Research Database. *BMJ* 342, d2139.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are available in packages of three blister packs (NDC 61916-090-30).

The film-coated tablets are rounded with "P" debossed on one side.

Each blister pack (28 film-coated tablets) contains in the following order:

- 24 active pink round, unscored, film-coated tablets debossed with a "P" on one side, each containing 3 mg drospirenone, USP (micronized) and 0.02 mg ethinyl estradiol, USP (micronized)
- 4 inert white round, unscored, film-coated tablets.

16.2 Storage

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

See "FDA-approved patient labeling (Patient Information)."

- Counsel patients that cigarette smoking increases the risk of serious cardiovascular events from COC use, and that women who are over 35 years old and smoke should not use COCs.
- Counsel patients that the increased risk of VTE compared to non-users of COCs is greatest after initially starting a COC or restarting (following a 4-week or greater pill-free interval) the same or a

different COC.

- Counsel patients about the information regarding the risk of VTE with DRSP-containing COCs compared to COCs that contain levonorgestrel or some other progestins.
- Counsel patients that Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg do not protect against HIV-infection (AIDS) and other sexually transmitted diseases.
- Counsel patients on Warnings and Precautions associated with COCs.
- Counsel patients that Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg contain DRSP. Drospirenone may increase potassium. Patients should be advised to inform their healthcare provider if they have kidney, liver or adrenal disease because the use of Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg in the presence of these conditions could cause serious heart and health problems. They should also inform their healthcare provider if they are currently on daily, long-term treatment (NSAIDs, potassium-sparing diuretics, potassium supplementation, ACE inhibitors, angiotensin-II receptor antagonists, heparin or aldosterone antagonists) for a chronic condition.
- Inform patients that Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are not indicated during pregnancy. If pregnancy occurs during treatment with Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg, instruct the patient to stop further intake.
- Counsel patients to take one tablet daily by mouth at the same time every day. Instruct patients what to do in the event pills are missed. See “**What to Do if You Miss Pills**” section in **FDA-Approved Patient Labeling**.
- Counsel patients to use a back-up or alternative method of contraception when enzyme inducers are used with COCs.
- Counsel patients who are breastfeeding or who desire to breastfeed that COCs may reduce breast milk production. This is less likely to occur if breastfeeding is well established.
- Counsel any patient who starts COCs postpartum, and who has not yet had a period, to use an additional method of contraception until she has taken a pink tablet for 7 consecutive days.
- Counsel patients that amenorrhea may occur. Rule out pregnancy in the event of amenorrhea in two or more consecutive cycles.

Manufactured by:

Pharmaceutics International, Inc. (Pii)

Hunt Valley, MD 21031

USA



FDA Approved Patient Labeling

Guide for Using Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg

WARNING TO WOMEN WHO SMOKE

Do not use Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if you smoke cigarettes and are over 35 years old. Smoking increases your risk of serious cardiovascular side effects (heart and blood vessel problems) from birth control pills, including death from heart attack, blood clots or stroke. This risk increases with age and the number of cigarettes you smoke.

Birth control pills help to lower the chances of becoming pregnant when taken as directed. They do not protect against HIV infection (AIDS) and other sexually transmitted diseases.

What Are Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg?

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are a birth control pill. It contains two female hormones, a synthetic estrogen called ethinyl estradiol and a progestin called drospirenone.

The progestin drospirenone may increase potassium. Therefore, you should not take Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if you have kidney, liver or adrenal disease because this could cause serious heart and health problems. Other drugs may also increase potassium. If you are currently on daily, long-term treatment for a chronic condition with any of the medications below, you should consult your healthcare provider about whether Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg are right for you, and during the first month that you take Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg, you should have a blood test to check your potassium level.

NSAIDs (ibuprofen [Motrin, Advil], naproxen [Aleve and others] when taken long-term and daily for treatment of arthritis or other problems)

Potassium-sparing diuretics (spironolactone and others)

Potassium supplementation

ACE inhibitors (Capoten, Vasotec, Zestril and others)

Angiotensin-II receptor antagonists (Cozaar, Diovan, Avapro and others)

Heparin

Aldosterone antagonists

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg may also be taken to treat moderate acne if all of the following are true:

Your healthcare provider says it is safe for you to use Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg

You are at least 14 years old.

You have started having menstrual periods.


You want to use a birth control pill to prevent pregnancy.

How Well Do Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg Work?

Your chance of getting pregnant depends on how well you follow the directions for taking your birth control pills. The better you follow the directions, the less chance you have of getting pregnant.

Based on the results of one clinical study, 1 to 2 women out of 100 women, may get pregnant during the first year they use Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg.

The following chart shows the chance of getting pregnant for women who use different methods of birth control. Each box on the chart contains a list of birth control methods that are similar in effectiveness. The most effective methods are at the top of the chart. The box on the bottom of the chart shows the chance of getting pregnant for women who do not use birth control and are trying to get pregnant.



How Do I Take Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg?

1. **Be sure to read these directions** before you start taking your pills or anytime you are not sure what to do.
2. The right way to take the pill is to take one pill every day at the same time in the order directed on the package. Preferably, take the pill after the evening meal or at bedtime, with some liquid, as needed. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg can be taken without regard to meals.
If you miss pills you could get pregnant. This includes starting the pack late. The more pills you miss, the more likely you are to get pregnant. See "WHAT TO DO IF YOU MISS PILLS" below.
3. Many women have spotting or light bleeding at unexpected times, or may feel sick to their stomach during the first 1-3 packs of pills.
If you do have spotting or light bleeding or feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If it does not go away, check with your healthcare provider.
4. Missing pills can also cause spotting or light bleeding, even when you make up these missed pills. On the days you take two pills, to make up for missed pills, you could also feel a little sick to your stomach.
5. If you have vomiting (within 3 to 4 hours after you take your pill), you should follow the instructions for "WHAT TO DO IF YOU MISS PILLS." If you have diarrhea or if you take certain medicines, including some antibiotics and some herbal products such as St. John's Wort, your pills may not work as well.
Use a back-up method (such as condoms and spermicides) until you check with your healthcare provider.

- If you have trouble remembering to take the pill, talk to your healthcare provider about how to make pill-taking easier or about using another method of birth control.
- If you have any questions or are unsure about the information in this leaflet, call your healthcare provider.**

Before You Start Taking Your Pills

- Decide What Time of Day You Want to Take Your Pill

It is important to take Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg in the order directed on the package at the same time every day, preferably after the evening meal or at bedtime, with some liquid, as needed. Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg can be taken without regard to meals.

- Look at Your Pill Pack - It has 28 Pills

The Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg-pill pack has 24 pink pills (with hormones) to be taken for 24 days, followed by 4 white pills (without hormones) to be taken for the next four days.

- Also look for:

- Where on the pack to start taking pills,
- In what order to take the pills (follow the arrows)



- Be sure you have ready at all times (a) another kind of birth control (such as condoms and spermicides) to use as a back-up in case you miss pills, and (b) an extra, full pill pack.

When To Start the First Pack of Pills

You have a choice for which day to start taking your first pack of pills. Decide with your healthcare provider which is the best day for you. Pick a time of day which will be easy to remember.

Day 1 Start:

- Take the first pink pill of the pack during the first 24 hours of your period.
- You will not need to use a back-up method of birth control, since you are starting the Pill at the beginning of your period. However, if you start Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg later than the first day of your period, you should use another method of birth control

(such as a condom and spermicide) as a back-up method until you have taken 7 pink pills.

Sunday Start:

1. Take the first pink pill of the pack on the Sunday after your period starts, even if you are still bleeding. If your period begins on Sunday, start the pack that same day.
2. Use another method of birth control (such as a condom and spermicide) as a back-up method if you have sex anytime from the Sunday you start your first pack until the next Sunday (7 days).

This also applies if you start Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg after having been pregnant, and you have not had a period since your pregnancy.

When You Switch From a Different Birth Control Pill

When switching from another birth control pill, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started on the same day that a new pack of the previous birth control pill would have been started.

When You Switch From Another Type of Birth Control Method

When switching from a transdermal patch or vaginal ring, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started when the next application would have been due. When switching from an injection, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started when the next dose would have been due. When switching from an intrauterine contraceptive or an implant, Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg should be started on the day of removal.

What to Do During the Month

1. Take one pill at the same time every day until the pack is empty.
Do not skip pills even if you are spotting or bleeding between monthly periods or feel sick to your stomach (nausea).
Do not skip pills even if you do not have sex very often.
2. When you finish a pack of pills, start the next pack on the day after your last white pill. Do not wait any days between packs.

What to Do if You Miss Pills

If you miss 1 pink pill of your pack:

1. Take it as soon as you remember. Take the next pill at your regular time. This means you may take two pills in one day.
2. You do not need to use a back-up birth control method if you have sex.

If you miss 2 pink pills in a row in Week 1 or Week 2 of your pack:

1. Take two pills on the day you remember and two pills the next day.
2. Then take one pill a day until you finish the pack.
3. You could become pregnant if you have sex in the 7 days after you restart your pills. You must use another birth control method (such as a condom and spermicide) as a back-up for those 7 days.

If you miss 2 pink pills in a row in Week 3 or Week 4 of your pack:

1. If you are a Day 1 Starter:

Throw out the rest of the pill pack and start a new pack that same day.

If you are a Sunday Starter:

Keep taking one pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack of pills that same day.

2. You could become pregnant if you have sex in the 7 days after you restart your pills. You must use

- another birth control method (such as a condom and spermicide) as a back-up for those 7 days.
3. You may not have your period this month but this is expected. However, if you miss your period two months in a row, call your healthcare provider because you might be pregnant.

If you miss 3 or more pink pills in a row during any week:

1. If you are a Day 1 Starter:

Throw out the rest of the pill pack and start a new pack that same day.

If you are a Sunday Starter:

Keep taking 1 pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack of pills that same day.

2. You could become pregnant if you have sex in the 7 days after you restart your pills. You must use another birth control method (such as condoms and spermicides) as a back-up for those 7 days.
3. Call your healthcare provider if you miss your period, because you might be pregnant.

If you miss any of the 4 white pills in Week 4:

Throw away the pills you missed.

Keep taking one pill each day until the pack is empty.

You do not need a back-up method.

Finally, if you are still not sure what to do about the pills you have missed:

Use a back-up method (such as condoms and spermicides) anytime you have sex.

Contact your healthcare provider and continue taking one active pink pill each day until otherwise directed.

WHO SHOULD NOT TAKE Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg?

Your healthcare provider will not give you Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if you:

- Ever had blood clots in your legs (deep vein thrombosis), lungs (pulmonary embolism), or eyes (retinal thrombosis)
- Ever had a stroke
- Ever had a heart attack
- Have certain heart valve problems or heart rhythm abnormalities that can cause blood clots to form in the heart
- Have an inherited problem with your blood that makes it clot more than normal
- Have high blood pressure that medicine can't control
- Have diabetes with kidney, eye, nerve, or blood vessel damage
- Ever had certain kinds of severe migraine headaches with aura, numbness, weakness or changes in vision
- Ever had breast cancer or any cancer that is sensitive to female hormones
- Have liver disease, including liver tumors
- Have kidney disease
- Have adrenal disease

Also, do not take birth control pills if you:

- Smoke and are over 35 years old
- Are or suspect you are pregnant

Birth control pills may not be a good choice for you if you have ever had jaundice (yellowing of the skin or eyes) caused by pregnancy (also called cholestasis of pregnancy).

Tell your healthcare provider if you have ever had any of the above conditions (your healthcare provider can recommend another method of birth control).

What Else Should I Know about Taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg?

Birth control pills do not protect you against any sexually transmitted disease, including HIV, the virus that causes AIDS.

Do not skip any pills, even if you do not have sex often.

If you miss a period, you could be pregnant. However, some women miss periods or have light periods on birth control pills, even when they are not pregnant. Contact your healthcare provider for advice if you:

- Think you are pregnant.
- Miss one period and have not taken your birth control pills every day
- Miss two periods in a row

Birth control pills should not be taken during pregnancy. However, birth control pills taken by accident during pregnancy are not known to cause birth defects.

You should stop Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg at least four weeks before you have major surgery and not restart it until at least two weeks after the surgery due to an increased risk of blood clots.

If you are breastfeeding, consider another birth control method until you are ready to stop breastfeeding. Birth control pills that contain estrogen, like Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg, may decrease the amount of milk you make. A small amount of the pill's hormones pass into breast milk.

If you have vomiting or diarrhea, your birth control pills may not work as well. Use another birth control method, like condoms and a spermicide, until you check with your healthcare provider.

If you are scheduled for any laboratory tests, tell your doctor you are taking birth-control pills. Certain blood tests may be affected by birth-control pills.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg may affect the way other medicines work, and other medicines may affect how well Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg works. Know the medicines you take.

Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

What are the Most Serious Risks of Taking Birth Control Pills?

Like pregnancy, birth control pills increase the risk of serious blood clots (see following graph), especially in women who have other risk factors, such as smoking, obesity, or age greater than 35. This increased risk is highest when you first start taking birth control pills and when you restart the same or different birth control pills after not using them for a month or more. Women who use birth control pills with drospirenone (like Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg) may have a higher risk of getting a blood clot. Some studies reported that the risk of blood clots was higher for women who use birth control pills that contain drospirenone than for women who use birth control pills that do not contain drospirenone.


Talk with your healthcare provider about your risk of getting a blood clot before deciding which birth control pill is right for you.

It is possible to die or be permanently disabled from a problem caused by a blood clot, such as a heart attack or a stroke. Some examples of serious clots are blood clots in the:

- Legs (deep vein thrombosis or DVT)
- Lungs (pulmonary embolus or PE)
- Eyes (loss of eyesight)
- Heart (heart attack)
- Brain (stroke)

To put the risk of developing a blood clot into perspective: If 10,000 women who are not pregnant and do not use birth control pills are followed for one year, between 1 and 5 of these women will develop a blood clot. The figure below shows the likelihood of developing a serious blood clot for women who are not pregnant and do not use birth control pills, for women who use birth control pills, for pregnant women, and for women in the first 12 weeks after delivering a baby.

Likelihood of Developing a Serious Blood Clot



* Pregnancy data based on actual duration of pregnancy in the reference studies. Based on a model assumption that pregnancy duration is nine months, the rate is 7 to 27 per 10,000 WY.

A few women who take birth control pills may get:

- High blood pressure
- Gallbladder problems
- Rare cancerous or noncancerous liver tumors

All of these events are uncommon in healthy women.

Call your healthcare provider right away if you have:

- Persistent leg pain
- Sudden shortness of breath
- Sudden blindness, partial or complete
- Severe pain in your chest

- Sudden, severe headache unlike your usual headaches
- Weakness or numbness in an arm or leg, or trouble speaking
- Yellowing of the skin or eyeballs

What are the Common Side Effects of Birth Control Pills?

The most common side effects of birth control pills are:

- Spotting or bleeding between menstrual periods
- Nausea
- Breast tenderness
- Headache

These side effects are usually mild and usually disappear with time.

Less common side effects are:

- Acne
- Less sexual desire
- Bloating or fluid retention
- Blotchy darkening of the skin, especially on the face
- High blood sugar, especially in women who already have diabetes
- High fat (cholesterol; triglyceride) levels in the blood
- Depression, especially if you have had depression in the past. Call your healthcare provider immediately if you have any thoughts of harming yourself.
- Problems tolerating contact lenses
- Weight changes

This is not a complete list of possible side effects. Talk to your healthcare provider if you develop any side effects that concern you. You may report side effects to the FDA at 1-800-FDA-1088.

No serious problems have been reported from a birth control pill overdose, even when accidentally taken by children.

Do Birth Control Pills Cause Cancer?

Birth control pills do not seem to cause breast cancer. However, if you have breast cancer now, or have had it in the past, do not use birth control pills because some breast cancers are sensitive to hormones.

Women who use birth control pills may have a slightly higher chance of getting cervical cancer. However, this may be due to other reasons such as having more sexual partners.

What Should I Know about My Period when Taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg?

Irregular vaginal bleeding or spotting may occur while you are taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg. Irregular bleeding may vary from slight staining between menstrual periods to breakthrough bleeding, which is a flow much like a regular period. Irregular bleeding occurs most often during the first few months of oral contraceptive use, but may also occur after you have been taking the pill for some time. Such bleeding may be temporary and usually does not indicate any serious problems. It is important to continue taking your pills on schedule. If the bleeding occurs in more than one cycle, is unusually heavy, or lasts for more than a few days, call your healthcare provider.

Some women may not have a menstrual period but this should not be cause for alarm as long as you have taken the pills according to direction.

What if I Miss My Scheduled Period when Taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg?

It is not uncommon to miss your period. However, if you miss two periods in a row or miss one period when you have not taken your birth control pills according to directions, call your healthcare provider. Also notify your healthcare provider if you have symptoms of pregnancy such as morning sickness or unusual breast tenderness. It is important that your healthcare provider checks you to find out if you are pregnant. Stop taking Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg if you are pregnant.

What If I Want to Become Pregnant?

You may stop taking the pill whenever you wish. Consider a visit with your healthcare provider for a pre-pregnancy checkup before you stop taking the pill.

General Advice about Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg

Your healthcare provider prescribed Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg for you. Please do not share Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg with anyone else. Keep Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02 mg out of the reach of children.

If you have concerns or questions, ask your healthcare provider. You may also ask your healthcare provider for a more detailed label written for medical professionals.

Pharmaceutics International, Inc. (Pii)

Revised March 2017

Principal Display Panel - Drospirenone and Ethinyl Estradiol Tablets Carton Label

NDC 61916-090-30

3 Blister Packs of 28 Tablets


Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02mg

Usual Dosage: One tablet daily for 28 consecutive days per menstrual cycle in the following order: 24 pink tablets followed by four white tablets as prescribed.

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Contains three (3) cartons, each containing one blister pack of 28 tablets (24 pink tablets, containing 3 mg drospirenone and 0.02 mg ethinyl estradiol; and 4 white inert tablets).



Principal Display Panel - Drospirenone and Ethinyl Estradiol Tablets Label

NDC 61916-090-30

Drospirenone and Ethinyl Estradiol Tablets, USP 3 mg/0.02mg

Contents: One blister pack of 28 tablets, 24 pink tablets and 4 white inert tablets.


Each pink tablet (24) contains 3 mg drospirenone and 0.02 mg ethinyl estradiol.

Each white tablet (4) contains inert ingredients.

Rx only

THIS PRODUCT (LIKE ALL ORAL CONTRACEPTIVES) IS INTENDED TO PREVENT PREGNANCY. IT DOES NOT PROTECT AGAINST HIV INFECTION (AIDS) AND OTHER SEXUALLY TRANSMITTED DISEASES.

Pii



DROSPIRENONE AND ETHINYL ESTRADIOL

drospirenone and ethinyl estradiol kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:61916-090
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61916-090-30	3 in 1 CARTON	10/26/2017	
1		1 in 1 BLISTER PACK		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		24
Part 2		4

Part 1 of 2

DROSPIRENONE AND ETHINYL ESTRADIOL

drospirenone and ethinyl estradiol tablet, film coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Drospirenone (UNII: N295J34A25) (Drospirenone - UNII:N295J34A25)	Drospirenone	3 mg
Ethinyl Estradiol (UNII: 423D2T571U) (Ethinyl Estradiol - UNII:423D2T571U)	Ethinyl Estradiol	0.02 mg

Inactive Ingredients

Ingredient Name	Strength
Lactose Monohydrate (UNII: EWQ57Q8I5X)	
Starch, Corn (UNII: O8232NY3SJ)	
Magnesium Stearate (UNII: 70097M6I30)	
Povidone, Unspecified (UNII: FZ989GH94E)	
Hypromellose, Unspecified (UNII: 3NXW29V3WO)	
Titanium Dioxide (UNII: 15FIX9V2JP)	
Polyethylene Glycol 400 (UNII: B697894SGQ)	
Ferric oxide red (UNII: 1K09F3G675)	

Product Characteristics

Color	pink (pink)	Score	no score
Shape	ROUND (ROUND)	Size	6mm
Flavor		Imprint Code	P
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203291	10/26/2017	

Part 2 of 2

INERT

inert tablet, film coated

Product Information

Route of Administration ORAL

Inactive Ingredients

Ingredient Name	Strength
Lactose Monohydrate (UNII: EWQ57Q8I5X)	
Starch, Corn (UNII: O8232NY3SJ)	
Magnesium Stearate (UNII: 70097M6I30)	
Povidone, Unspecified (UNII: FZ989GH94E)	
Polyvinyl Alcohol, Unspecified (UNII: 532B59J990)	
Titanium Dioxide (UNII: 15FIX9V2JP)	
Talc (UNII: 7SEV7J4R1U)	
Polyethylene Glycol 4000 (UNII: 4R4HF16D95)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	6mm
Flavor		Imprint Code	
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203291	10/26/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203291	10/26/2017	

Labeler - Pharmaceutics International, Inc. (Pii) (878265586)

Establishment

Name	Address	ID/FEI	Business Operations
Pharmaceutics International, Inc. (Pii)		878265586	MANUFACTURE(61916-090) , ANALYSIS(61916-090)

Revised: 7/2017

Pharmaceutics International, Inc. (Pii)