

PROLAYED LIDOCAINE MALE GENITAL DESENSITIZER- lidocaine ointment
ViaDerma Distribution, Inc.

Prolayed Lidocaine Ointment-Male Genital Desensitizer

Drug Facts

Active Ingredient

Lidocaine 10 mg

Purpose

Male genital desensitizer

Use

Helps in temporarily slowing the onset of ejaculation.

Warnings

For external use only

When using this product

- avoid contact with the eyes.

Stop use and ask a doctor if

- this product, used as directed, does not provide relief. Premature ejaculation may be due to a condition requiring medical supervision
- you or your partner develop a rash or irritation, such as burning or itching.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply 3 or more sprays, not to exceed 10, to head and shaft of penis before intercourse, or use as directed by a doctor.
- Wash product off after intercourse.

Inactive Ingredients

acetic acid, ascorbic acid, chlorhexidine gluconate, cholecalciferol, dimethyl sulfoxide, dipropylene glycol, glucono delta lactone, glycerin, histidine, hydroxethyl-cellulose, magnesium stearate, methylparaben, sodium hydroxide, sorbic acid, stearic acid, water

Package Labeling:

Drug Facts

Active Ingredient (in each spray) **Purpose**
Lidocaine 10 mg. Male genital desensitizer

Use
Helps in temporarily slowing the onset of ejaculation.

Warnings
For external use only

When using this product
■ avoid contact with the eyes.

Stop use and ask a doctor if
■ this product, used as directed, does not provide relief. Premature ejaculation may be due to a condition requiring medical supervision
■ you or your partner develop a rash or irritation, such as burning or itching.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
■ Apply 3 or more sprays, not to exceed 10, to head and shaft of penis before intercourse, or use as directed by a doctor.
■ Wash product off after intercourse.

Inactive Ingredients
acetic acid, ascorbic acid, chlorhexidine gluconate, cholecalciferol, dimethyl sulfoxide, dipropylene glycol, glucono delta lactone, glycerin, histidine, hydroxyethylcellulose, magnesium stearate, methylparaben, sodium hydroxide, sorbic acid, stearic acid, water

topical spray
male genital desensitizer
PROLAYED
VIA DERMA[®] Distribution

VIA DERMA[®] Distribution
PROLAYED[™]
male genital desensitizer
topical spray
Lidocaine
10mg in 1 gram
(active ingredient)



non-staining formula

Keep this and all drugs away from children. In case of accidental ingestion, seek medical assistance or contact a Poison Control Center immediately.

MADE IN USA
VIA DERMA[®] Distribution
PROLAYED[™]
male genital desensitizer
topical spray
Lidocaine
10mg in 1 gram
(active ingredient)

non-staining formula

0.5 FL OZ (15ml)

VIA DERMA[®] Distribution
PROLAYED[™]
male genital desensitizer
topical spray
Lidocaine
10mg in 1 gram
(active ingredient)



non-staining formula

Side Effect: Same as any other ingredients in Triastem[™] products. See back panel for additional information.

Licensed to ViaDerma Distribution by: ViaDerma, Inc.
Distributed by: ViaDerma Distribution. Report any side effects to: ViaDerma Distribution 2235 E. Flamingo Rd, Suite 152 Las Vegas, NV 89119 Tel: 1-800-585-8685 www.ViaDermaInc.com

PATENT PENDING



8 65393 00040 8
NDC# 71262-002-15

Licensed to ViaDerma Distribution by:
ViaDerma, Inc. Distributed by: ViaDerma
Distribution Report any side effects to:
ViaDerma Distribution 2235 E. Flamingo Rd,
Suite 152 Las Vegas, NV 89119
Tel: 1-800-585-8685 | NDC# 71262-002-01



PROLAYED™

male genital desensitizer

topical spray

Lidocaine
10mg in 1 gram



0.5 FL OZ (15ml)

KEEP THIS AND ALL DRUGS AWAY FROM
CHILDREN. IN CASE OF ACCIDENTAL INGESTION,
SEEK MEDICAL ASSISTANCE OR CONTACT
A POISON CONTROL IMMEDIATELY.

PROLAYED LIDOCAINE MALE GENITAL DESENSITIZER

lidocaine ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71262-006
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SORBIC ACID (UNII: X045VJ989B)	
ACETIC ACID (UNII: Q40Q9N063P)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E)	
CHOLECALCIFEROL (UNII: 1C6V77QF41)	
DIMETHYL SULFOXIDE (UNII: YOW8V9698H)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
GLUCONOLACTONE (UNII: WQ29KQ9POT)	
GLYCERIN (UNII: PDC6A3C0OX)	
HISTIDINE (UNII: 4QD397987E)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71262-006-15	1 in 1 BOX	06/23/2017	
1		15 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	06/23/2017	

Labeler - ViaDerma Distribution, Inc. (081113521)

Revised: 12/2023

ViaDerma Distribution, Inc.