

LOW DOSE ASPIRIN ENTERIC SAFETY COATED- aspirin tablet, delayed release
Time-Cap Labs, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

381R ASA 81 DR TABS

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID*).....Pain reliever

*nonsteroidal anti-inflammatory drug

anhydrous lactose, black iron oxide, brown iron oxide, colloidal silicon dioxide, croscarmellose sodium, d&C yellow #10 aluminum lake, hypromellose, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, Polysorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Purpose: Pain Reliever

Uses

- temporarily relieves minor aches and pains
 - for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effects may occur
-

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
-

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
-

-
-

Ask a doctor before use if

- stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 - you are taking a diuretic
 - you have asthma
 - you have not been drinking fluids
 - you have lost a lot of fluid due to vomiting or diarrhea
-

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
 - taking any other drug
 - under a doctor's care for any serious condition
-

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - ringing in the ears or a loss of hearing occurs
-

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose

- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Time-Cap Labs, Inc. NDC 49483-381-10
 **Compare to the active ingredient in Bayer® Low Dose Aspirin 81 mg (NSAID) Tablets, USP.

LOW DOSE ASPIRIN ENTERIC SAFETY COATED

Pain Reliever (NSAID)
 SEE NEW WARNINGS INFORMATION
 THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN
1000 ENTERIC COATED TABLETS New Improved Formula

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID) Pain reliever nonsteroidal anti-inflammatory drug</p> <p>Uses • temporarily relieves minor aches and pains • for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effects may occur</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: • hives • facial swelling • shock • asthma (wheezing) Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed</p>	<p>Drug Facts (continued)</p> <p>• Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) Do not use • if you are allergic to aspirin or any other pain reliever/fever reducer Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have not been drinking fluids • you have lost a lot of fluid due to vomiting or diarrhea Ask a doctor or pharmacist before use if you are • taking a prescription drug for diabetes, gout, or arthritis • taking any other drug • under a doctor's care for any serious condition Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding: • feel faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear • ringing in the ears or a loss of hearing occurs</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions • drink a full glass of water with each dose • adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years: consult a doctor Other information • store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) • use by expiration date on package Inactive ingredients anhydrous lactose, black iron oxide, brown iron oxide, colloidal silicon dioxide, croscarmellose sodium, DCC yellow #10 aluminum lake, hypromellose, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, poly sorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate</p> <p>***This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Low Dose, 81 mg OTC.</p>
--	---	--

Manufactured by Time-Cap Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735
NEVER EXCEED. DO NOT USE THIS PRODUCT IF THE IMPRINTED OR SEAL OVER THE BARCODE OF THIS TABLET IS CRACKED, BLENDED, OR DAMAGED.
 LOT #: 381R 0710
 EXP. DATE: 49881108

LOW DOSE ASPIRIN ENTERIC SAFETY COATED

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-381
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B710)	
TALC (UNII: 7SEV7J4R1U)	

METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)
SODIUM HYDROXIDE (UNII: 55X04QC32I)
DIMETHICONE (UNII: 92RU3N3Y1O)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
STARCH, CORN (UNII: O8232NY3SJ)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)
POLYSORBATE 80 (UNII: 6OZP39ZG8H)
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
BROWN IRON OXIDE (UNII: 1N032N7MFO)
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)
HYPROMELLOSES (UNII: 3NXW29V3WO)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
MINERAL OIL (UNII: T5L8T28FGP)
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	T81
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-381-12	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/17/2018	
2	NDC:49483-381-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/17/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	08/10/2010	

Labeler - Time-Cap Labs, Inc (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
Time-Cap Labs, Inc		037052099	manufacture(49483-381)