

**ASPIRIN 325 MG EC- aspirin tablet  
Bryant Ranch Prepack**

-----  
**ASPIRIN 325 MG EC TABLET**

***Drug Facts***

Aspirin USP, 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

Pain Reliever/Fever Reducer

temporarily relieves

- headache
- menstrual pain
- minor pain of arthritis
- muscle pain
- toothache
- pain and fever of colds

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

***Drug Facts*** continued on back of label

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout or arthritis
- under a doctor's care for any serious condition
- taking any other drug

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
    - pain gets worse or lasts more than 10 days
    - fever gets worse or lasts more than 3 days
    - redness or swelling is present in the painful area
    - any new symptoms appear
    - ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

DRINK A FULL GLASS OF WATER WITH EACH DOSE

- **adults and children 12 years and over:** 1-2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- **children under 12 years:** ask a doctor

### **Other information**

- store at room temperature 15°-30°C (59°-86°F)

### ***Inactive ingredients***

croscarmellose sodium, D&C yellow #10 (Al-lake), FD&C yellow #6 (Al-lake), HPMC, methacrylic acid copolymer, polyethylene glycol, propylene glycol, povidone, sodium lauryl sulfate, starch, stearic acid, talc, titanium dioxide

**Questions or comments?**

**Questions or comments ? call 516-341-0666, 8:30 am - 4:30 pm ET, Monday - Friday**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark Ecotrin®.

Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**

**www.reliable1labs.com**

**HOW SUPPLIED**

- NDC: 71335-1800-1: 30 Tablets in a BOTTLE
- NDC: 71335-1800-2: 60 Tablets in a BOTTLE
- NDC: 71335-1800-3: 100 Tablets in a BOTTLE
- NDC: 71335-1800-4: 90 Tablets in a BOTTLE
- NDC: 71335-1800-5: 20 Tablets in a BOTTLE
- NDC: 71335-1800-6: 45 Tablets in a BOTTLE
- NDC: 71335-1800-7: 120 Tablets in a BOTTLE
- NDC: 71335-1800-8: 7 Tablets in a BOTTLE

**Aspirin E.C 325mg Tablet**



GTIN 0037335180010  
 Lot 20820  
 Exp 1/25/2026  
 SN 0123456789

Drug Facts	
<b>Active ingredient (in each tablet)</b>	<b>Purpose</b>
Aspirin 325 mg (NSAID)	Pain reliever
nonsteroidal anti-inflammatory drug	
<b>Uses</b>	
Temporarily relieves minor aches and pains due to: •headache •muscle pain •toothache •menstrual pain •colds •minor arthritis pain or as recommended by a doctor	
<b>Warnings</b>	
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: •hives •facial swelling •hitch •asthma (wheezing) •stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: •are age 60 or older •have had stomach ulcers or bleeding problems •take a blood thinner (anticoagulant) or steroid drug •take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) •have 3 or more alcoholic drinks every day while using this product •take more or for a longer time than directed. Do not use if you are allergic to aspirin or any other pain reliever/fever reducer. Stop use and ask a doctor if: •an allergic reaction occurs. Seek medical help right away. •you experience any of the following signs of stomach bleeding: •feel faint, have bloody or black stools, vomit blood, have stomach pain that does not get better, •pain gets worse or lasts more than 10 days •redness or swelling is present •new symptoms occur •ringing in the ears or a loss of hearing occurs. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
<b>Other information</b>	
•Each tablet contains sodium 1 mg (VERY LOW SODIUM)	
•Store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)	
•Use by expiration date on package	
<b>Directions</b>	
•Drink a full glass of water with each dose	
•Adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor	
•Children under 12 years: consult a doctor	
<b>Inactive ingredients</b>	
Corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate	

**NDC 71335-1800-1**

**Aspirin Tablets USP**

**325 mg**

**30 Tablets**



**Repackaged by:**  
 Bryant Ranch Prepack, Inc.  
 Burbank, CA 91504 USA

**Manufactured by:**  
 Reliable 1  
 Laboratories LLC



**ASPIRIN 325 MG EC**

aspirin tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71335-1800(NDC:69618-015)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>POLYETHYLENE GLYCOL 1000</b> (UNII: U076Q6Q621)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	AP;110
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71335-1800-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
2	NDC:71335-1800-2	60 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
3	NDC:71335-1800-3	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
4	NDC:71335-1800-4	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
5	NDC:71335-1800-5	20 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	

6	NDC:71335-1800-6	45 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
7	NDC:71335-1800-7	120 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
8	NDC:71335-1800-8	7 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	



# ASPIRIN 325mg EC Tablet

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/01/2015	

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

**Establishment**

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(71335-1800) , RELABEL(71335-1800)

Revised: 1/2024

Bryant Ranch Prepack