

ACETAMINOPHEN REGULAR STRENGTH- acetaminophen tablet
Magno-Humphries, Inc.

Acetaminophen Regular Strength

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains associated with:
- headache
- muscular aches
- backache
- arthritis
- common cold
- toothache
- menstrual cramps
- temporarily reduces fever

Warnings

This product contains acetaminophen. Severe liver damage may occur if you take **Liver warning:**

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 tablets every 4 to 6 hours as needed• do not take more than 8 tablets in 24 hours• do not take for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">• take 1 tablets every 4 to 6 hours as needed• do not take more than 4 tablets in 24 hours• do not take for more than 5 days unless directed by a doctor
children under 6 years	do not use this adult Regular Strength product in children under 6 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage.

Other information

- store at room temperature. Protect from moisture.
- avoid excessive heat 40°C (above 104°F)


Inactive ingredients

Corn starch, pregelatinized starch, stearic acid, may contain povidone, sodium starch glycolate


Questions?

call toll-free 1-800-935-6737

Package Labeling:



0 43292156451 5




NDC 54257-720-02

Compare to the active ingredient in Tylenol® Regular Strength*

ACETAMINOPHEN

REGULAR STRENGTH



- PAIN RELIEVER
- FEVER REDUCER

100 TABLETS • 325 MG EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each tablet) Purpose
Acetaminophen 325 mgPain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
headache ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

PEEL FOR DIRECTIONS 67206-100-01-0

LIFT HERE

Distributed by:
Magno-Humphries Labs
Tigard, OH 5723 U.S.A.

Visit our website:
www.magno-humphries.com

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

adults and children 12 years and over	take 2 tablets every 4 to 6 hours while symptoms last; do not take more than 10 tablets in 24 hours; unless directed by a doctor, do not use for more than 10 days unless directed by a doctor
children 6 years to under 12 years	take 1 tablet every 4 to 6 hours while symptoms last; do not take more than 5 tablets in 24 hours; do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Other information

- store at room temperature
- avoid excessive heat, 40° C (above 104°F)
- protect from moisture

Inactive ingredients
corn starch, pregelatinized starch, stearic acid, may contain polydione, sodium starch glycolate

Questions? call toll-free 1-800-935-6737

*This product is not manufactured or distributed by the owner of the registered trademark Tylenol® Regular Strength

ACETAMINOPHEN REGULAR STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54257-720
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength

STARCH, CORN (UNII: O8232NY3SJ)

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ010
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54257-720-02	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/24/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/24/2020	

Labeler - Magno-Humphries, Inc. (063251433)

Revised: 10/2023

Magno-Humphries, Inc.