

**HEALTH MART MIGRAINE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated**  
**Strategic Sourcing Services LLC**

-----  
**McKesson Migraine Relief Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever

Pain reliever aid

**Use**

- treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
-

### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
- diabetes

- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - your migraine is not relieved or worsens after first dose
  - new or unexpected symptoms occur
  - ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not use more than directed**
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

### **Other information**

- store at 20°-25° C (68°-77° F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

### **Inactive ingredients**

carnauba wax, crospovidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Compare to Excedrin<sup>®</sup> Migraine  
active ingredients

MIGRAINE FORMULA

Migraine Relief

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets

Pain Reliever/Pain Reliever Aid

ACTUAL SIZE

Gluten Free

100 Coated Caplets\*\*

(\*\*Capsule-Shaped Tablets)



NDC 62011-0243-1  
Compare to Excedrin<sup>®</sup> Migraine  
active ingredients\*

MIGRAINE FORMULA

# Migraine Relief

Acetaminophen, Aspirin (NSAID)  
and Caffeine Tablets



ACTUAL SIZE

Pain Reliever/Pain Reliever Aid

Gluten Free

100 Coated Caplets\*\* (\*\*Capsule-Shaped Tablets)

37478 W6 C6



### Drug Facts

#### Active ingredients (in each caplet)

Active ingredients	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	Pain reliever aid

\*nonsteroidal anti-inflammatory drug

#### Use

- treats migraine

#### Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash
- If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

■ have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

### Drug Facts (continued)

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month. Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medication, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

#### Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

#### Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers

■ you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
  - diabetes ■ gout ■ asthma
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

#### Other information

- store at 20°-25°C (68°-77°F)
- dose cap tightly after use
- read all product information before using. Keep this box for important information.

**Inactive ingredients** camcuba wax, croscopolone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, polyethylene glycol, polyethylene glycol, saccharin sodium, stearic acid, titanium dioxide

**Questions or comments?** 1-800-719-9260

KEEP BOX FOR REFERENCE

\*This product is not a substituted or diluted by GSK Consumer Healthcare, distributor of Excedrin<sup>®</sup> Migraine.

**DON'T USE IF PRINTED FOLLOWS**  
**IS BROKEN OR MISSING**

**MEKESON**

Another Quality Product  
Distributed by Mekecon  
6355 State Highway 161, Irving, TX 75039  
Money Back Guarantee  
health.mart.com

# HEALTH MART MIGRAINE RELIEF

acetaminophen, aspirin, caffeine tablet, film coated

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0243
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

## Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L374
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0243-1	1 in 1 CARTON	04/22/2014	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	04/22/2014	

