

ACETAMINOPHEN- pain reliever tablet McKesson

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Acetaminophen Tablets 500 mg

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

For the temporary relief of minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- common cold
- toothache
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a

drug contains acetaminophen, ask a doctor or pharmacist.

- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed

Adults and children: (12 years and older) Take 2 tablets every 4 to 6 hours as needed. Do not take more than 8 tablets in 24 hours.

Children under 12 years: Do not give to children under 12 years of age

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, maltodextrin*, microcrystalline cellulose*, polyethylene glycol, povidone*, pregelatinized starch*, sodium starch glycolate*, stearic acid, titanium dioxide*.

* may contain

Questions or comments? 1-800-777-4908

NDC# 68599-4681-3

McKesson

Acetaminophen Tablets 500mg

Non-Aspirin

Used to reduce fever / Temporarily relieves aches and pains

2 per Pack / 250 packs per Box

MFR# 82468

NDC 68599-4681-3

McKESSON

Acetaminophen Tablets 500mg NON-ASPIRIN

2

PER PACK

MFR # 82468

| Active ingredient (in each tablet) | Purpose |
|------------------------------------|-----------------------------|
| Acetaminophen..... | Pain reliever/fever reducer |

Uses For the temporary relief of minor aches and pains associated with

- headache
- toothache
- muscular aches
- common cold
- minor arthritis pain
- menstrual cramps

For the reduction of fever.

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 12 tablets in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Store at room temperature 59-86°F (15-30°C)
 - Tamper evident sealed packets
 - Do not use any opened or torn packets

Warnings (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed

Adults and children: (12 years and older)

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

Children under 12 years:

Do not give to children under 12 years of age

Questions or comments? Call 800-777-4908

SEE BOX FOR COMPLETE PRODUCT INFORMATION

Retain carton for complete product information

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

McKESSON

Drug Facts

| Active ingredient (in each tablet) | Purpose |
|------------------------------------|-----------------------------|
| Acetaminophen 500mg..... | Pain reliever/fever reducer |

Uses
For the temporary relief of minor aches and pains associated with

- headache
- toothache
- common cold
- muscular aches
- minor arthritis pain
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if
■ you are taking the blood thinning drug warfarin

Drug Facts (continued)

Stop using and ask a doctor if

- symptoms do not improve
- pain or fever persists or gets worse
- new symptoms occur
- redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not use more than directed

Adults and children: (12 years and older) Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

Children under 12 years: Do not give to children under 12 years of age

Other information

- store at room temperature 59-86°F (15-30°C)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, maltodextrin*, microcrystalline cellulose*, polyethylene glycol, povidone*, pregelatinized starch*, sodium starch glycolate*, stearic acid, titanium dioxide*

*may contain

Questions or comments? Call 800,777,4908

Satisfaction Guaranteed

For complete details, please visit
mms.mckesson.com/mckesson-brands.

Distributed by McKesson Medical-Surgical Inc.
Richmond, VA 23233
PVN A0120
Made in USA

LOT

EXP

MFR # 82468



NDC 68599-4681-3

3³¹/₃₂

McKESSON

Acetaminophen Tablets 500mg NON-ASPIRIN

USED TO REDUCE FEVER
TEMPORARILY RELIEVES ACHEs AND PAINS

2

PER PACK

250

PACKS PER BOX

MFR #

82468

NDC 68599-4681-3

McKesson

82468

Acetaminophen Tablets

500mg | NON-ASPIRIN

USED TO REDUCE FEVER | TEMPORARILY RELIEVES ACHES AND PAINS

2 | **250** | **12**
PER PACK | PACKS PER BOX | BOXES PER CASE

MFR #
82468

Store at room temperature 59-86°F (15-30°C).
Not made with natural rubber latex.

Distributed by McKesson Medical-Surgical Inc.
Richmond, VA 23233
PVN A0120 Made in USA



LOT

EXP

ACETAMINOPHEN

pain reliever tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68599-4681 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|--------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 0.5 g in 1 g |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| GLYCOLIC ACID (UNII: 0WT12SX38S) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE (UNII: FZ989GH94E) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|--------------------------|
| Color | white | Score | score with uneven pieces |
| Shape | ROUND | Size | 12mm |
| Flavor | | Imprint Code | FR;33 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:68599-4681-3 | 0.5 g in 1 POUCH; Type 0: Not a Combination Product | 01/30/2020 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part343 | 01/30/2020 | |

Labeler - McKesson (023904428)

Establishment

| Name | Address | ID/FEI | Business Operations |
|------------------|---------|-----------|---------------------|
| Medique Products | | 086911794 | pack(68599-4681) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|------------------|---------|-----------|-------------------------|
| Allegiant Health | | 079501930 | manufacture(68599-4681) |

Revised: 3/2023

McKesson