LEUCOVORIN CALCIUM- leucovorin calcium injection, powder, lyophilized, for solution

Meitheal Pharmaceuticals Inc.

-----

Leucovorin Calcium for Injection, USP

meitheal<sub>®</sub> Rx only

#### DESCRIPTION

Leucovorin is one of several active, chemically reduced derivatives of folic acid. It is useful as an antidote to drugs which act as folic acid antagonists.

Also known as folinic acid, Citrovorum factor, or 5-formyl-5,6,7,8-tetrahydrofolic acid, this compound has the chemical designation of L-Glutamic acid, N-[4-[[(2-amino-5-formyl-1,4,5,6,7,8-hexahydro-4-oxo-6-pteridinyl)methyl]amino]benzoyl]-calcium salt (1:1). The formula weight is 511.51 and the structural formula of leucovorin calcium is:



C20H21CaN7O7

Leucovorin Calcium for Injection is a sterile product indicated for intramuscular (IM) or intravenous (IV) administration and is supplied in 50 mg, 100 mg, 200 mg, 350 mg and 500 mg vials.

Each 50 mg vial of Leucovorin Calcium for Injection, when reconstituted with 5 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

Each 100 mg vial of Leucovorin Calcium for Injection, when reconstituted with 10 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

Each 200 mg vial of Leucovorin Calcium for Injection, when reconstituted with 20 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

Each 350 mg vial of Leucovorin Calcium for Injection, when reconstituted with 17.5 mL of sterile diluent, contains leucovorin (as the calcium salt) 20 mg/mL.

Each 500 mg vial of Leucovorin Calcium for Injection, when reconstituted with 50 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

In each dosage form, one milligram of leucovorin calcium contains 0.002 mmol of leucovorin and 0.002 mmol of calcium.

These lyophilized products contain no preservative. The inactive ingredient is Sodium Chloride, USP, added to adjust tonicity and sodium hydroxide and/or hydrochloric acid for pH adjustment. pH adjusted to 7.0 to 8.5.

Reconstitute with Bacteriostatic Water for Injection, USP, which contains benzyl alcohol (see **WARNINGS** section), or with Sterile Water for Injection, USP.

## CLINICAL PHARMACOLOGY

Leucovorin is a mixture of the diastereoisomers of the 5-formyl derivative of tetrahydrofolic acid (THF). The biologically active compound of the mixture is the (-)-*l*-isomer, known as Citrovorum factor, or (-)-folinic acid. Leucovorin does not require reduction by the enzyme dihydrofolate reductase in order to participate in reactions utilizing folates as a source of "one-carbon" moieties. *l*-Leucovorin (*l*-5-formyltetrahydrofolate) is rapidly metabolized (via 5,10-methenyltetrahydrofolate then 5,10-methylenetetrahydrofolate) to *l*-5-methyltetrahydrofolate. *l*-5-Methyltetrahydrofolate can in turn be metabolized via other pathways back to 5,10-methylenetetrahydrofolate, which is converted to 5-methyltetrahydrofolate by an irreversible, enzyme catalyzed reduction using the cofactors FADH<sub>2</sub> and NADPH.

Administration of leucovorin calcium can counteract the therapeutic and toxic effects of folic acid antagonists such as methotrexate, which act by inhibiting dihydrofolate reductase.

In contrast, leucovorin calcium can enhance the therapeutic and toxic effects of fluoropyrimidines used in cancer therapy, such as 5-fluorouracil. Concurrent administration of leucovorin calcium does not appear to alter the plasma pharmacokinetics of 5-fluorouracil. 5-Fluorouracil is metabolized to fluorodeoxyuridylic acid, which binds to and inhibits the enzyme thymidylate synthase (an enzyme important in DNA repair and replication).

Leucovorin is readily converted to another reduced folate, 5,10methylenetetrahydrofolate, which acts to stabilize the binding of fluorodeoxyuridylic acid to thymidylate synthase and thereby enhances the inhibition of this enzyme.

The pharmacokinetics after intravenous, intramuscular, and oral administration of a 25 mg dose of leucovorin calcium were studied in male volunteers. After intravenous administration, serum total reduced folates (as measured by *Lactobacillus casei* assay) reached a mean peak of 1259 ng/mL (range 897 – 1625). The mean time to peak was 10 minutes. This initial rise in total reduced folates was primarily due to the parent compound 5-formyl-THF (measured by *Streptococcus faecalis* assay) which rose to 1206 ng/mL at 10 minutes. A sharp drop in parent compound followed and coincided with the appearance of the active metabolite 5-methyl-THF which became the predominant circulating form of the drug.

The mean peak of 5-methyl-THF was 258 ng/mL and occurred at 1.3 hours. The terminal half-life for total reduced folates was 6.2 hours. The area under the concentration versus time curves (AUCs) for *l*-leucovorin, *d*-leucovorin and 5-methyltetrahydrofolate were 28.4  $\pm$  3.5, 956  $\pm$  97 and 129  $\pm$  12 (mg/min/L  $\pm$  S.E.). When a higher dose of *d*,*l*-leucovorin (200 mg/m<sup>2</sup>) was used, similar results were obtained. The *d*-isomer persisted in plasma at concentrations greatly exceeding those of the *l*-isomer.

After intramuscular injection, the mean peak of serum total reduced folates was 436

ng/mL (range 240-725) and occurred at 52 minutes. Similar to IV administration, the initial sharp rise was due to the parent compound. The mean peak of 5-formyl-THF was 360 ng/mL and occurred at 28 minutes. The level of the metabolite 5-methyl-THF increased subsequently over time until at 1.5 hours it represented 50% of the circulating total folates. The mean peak of 5-methyl-THF was 226 ng/mL at 2.8 hours. The terminal half-life of total reduced folates was 6.2 hours. There was no difference of statistical significance between IM and IV administration in the AUC for total reduced folates, 5-formyl-THF, or 5-methyl-THF.

After oral administration of leucovorin calcium reconstituted with aromatic elixir, the mean peak concentration of serum total reduced folates was 393 ng/mL (range 160-550). The mean time to peak was 2.3 hours and the terminal half-life was 5.7 hours. The major component was the metabolite 5-methyltetrahydrofolate to which leucovorin is primarily converted in the intestinal mucosa. The mean peak of 5-methyl-THF was 367 ng/mL at 2.4 hours. The peak level of the parent compound was 51 ng/mL at 1.2 hours. The AUC of total reduced folates after oral administration of the 25 mg dose was 92% of the AUC after intravenous administration.

Following oral administration, leucovorin calcium is rapidly absorbed and expands the serum pool of reduced folates. At a dose of 25 mg, almost 100% of the *I*-isomer but only 20% of the *d*-isomer is absorbed. Oral absorption of leucovorin calcium is saturable at doses above 25 mg. The apparent bioavailability of leucovorin was 97% for 25 mg, 75% for 50 mg, and 37% for 100 mg.

In a randomized clinical study conducted by the Mayo Clinic and the North Central Cancer Treatment Group (Mayo/NCCTG) in patients with advanced metastatic colorectal cancer three treatment regimens were compared: Leucovorin calcium (LV) 200 mg/m<sup>2</sup> and 5-fluorouracil (5-FU) 370 mg/m<sup>2</sup> versus LV 20 mg/m<sup>2</sup> and 5-FU 425 mg/m<sup>2</sup> versus 5-FU 500 mg/m<sup>2</sup>. All drugs were administered by slow intravenous infusion daily for 5 days repeated every 28-35 days. Response rates were 26% (p = 0.04 versus 5-FU alone), 43% (p = 0.001 versus 5-FU alone) and 10% for the high dose leucovorin calcium, low dose leucovorin calcium and 5-FU alone groups respectively. Respective median survival times were 12.2 months (p = 0.037), 12 months (p = 0.050), and 7.7 months. The low dose LV regimen gave a statistically significant improvement in weight gain of more than 5%, relief of symptoms, and improvement in performance status. The high dose LV regimen gave a statistically significant improvement in performance status and trended toward improvement in weight gain and in relief of symptoms but these were not statistically significant.

In a second Mayo/NCCTG randomized clinical study the 5-FU alone arm was replaced by a regimen of sequentially administered methotrexate (MTX), 5-FU, and LV. Response rates with LV 200 mg/m<sup>2</sup> and 5-FU 370 mg/m<sup>2</sup> versus LV 20 mg/m<sup>2</sup> and 5-FU 425 mg/m<sup>2</sup> versus sequential MTX and 5-FU and LV were respectively 31% (p = < 0.01), 42% (p = < 0.01), and 14%. Respective median survival times were 12.7 months (p = < 0.04), 12.7 months (p = < 0.01), and 8.4 months. No statistically significant difference in weight gain of more than 5% or in improvement in performance status was seen between the treatment arms.

The pharmacokinetics of 200 mg doses of leucovorin calcium administered intravenously and orally (reconstituted powder, not tablets) have been evaluated in healthy male subjects. The serum clearance corrected for bioavailability, terminal half-life, and apparent volume of distribution of total folate were not significantly different between the routes of administration. The oral bioavailability of the 200 mg dose was 31%. Eighty-three percent of the biologically active IV dose was recovered in the urine within 24 hours, 31% as 5-methyltetrahydrofolate. Twenty percent of the same oral dose was excreted in 24 hours, 16% as 5-methyltetrahydrofolate.

## INDICATIONS AND USAGE

Leucovorin Calcium for Injection rescue is indicated after high dose methotrexate therapy in osteosarcoma. Leucovorin Calcium for Injection is also indicated to diminish the toxicity and counteract the effects of impaired methotrexate elimination and of inadvertent overdosages of folic acid antagonists.

Leucovorin Calcium for Injection is indicated in the treatment of megaloblastic anemias due to folic acid deficiency when oral therapy is not feasible.

Leucovorin Calcium for Injection is also indicated for use in combination with 5fluorouracil to prolong survival in the palliative treatment of patients with advanced colorectal cancer. Leucovorin Calcium for Injection should not be mixed in the same infusion as 5-fluorouracil because a precipitate may form.

## CONTRAINDICATIONS

Leucovorin calcium is improper therapy for pernicious anemia and other megaloblastic anemias secondary to the lack of vitamin  $B_{12}$ . A hematologic remission may occur while neurologic manifestations continue to progress.

## WARNINGS

In the treatment of accidental overdosages of folic acid antagonists, intravenous leucovorin calcium should be administered as promptly as possible. As the time interval between antifolate administration (e.g., methotrexate) and leucovorin rescue increases, leucovorin calcium's effectiveness in counteracting toxicity decreases. In the treatment of accidental overdosages of intrathecally administered folic acid antagonists, do not administer leucovorin calcium intrathecally. LEUCOVORIN CALCIUM MAY BE HARMFUL OR FATAL IF GIVEN INTRATHECALLY.

Monitoring of the serum methotrexate concentration is essential in determining the optimal dose and duration of treatment with leucovorin calcium.

Delayed methotrexate excretion may be caused by a third space fluid accumulation (ie, ascites, pleural effusion), renal insufficiency, or inadequate hydration. Under such circumstances, higher doses of leucovorin calcium or prolonged administration may be indicated. Doses higher than those recommended for oral use must be given intravenously.

Because of the benzyl alcohol contained in certain diluents used for reconstituting Leucovorin Calcium for Injection, when doses greater than 10 mg/m<sup>2</sup> are administered, leucovorin calcium for injection should be reconstituted with Sterile Water for Injection, USP, and used immediately. (See **DOSAGE AND ADMINISTRATION**.)

Because of the calcium content of the leucovorin calcium solution, no more than 160 mg of leucovorin should be injected intravenously per minute (16 mL of a 10 mg per mL, or

8 mL of a 20 mg per mL solution per minute).

Leucovorin calcium enhances the toxicity of 5-fluorouracil. When these drugs are administered concurrently in the palliative therapy of advanced colorectal cancer, the dosage of 5-fluorouracil must be lower than usually administered. Although the toxicities observed in patients treated with the combination of leucovorin calcium plus 5fluorouracil are qualitatively similar to those observed in patients treated with 5fluorouracil alone, gastrointestinal toxicities (particularly stomatitis and diarrhea) are observed more commonly and may be more severe and of prolonged duration in patients treated with the combination.

In the first Mayo/NCCTG controlled trial, toxicity, primarily gastrointestinal, resulted in 7% of patients requiring hospitalization when treated with 5-fluorouracil alone or 5-fluorouracil in combination with 200 mg/m<sup>2</sup> of leucovorin calcium and 20% when treated with 5-fluorouracil in combination with 20 mg/m<sup>2</sup> of leucovorin calcium. In the second Mayo/NCCTG trial, hospitalizations related to treatment toxicity also appeared to occur more often in patients treated with the low dose leucovorin calcium/5-fluorouracil combination than in patients treated with the high dose combination — 11% versus 3%. Therapy with leucovorin calcium and 5-fluorouracil must not be initiated or continued in patients who have symptoms of gastrointestinal toxicity of any severity, until those symptoms have completely resolved. Patients with diarrhea must be monitored with particular care until the diarrhea has resolved, as rapid clinical deterioration leading to death can occur. In an additional study utilizing higher weekly doses of 5-fluorouracil and leucovorin calcium, elderly and/or debilitated patients were found to be at greater risk for severe gastrointestinal toxicity.

Seizures and/or syncope have been reported rarely in cancer patients receiving leucovorin calcium, usually in association with fluoropyrimidine administration, and most commonly in those with CNS metastases or other predisposing factors, however, a causal relationship has not been established.

The concomitant use of leucovorin calcium with trimethoprim-sulfamethoxazole for the acute treatment of Pneumocystis carinii pneumonia in patients with HIV infection was associated with increased rates of treatment failure and morbidity in a placebo-controlled study.

## PRECAUTIONS

## General

Parenteral administration is preferable to oral dosing if there is a possibility that the patient may vomit and not absorb the leucovorin. Leucovorin has no effect on non-hematologic toxicities of methotrexate, such as the nephrotoxicity resulting from drug and/or metabolite precipitation in the kidney.

Since leucovorin calcium enhances the toxicity of fluorouracil, leucovorin calcium/5-fluorouracil combination therapy for advanced colorectal cancer should be administered under the supervision of a physician experienced in the use of antimetabolite cancer chemotherapy. Particular care should be taken in the treatment of elderly or debilitated colorectal cancer patients, as these patients may be at increased risk of severe toxicity.

## Laboratory Tests

Patients being treated with the leucovorin calcium/5-fluorouracil combination should have a CBC with differential and platelets prior to each treatment. During the first two courses a CBC with differential and platelets has to be repeated weekly and thereafter once each cycle at the time of anticipated WBC nadir. Electrolytes and liver function tests should be performed prior to each treatment for the first three cycles then prior to every other cycle. Dosage modifications of fluorouracil should be instituted as follows, based on the most severe toxicities:

Diarrhea and/or Stomatitis	WBC/mm <sup>3</sup> Nadir	Platelets/mm <sup>3</sup> Nadir	5-FU Dose
Moderate	1,000-1,900	25-75,000	decrease 20%
Severe	< 1,000	< 25,000	decrease 30%

If no toxicity occurs, the 5-fluorouracil dose may increase 10%. Treatment should be deferred until WBC's are 4,000/mm<sup>3</sup> and platelets 130,000/mm<sup>3</sup>. If blood counts do not reach these levels within two weeks, treatment should be discontinued. Patients should be followed up with physical examination prior to each treatment course and appropriate radiological examination as needed. Treatment should be discontinued when there is clear evidence of tumor progression.

## **Drug Interactions**

Folic acid in large amounts may counteract the antiepileptic effect of phenobarbital, phenytoin and primidone, and increase the frequency of seizures in susceptible pediatric patients.

Preliminary animal and human studies have shown that small quantities of systemically administered leucovorin calcium enter the CSF primarily as 5-methyltetrahydrofolate and, in humans, remain 1-3 orders of magnitude lower than the usual methotrexate concentrations following intrathecal administration. However, high doses of leucovorin calcium may reduce the efficacy of intrathecally administered methotrexate.

Leucovorin calcium may enhance the toxicity of 5-fluorouracil. (See WARNINGS.)

## Pregnancy

## Teratogenic Effects

Adequate animal reproduction studies have not been conducted with leucovorin calcium. It is also not known whether leucovorin calcium can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Leucovorin calcium should be given to a pregnant woman only if clearly needed.

## Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when leucovorin calcium is administered to a nursing mother.

## Pediatric Use

## See PRECAUTIONS, Drug Interactions.

## Geriatric Use

Clinical studies of leucovorin calcium did not show differences in safety or effectiveness between subjects over 65 and younger subjects. Other clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older patients cannot be ruled out. This drug is known to be excreted by the kidney and the risk of toxic reactions to the drug may be greater in patients with impaired renal function. Because elder patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **ADVERSE REACTIONS**

Diarrhea

Allergic sensitization, including anaphylactoid reactions and urticaria, has been reported following administration of both oral and parenteral leucovorin calcium. No other adverse reactions have been attributed to the use of leucovorin calcium per se.

The following table summarizes significant adverse events occurring in 316 patients treated with the leucovorin calcium/5-fluorouracil combinations compared against 70 patients treated with 5-fluorouracil alone for advanced colorectal carcinoma. These data are taken from the Mayo/NCCTG large multicenter prospective trial evaluating the efficacy and safety of the combination regimen.

PERCENTAGE OF PATI	ENTS TREA	ATED WITH L	EUCOV	ORIN CALCIU	M/FLUC	ROURACIL
FC	OR ADVAN	CED COLOR	ECTAL C	ARCINOMA		
REPORTING AD	VERSE EX	PERIENCES (	or hose	PITALIZED FO	OR TOXI	CITY
	(Higł	า LV)/5-FU	(Lov	v LV)/5-FU	5-l	FU Alone
	1)	N=155)	1)	N=161)	(	N=70)
	Any	Grade3+	Any	Grade 3+	Any	Grade 3+
	<u>(%)</u>	<u>(%)</u>	<u>(%)</u>	<u>(%)</u>	<u>(%)</u>	<u>(%)</u>
Leukopenia	69	14	83	23	93	48
Thrombocytopenia	8	2	8	1	18	3
Infection	8	1	3	1	7	2
Nausea	74	10	80	9	60	6
Vomiting	46	8	44	9	40	7

Blaithea	00	±0	07		15			
Stomatitis	75	27	84	29	59			
Constipation	3	0	4	0	1			
Lethargy/Malaise/Fatigue	13	3	12	2	6			
Alopecia	42	5	43	6	37			
Dermatitis	21	2	25	1	13			
Anorexia	14	1	22	4	14			
Hospitalization for Toxicity		5%	]	L5%		7%		
High LV= Leucovorin calcium 200 mg/m <sup>2</sup>								
Low LV=Leucovorin calcium 20 mg/m <sup>2</sup>								
Any= percentage of patients re	eporting	toxicity of any	severity					

18

67

14

43

66

To report SUSPECTED ADVERSE REACTIONS, contact Meitheal Pharmaceuticals Inc. at 1-844-824-8426 or FDA at 1-800-FDA-1088 or <u>www.fda.gov/medwatch</u>.

## OVERDOSAGE

Excessive amounts of leucovorin calcium may nullify the chemotherapeutic effect of folic acid antagonists.

## DOSAGE AND ADMINISTRATION

## Advanced Colorectal Cancer

Either of the following two regimens is recommended:

- 1. Leucovorin calcium for injection is administered at 200 mg/m<sup>2</sup> by slow intravenous injection over a minimum of 3 minutes, followed by 5-fluorouracil at 370 mg/m<sup>2</sup> by intravenous injection.
- 2. Leucovorin calcium for injection is administered at 20 mg/m<sup>2</sup> by intravenous injection followed by 5-fluorouracil at 425 mg/m<sup>2</sup> by intravenous injection.

5-Fluorouracil and leucovorin calcium for injection should be administered separately to avoid the formation of a precipitate.

Treatment is repeated daily for five days. This five-day treatment course may be repeated at 4 week (28-day) intervals, for 2 courses and then repeated at 4-5 week (28-35 day) intervals provided that the patient has completely recovered from the toxic effects of the prior treatment course.

In subsequent treatment courses, the dosage of 5-fluorouracil should be adjusted based on patient tolerance of the prior treatment course. The daily dosage of 5fluorouracil should be reduced by 20% for patients who experienced moderate hematologic or gastrointestinal toxicity in the prior treatment course, and by 30% for patients who experienced severe toxicity (see **PRECAUTIONS: Laboratory Tests**). For patients who experienced no toxicity in the prior treatment course, 5-fluorouracil dosages may be increased by 10%. Leucovorin calcium for injection dosages are not adjusted for toxicity.

## Leucovorin Rescue After High-Dose Methotrexate Therapy

The recommendations for leucovorin rescue are based on a methotrexate dose of 12-15 grams/m<sup>2</sup> administered by intravenous infusion over 4 hours (see methotrexate package insert for full prescribing information).

Leucovorin rescue at a dose of 15 mg (approximately 10 mg/m<sup>2</sup>) every 6 hours for 10 doses starts 24 hours after the beginning of the methotrexate infusion. In the presence of gastrointestinal toxicity, nausea or vomiting, leucovorin calcium for injection should be administered parenterally. Do not administer leucovorin calcium for injection intrathecally.

Serum creatinine and methotrexate levels should be determined at least once daily.

Leucovorin calcium for injection administration, hydration, and urinary alkalization (pH of 7.0 or greater) should be continued until the methotrexate level is below  $5 \times 10^{-8}$  M (0.05 micromolar). The leucovorin calcium for injection dose should be adjusted or leucovorin rescue extended based on the following guidelines:

# GUIDELINES FOR LEUCOVORIN CALCIUM FOR INJECTION DOSAGE AND ADMINISTRATION

DO NOT ADMINISTER LEUCOVORIN CALCIUM FOR INJECTION INTRATHECALLY

Clinical Situation	Laboratory Findings	Leucovorin Calcium for Injection Dosage and Duration
Normal Methotrexate Elimination	Serum methotrexate level approximately 10 micromolar at 24 hours after administration, 1 micromolar at 48 hours, and less than 0.2 micromolar at 72 hours.	15 mg PO, IM, or IV q 6 hours for 60 hours (10 doses starting at 24 hours after start of methotrexate infusion).
Delayed Late Methotrexate Elimination	Serum methotrexate level remaining above 0.2 emicromolar at 72 hours, and more than 0.05 micromolar at 96 hours after administration.	Continue 15 mg PO, IM, or IV q 6 hours, until methotrexate level is less than 0.05 micromolar.
Delayed Early Methotrexate Elimination and/or Evidence of Acute Renal Injury	Serum methotrexate level of 50 micromolar or emore at 24 hours, or 5 micromolar or more at 48 hours after administration, OR; a 100% or greater increase in serum creatinine level at 24 hours after methotrexate administration (eg, an increase from 0.5 mg/dL to a level of 1 mg/dL or more)	150 mg IV q 3 hours, until methotrexate level is less than 1 micromolar; then 15 mg IV q 3 hours until methotrexate level is less than 0.05 micromolar.

Patients who experience delayed early methotrexate elimination are likely to develop reversible renal failure. In addition to appropriate leucovorin calcium for injection therapy, these patients require continuing hydration and urinary alkalinization, and close monitoring of fluid and electrolyte status, until the serum methotrexate level has fallen to below 0.05 micromolar and the renal failure has resolved.

Some patients will have abnormalities in methotrexate elimination or renal function following methotrexate administration, which are significant but less severe than the abnormalities described in the table above. These abnormalities may or may not be associated with significant clinical toxicity. If significant clinical toxicity is observed, leucovorin rescue should be extended for an additional 24 hours (total of 14 doses over 84 hours) in subsequent courses of therapy. The possibility that the patient is taking other medications which interact with methotrexate (eg, medications which may interfere with methotrexate elimination or binding to serum albumin) should always be reconsidered when laboratory abnormalities or clinical toxicities are observed.

## Impaired Methotrexate Elimination or Inadvertent Overdosage

Leucovorin rescue should begin as soon as possible after an inadvertent overdosage and within 24 hours of methotrexate administration when there is delayed excretion (see **WARNINGS**). Leucovorin calcium for injection 10 mg/m<sup>2</sup> should be administered IV, IM, or PO every 6 hours until the serum methotrexate level is less than 10<sup>-8</sup> M. In the presence of gastrointestinal toxicity, nausea, or vomiting, leucovorin calcium for injection should be administered parenterally. Do not administer leucovorin calcium for injection intrathecally.

Serum creatinine and methotrexate levels should be determined at 24 hour intervals. If the 24 hour serum creatinine has increased 50% over baseline or if the 24 hour methotrexate level is greater than 5 x  $10^{-6}$  M or the 48 hour level is greater than 9 x  $10^{-7}$  M, the dose of leucovorin calcium for injection should be increased to 100 mg/m<sup>2</sup> IV every 3 hours until the methotrexate level is less than  $10^{-8}$  M.

Hydration (3 L/d) and urinary alkalinization with sodium bicarbonate solution should be employed concomitantly. The bicarbonate dose should be adjusted to maintain the urine pH at 7.0 or greater.

## Megaloblastic Anemia Due to Folic Acid Deficiency

Up to 1 mg daily. There is no evidence that doses greater than 1 mg/day have greater efficacy than those of 1 mg; additionally, loss of folate in urine becomes roughly logarithmic as the amount administered exceeds 1 mg.

Each 50 mg, 100 mg, 200 mg, and 500 mg vial of leucovorin calcium for injection when reconstituted with 5 mL, 10 mL, 20 mL, and 50 mL, respectively, of sterile diluent yields a leucovorin concentration of 10 mg per mL. Each 350 mg vial of leucovorin calcium for injection when reconstituted with 17.5 mL of sterile diluent yields a leucovorin concentration of 20 mg per mL. Leucovorin calcium for injection contains no preservative. Reconstitute the lyophilized vial products with Bacteriostatic Water for Injection, USP (benzyl alcohol preserved), or Sterile Water for Injection, USP. When reconstituted with Bacteriostatic Water for Injection, USP, the resulting solution must be used within 7 days. If the product is reconstituted with Sterile Water for Injection, USP, use immediately and discard any unused portion.

Because of the benzyl alcohol contained in Bacteriostatic Water for Injection, USP, when doses greater than 10 mg/m<sup>2</sup> are administered leucovorin calcium for injection should be reconstituted with Sterile Water for Injection, USP, and used immediately. (See **WARNINGS**.) Because of the calcium content of the leucovorin calcium for injection solution, no more than 160 mg of leucovorin should be injected intravenously per minute (16 mL of a 10 mg per mL, or 8 mL of a 20 mg per mL solution per minute).

Parenteral products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Leucovorin calcium for injection should not be mixed in the same infusion as 5-fluorouracil, since this may lead to the formation of a precipitate.

## HOW SUPPLIED

Leucovorin Calcium for Injection, USP is supplied as a sterile lyophilized powder as follows:

## NDC

#### Leucovorin Calcium for Injection, USP Package Factor

71288-**160**-10 71288-**161**-20 71288-**162**-30

50 mg Single-Dose Vial 100 mg Single-Dose Vial 200 mg Single-Dose Vial 1 vial per carton 1 vial per carton 1 vial per carton

71288- <b>163</b> -30	3
71288- <b>164</b> -50	5

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. **Protect from light.** Retain in carton until time of use.

Discard unused portion.

## Sterile, Nonpyrogenic, Preservative-free.

## The container closure is not made with natural rubber latex.

meitheal<sub>®</sub> Mfd. for Meitheal Pharmaceuticals Chicago, IL 60631 (USA) © 2024 Meitheal Pharmaceuticals Inc.

Mfd. by Kindos Pharmaceuticals Co., Ltd. Chengdu, China 611731

October 2024

LB-462-V3

#### PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 50 mg Vial Label

NDC 71288-**160**-10

Rx Only

Leucovorin Calcium for Injection, USP

50 mg per vial

For Intravenous or Intramuscular Use

Lyophilized

Single-Dose Vial



## PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 50 mg Carton

NDC 71288-**160**-10

Rx Only

1 Single-Dose Vial

Leucovorin Calcium for Injection, USP

50 mg per vial

For Intravenous or Intramuscular Use



PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 100 mg Vial Label

NDC 71288-161-20

**Rx Only** 

Leucovorin Calcium for Injection, USP

100 mg per vial

## For Intravenous or Intramuscular Use

## Lyophilized

Single-Dose Vial



## PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 100 mg Carton

NDC 71288-**161**-20

Rx Only

1 Single-Dose Vial

Leucovorin Calcium for Injection, USP

100 mg per vial

For Intravenous or Intramuscular Use



PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 200 mg Vial Label

NDC 71288-162-30

**Rx Only** 

Leucovorin Calcium for Injection, USP

200 mg per vial

For Intravenous or Intramuscular Use



## PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 200 mg Carton

NDC 71288-**162**-30

Rx Only

1 Single-Dose Vial

Leucovorin Calcium for Injection, USP

200 mg per vial

For Intravenous or Intramuscular Use



## PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 350 mg Vial Label

NDC 71288-**163**-30

Rx Only

## Leucovorin Calcium for Injection, USP

350 mg per vial

For Intravenous or Intramuscular Use

## Lyophilized

Single-Dose Vial

NDC 71288-163-30 Rx Only	For doses er for ochloric acid 0 to 8.5. 0 to 8.5. ains ains
Leucovorin Calcium for Injection, USP	information. Information. Informately 7. Informately 7. In
> 350 mg per vial	rescribing tre only will hy. (See Willium hydrow illum hydrow indjusted to a inter for nzyl alcohol g leucovori g leucovori 12881633
For Intravenous or Intramuscular Use	if for complete p gy/m², reconstitu i use immediate ucovorin Calcium dium chloride. Sod adjustment. Pl a di reconstituted so in Temperature]. Pr ed. with 17.5 mL Steri preserved with bei preserved with b
Lyophilized	used for pharman 10 m and 10 m son and add for pharmacely marked port an instituted Roor and after and the set pharman and the set pharman and the set pharmacely and a set of the set o
Single-Dose Vial	See packa greater th Injection, Each vial cc base and 1, may be add Store dry p USP Contro When recoi When recoi Use contei Leucovorin Do not use Mid. for Meii Chicago, IL 66 Made in Chin Exp.: I

PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 350 mg Carton

NDC 71288-**163**-30

Rx Only

1 Single-Dose Vial

Leucovorin Calcium for Injection, USP

350 mg per vial

For Intravenous or Intramuscular Use



PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 500 mg Vial Label

NDC 71288-164-50

**Rx Only** 

## Leucovorin Calcium for Injection, USP

500 mg per vial

## For Intravenous or Intramuscular Use

## Lyophilized

Single-Dose Vial



## PRINCIPAL DISPLAY PANEL - Leucovorin Calcium for Injection, USP 500 mg Carton

NDC 71288-**164**-50

Rx Only

1 Single-Dose Vial

Leucovorin Calcium for Injection, USP

500 mg per vial

For Intravenous or Intramuscular Use



LEUCOVORIN CALCIUM							
leucovorin calcium injection, p	owder, lyophilized, for solution						
Product Information							
Product Type	HUMAN PRESCRIPTION DRUG	ltem	Code (Source)	NDC:71288	-160		
Route of Administration	INTRAVENOUS, INTRAMUSCULAR						
Active Ingredient/Active Moiety							
Ingre	dient Name		Basis of Strengt	th Stren	gth		
LEUCOVORIN CALCIUM (UNII: RPF	(LEUCOVORIN - UNII:Q573)	9DVLP)	LEUCOVORIN	50 mg ir	ו 5 mL		

Ir	nactive Ingr	edients					
	Ingredient Name Strength						
s		<b>DE</b> (UNII: 451W47	IQ8X)				
H	<b>YDROCHLORIC</b>	ACID (UNII: QTT)	7582CB)				
S	DDIUM HYDROX	(IDE (UNII: 55X04	QC32I)				
Ρ	ackaging						
#	Item Code	Pa	ackage Description	Marketing Sta Date	rt Marketing End Date		
1	NDC:71288- 160-10	1 in 1 CARTON		07/19/2023			
1		5 mL in 1 VIAL, 9 Combination Pro	SINGLE-DOSE; Type 0: Not a oduct				
M	larketing	Informat	ion				
	Marketing	Applicat	tion Number or Monograph	Marketing Star	Markoting End		
	Category	Applica	Citation	Date	Date		
AN	Category	ANDA21659	<b>Citation</b>	Date 07/19/2023	Date		
AN	<b>Category</b> IDA	ANDA21659	<b>Citation</b>	Date	Date		
AN	<b>Category</b> IDA	ANDA21659	Citation (	Date	Date		
	Category IDA EUCOVOR	ANDA216590	Citation (	Date 07/19/2023	Date		
	Category IDA EUCOVOR	ANDA216590	Citation D JM powder, lyophilized, for solution	Date 07/19/2023	Date		
	Category IDA EUCOVOR	ANDA216590	<b>Citation</b> <b>JM</b> bowder, lyophilized, for solution	Date 07/19/2023	Date		
AN LI let	Category IDA EUCOVOR JCOVORIN Calciu roduct Info	ANDA216590 IN CALCIU um injection, p rmation	<b>Citation</b> D <b>JM</b> bowder, lyophilized, for solution	Date 07/19/2023	Date		
AN LI let P	Category IDA EUCOVOR JCOVORIN Calciu Iroduct Info roduct Type	ANDA216590 IN CALCIU um injection, p rmation	Citation Citation JM bowder, lyophilized, for solution HUMAN PRESCRIPTION DRUG	Date 07/19/2023 Item Code (Sour	<b>ce)</b> NDC:71288-161		
AN LI let P R	Category IDA EUCOVOR JCOVOR JCOVORIN CAICIN TODUCT Info TODUCT Type oute of Admir	ANDA216590 IN CALCIU um injection, p rmation	Citation Citation JM powder, lyophilized, for solution HUMAN PRESCRIPTION DRUG INTRAVENOUS, INTRAMUSCULAR	Date D7/19/2023	ce) NDC:71288-161		
AN let P R	Category IDA EUCOVOR JCOVORIN Calciu Incovorin calciu Incoduct Info roduct Type oute of Admir	ANDA216590 IN CALCIU um injection, p rmation	Citation Citation D D D D D D D D D D D D D D D D D D D	Date D7/19/2023	ce) NDC:71288-161		
AN LI let R	Category IDA EUCOVOR JCOVORIN Calciu Covorin Covorin Calciu Covorin Covorin Calciu Covorin Covorin Calciu Covorin Covorin Calciu Covorin Covori	ANDA216590 IN CALCIU um injection, p rmation histration	Citation Citation D D D D D D D D D D D D D D D D D D D	Date 07/19/2023	ce) NDC:71288-161		
AN LI let P R	Category IDA EUCOVOR JCOVORIN Calciu roduct Info roduct Type oute of Admir	ANDA216590 IN CALCIU um injection, p rmation histration lient/Active Ingred	Citation Citation D D D D D D D D D D D D D D D D D D D	Date D7/19/2023 Item Code (Sour Basis o Strengt	ce) NDC:71288-161		

Inactive Ingredients	
Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
Packaging	
	NAME AND A DESCRIPTION OF THE OWNER

#	Item Code		Package Description	Marketing Start Date	Marketing End Date
1	NDC:71288- 161-20	1 in	1 CARTON	07/19/2023	
1		10 Cor	mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a nbination Product		
P	larketing	In	formation		
	Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
AN	NDA		ANDA216590	07/19/2023	
L	EUCOVOR	IN	CALCIUM		
leı	ucovorin calci	um	injection, powder, lyophilized, for solution	n	
P	roduct Info	rm	ation		

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71288-162
Route of Administration	INTRAVENOUS, INTRAMUSCULAR		

Active Ingredient/Active Moiety						
Ingredient Name	Basis of Strength	Strength				
LEUCOVORIN CALCIUM (UNII: RPR1R4C0P4) (LEUCOVORIN - UNII:Q573I9DVLP)	LEUCOVORIN	200 mg in 20 mL				

Inactive Ingredients					
Ingredient Name	Strength				
SODIUM CHLORIDE (UNII: 451W47IQ8X)					
HYDROCHLORIC ACID (UNII: QTT17582CB)					
SODIUM HYDROXIDE (UNII: 55X04QC32I)					

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71288- 162-30	1 in 1 CARTON	07/19/2023	
1		20 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
ANDA	ANDA216590	07/19/2023	

LEUCOVOR	IN CALCIU	JM				
leucovorin calciu	um injection, p	oowder, lyophilized, for solution	n			
Product Info	rmation					
Product Type		HUMAN PRESCRIPTION DRUG	lten	n Code (Sourc	e)	NDC:71288-163
Route of Admir	nistration	INTRAVENOUS, INTRAMUSCULAR				
Active Ingred	lient/Active	Moiety				
	Ingred	ient Name	Basis of Strength			Strength
LEUCOVORIN CA UNII:Q573I9DVLP)	LCIUM (UNII: RPF	R1R4C0P4) (LEUCOVORIN -	L	EUCOVORIN		350 mg in 17.5 mL
Inactive Ingr	edients					
		Ingredient Name				Strength
SODIUM CHLORI	<b>DE</b> (UNII: 451W47	/IQ8X)				
HYDROCHLORIC	ACID (UNII: QTT)	L7582CB)				
SODIUM HYDROX	(IDE (UNII: 55X04	IQC32I)				
Packaging						
# Item Code	Pa	ackage Description	М	Marketing Star Date		Marketing End Date
<b>1</b> NDC:71288- 163-30	1 in 1 CARTON		07/19/2023			
1	17.5 mL in 1 VIA Combination Pro	L, SINGLE-DOSE; Type 0: Not a oduct				
Marketing	Informat	ion				
Marketing	Annlisa	tion Number or Menegraph	Ma	rkating Start		As ricoting End
Marketing Applica Category		Citation	Citation Date		r	Date
ANDA ANDA21659		0	07/19/2023			
LEUCOVOR leucovorin calciu	IN CALCIU um injection, p	<b>JM</b> bowder, lyophilized, for solution	n			
Product Info	rmation					
Product Type		HUMAN PRESCRIPTION DRUG	lten	n Code (Sourc	e)	NDC:71288-164

Active Ingredient/Active Moiety							
Ingredient Name Basis of Strengt							Strength
LEUCOVORIN CALCIUM (UNII: RPR1R4C0P4) (LEUCOVORIN - UNII:Q573I9DVLP) LEUCOVORIN							500 mg in 50 mL
In	active Ingr	edie	nts				
			Ingredient Name				Strength
sc	DIUM CHLORI	DE (UI	NII: 451W47IQ8X)				
ΗY	DROCHLORIC	ACID	(UNII: QTT17582CB)				
sc	DIUM HYDRO	KIDE (	UNII: 55X04QC32I)				
Pa	ackaging						
#	ltem Code	de Package Description Marketing Sta			arketing Start Date	: <b>M</b>	arketing End Date
1	NDC:71288- 164-50	1288- 0 1 in 1 CARTON 07/19/2023					
1	1 50 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product						
Marketing Information							
	Marketing Category		Application Number or Monograph Citation	Ma	rketing Start Date	М	arketing End Date
AN	DA	А	NDA216590	07/19	/2023		

Labeler - Meitheal Pharmaceuticals Inc. (080548348)

## Establishment

Name	Address	ID/FEI	<b>Business Operations</b>
Kindos Pharmaceuticals Co., LTD		529111185	MANUFACTURE(71288-160, 71288-161, 71288-162, 71288-163, 71288-164)

Revised: 3/2025

Meitheal Pharmaceuticals Inc.