

**SINUS AND COLD D- naproxen sodium, pseudoephedrine hydrochloride tablet, film coated, extended release  
CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED**

-----  
**CVS Pharmacy, Inc. Sinus & Cold-D Drug Facts**

**Active ingredients (in each caplet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*

Pseudoephedrine HCl 120 mg, extended-release

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

Nasal decongestant

**Uses**

temporarily relieves these cold, sinus, and flu symptoms:

- sinus pressure
- minor body aches and pains
- headache
- nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose)
- fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools

- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- redness or swelling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- **swallow whole;** do not crush or chew
- **drink a full glass of water with each dose**
- adults and children 12 years and older: **1 caplet every 12 hours;** do not take more than 2 caplets in 24 hours
- children under 12 years: do not use

### **Other information**

- **each caplet contains:** sodium 21 mg
- meets USP *Dissolution Test 2*
- store at 20-25°C (68-77°F)
- store in a dry place

### **Inactive ingredients**

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

## Questions or comments?

1-800-719-9260

### Package/Label Principal Display Panel

CVS Health®

Compare to the active ingredients in Aleve-D® Sinus & Cold  
Caplets\*\*

Non-Drowsy

Sinus & Cold-D

NAPROXEN SODIUM, 220 mg

PSEUDOEPHEDRINE HYDROCHLORIDE, 120 mg

EXTENDED-RELEASE TABLET

Pain reliever/Fever reducer

(NSAID)

Nasal decongestant

1 CAPLET 12 HOURS

12 Hour multi-symptom relief of:

Sinus pressure

Headache

Sinus congestion

Nasal congestion

Body aches

SINUS & COLD

20 CAPLETS\*\*

\*\*Capsule-shaped tablets

Actual Size

CVS Health.

Caplets\*\*

# Non-Drowsy Sinus & Cold-D



CVS Health.

Compare to the active ingredients  
in Aleve-D® Sinus & Cold†

Caplets\*\*

NDC 51316-339-01

# Non-Drowsy Sinus & Cold-D

NAPROXEN SODIUM, 220 mg  
PSEUDOEPHEDRINE HYDROCHLORIDE, 120 mg  
EXTENDED-RELEASE TABLET

Pain reliever/Fever reducer  
(NSAID)

Nasal decongestant

12 Hour multi-symptom  
relief of:

- Sinus pressure
- Headache
- Sinus congestion
- Nasal congestion
- Body aches



SINUS  
& COLD



20 CAPLETS\*\*

\*\*Capsule-shaped tablets



Actual  
Size

DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS BROKEN

417C9 17 C3

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2023 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
V-14017

#619363

100% money back  
guaranteed.  
CVS.com/returnpolicy



## Drug Facts

### Active ingredients (in each caplet)

Naproxen sodium 220 mg  
(naproxen 200 mg) (NSAID)\*.....Pain reliever/fever reducer  
Pseudoephedrine HCl 120 mg,  
extended-release.....Nasal decongestant  
\*nonsteroidal anti-inflammatory drug

### Purposes

**Uses** temporarily relieves these cold, sinus, and flu symptoms:  
■ sinus pressure ■ minor body aches and pains  
■ headache  
■ nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose)  
■ fever

### Warnings

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed ▶

## Drug Facts (continued)

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ in children under 12 years of age

### Ask a doctor before use if

■ the stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke  
■ you are taking a diuretic  
■ you have problems or serious side effects from taking pain relievers or fever reducers

### Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin  
■ taking any other drug

### When using this product

■ take with food or milk if stomach upset occurs ▼

LIFT  
HERE

LIFT  
HERE

## Drug Facts (continued)

### Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain  
■ trouble breathing  
■ weakness in one part or side of body  
■ slurred speech  
■ legs swelling  
■ redness or swelling is present in the painful area  
■ any new symptoms appear  
■ fever gets worse or lasts more than 3 days  
■ you have difficulty swallowing or the caplet feels stuck in your throat  
■ you get nervous, dizzy, or sleepless  
■ nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). ▶

## Drug Facts (continued)

### Directions

■ do not take more than directed  
■ the smallest effective dose should be used  
■ swallow whole; do not crush or chew  
■ drink a full glass of water with each dose  
■ adults and children 12 years and older: 1 caplet every 12 hours; do not take more than 2 caplets in 24 hours  
■ children under 12 years: do not use

### Other information

■ each caplet contains: sodium 21 mg  
■ meets USP *Dissolution Test 2*  
■ store at 20-25°C (68-77°F)  
■ store in a dry place

**Inactive ingredients** colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyorbate 80, povidone, talc, titanium dioxide

### Questions or comments? 1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve-D® Sinus & Cold.

Glue Area

# SINUS AND COLD D

naproxen sodium, pseudoephedrine hydrochloride tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:51316-339
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NAPROXEN SODIUM</b> (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	L417
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-339-01	20 in 1 CARTON	05/08/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
ANDA	ANDA076518	05/08/2024	

**Labeler** - CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED (062312574)

Revised: 5/2024

CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED