

**CARING MILL IBUPROFEN- ibuprofen tablet, film coated
FSA Store Inc.**

FSA Store Inc. Ibuprofen Tablet Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Ibuprofen Tablets 200 mg

Pain Reliever/Fever Reducer (NSAID)

Actual Size

100 Coated Tablets

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Pain Reliever/Fever Reducer (NSAID)

Ibuprofen Tablets

200 mg

caring mill™



Drug Facts (continued)

Do not use if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer.

Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers, if you have a history of stomach problems, such as heartburn, if you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, if you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease its benefit, aspirin, under a doctor's care for any serious condition, taking any other drug in the painful area, or taking with food or milk if stomach upset occurs.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: **feeling** vomit/blood, **have** bloody or black stools, **have** stomach pain that does not get better, **you** have symptoms of heart problems or stroke, **chest** pain, **trouble** breathing, **weakeness** in one part or side of body, **stained** speech, **leg** swelling, **pain** gets worse or lasts more than 3 days, **redness** or swelling is present in the painful area, **any** new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Do not take more than directed while symptoms persist. **1** tablet, 2 tablets may be used, **do not exceed 6** tablets in 24 hours, unless directed by a doctor.

ask a doctor

children under 12 years

Other information **read all warnings and directions before use.** Store at 20°-25° (68°-77°F) avoiding humidity and excessive heat above 40° (104°F). See end panel for lot number and expiration date.

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, titanium dioxide.

Questions or comments? 1-800-719-9260

Manufactured for FSA Store Inc.
240 W. 37th St. 6th Floor
New York, NY 10018
Customer Care Help Line
1-800-719-9260

KEEP CARTON FOR REFERENCE



3 70030 01420 0

60478 JV C1



Drug Facts

Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)*. *Nonsteroidal anti-inflammatory drug.

Purposes Ibuprofen 200 mg (NSAID)*. *Nonsteroidal anti-inflammatory drug.

Uses temporarily relieves minor aches and pains due to:

- headache
- muscle aches
- minor pain of arthritis
- toothache
- backache
- menstrual cramps
- temporarily reduces fever

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- skin redness
- shock
- skin redness
- asthma (wheeze and cough)
- itching
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take other drugs containing aspirin or non-steroidal anti-inflammatory drugs (NSAIDs) (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.



NDC 81522-604-78

 caring mill™

Ibuprofen Tablets

200 mg

Pain Reliever/Fever Reducer (NSAID)



Actual Size

100 Coated Tablets



CARING MILL IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81522-604
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81522-604-71	1 in 1 CARTON	07/27/2021	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:81522-604-78	1 in 1 CARTON	07/27/2021	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:81522-604-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/29/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	07/27/2021	

Labeler - FSA Store Inc. (049283340)

Revised: 7/2021

FSA Store Inc.