

## **NYSTATIN- nystatin suspension** **Bryant Ranch Prepack**


-----

### **NYSTATIN ORAL SUSPENSION, USP**

#### **Rx Only**

#### **DESCRIPTION**

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*.  
Structural formula:



Nystatin Oral Suspension, USP, for oral administration contains 100,000 Nystatin units per mL. In addition, the yellow opaque suspension contains the following inactive ingredients : Alcohol (0.5% v/v), USP, Alcohol free Bubble Gum Flavoring, Carboxymethylcellulose Sodium, USP, Dibasic Sodium Phosphate, USP, Glycerin Natural 99.5%, USP, Methylparaben, NF, (Preservative), Monobasic Sodium Phosphate, USP, Propylparaben, NF, (Preservative), Purified Water, USP, Saccharin Sodium, USP, and Sucrose, NF.

#### **CLINICAL PHARMACOLOGY**

##### **Pharmacokinetics**

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

##### **Microbiology**

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida* species become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

## **INDICATIONS AND USAGE**

Nystatin Oral Suspension, USP, is indicated for the treatment of candidiasis in the oral cavity.

## **CONTRAINDICATIONS**

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

## **PRECAUTIONS**

### **General**

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

### **Pregnancy**

Teratogenic Effects

#### *Category C*

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

### **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

### **Pediatric Use**

See **DOSAGE AND ADMINISTRATION**.

## **ADVERSE REACTIONS**

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General**).

*Gastrointestinal:* Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

*Dermatologic:* Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

*Other:* Tachycardia, bronchospasm, facial swelling, and nonspecific myalgia have also been rarely reported.

To report SUSPECTED ADVERSE EVENTS, contact FDA at 1-800-FDA-1088 or [www.fda.gov](http://www.fda.gov).

## **OVERDOSAGE**

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects or superinfections (see **CLINICAL PHARMACOLOGY: Pharmacokinetics**).

## **DOSAGE AND ADMINISTRATION**

### **Infants**

2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

NOTE: Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

### **Children and Adults**

4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

## **HOW SUPPLIED**

Nystatin Oral Suspension, USP, 100,000 Nystatin units per mL, is available as a bubble gum flavored, yellow opaque, ready-to-use suspension.

Nystatin Oral Suspension, USP, is available as follows:

NDC: 63629-8519-1: 16 fl. oz. bottle (480 mL).

## **Storage**

Store at 20°C to 25°C (68° to 77°F) [See USP Controlled Room Temperature]

AVOID FREEZING

**Rx Only**

**Manufactured By:**

**VistaPharm®**

Largo, FL 33771

VP1086R2

06/19

**Nystatin 100000 Suspension, #480**



Each 1 mL contains: 100,000 units Nystatin, USP. Contains: Alcohol 0.5% v/v (Bubblegum Flavored)

Keep this and all medication out of the reach of children.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Avoid Freezing.

Dispense in tight, light-resistant containers as defined in the USP.

Shake well before using.

**NDC 63629-8519-1**

**Nystatin Oral Suspension, USP**

**100,000 units/mL**



Relabeled by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504 USA

Rx only  
**16 fl. oz. (480 mL)**

Manufactured by:  
VistaPharm, Inc.



## NYSTATIN

nystatin suspension

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63629-8519(NDC:66689-008)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USP'U] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	

<b>SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM</b> (UNII: GR686LBA74)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SUCROSE</b> (UNII: C151H8M554)	

### Product Characteristics

<b>Color</b>	YELLOW (Light yellow)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629-8519-1	480 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/19/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA064142	05/01/2012	

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

### Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(63629-8519) , RELABEL(63629-8519)

Revised: 6/2023

Bryant Ranch Prepack