

ASPIRIN- aspirin tablet
Cardinal Health

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Aspirin 325 mg

Hospak® Unit Dose Products

Drug Facts

ASPIRIN 325 mg (5gr) (NSAID)*

*nonsteroidal anti-inflammatory drug

MICRO-COATED TABLETS

Active ingredient

Aspirin 325 mg (in each tablet)

Purpose

Pain reliever

Uses

-for the temporary relief of minor aches and pains due to:

- Headache
- Colds
- Menstrual pain
- Muscle pain
- Toothache
- Minor pain of arthritis
- Or as directed by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- Hives
- Facial swelling
- Shock
- Asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach

bleeding. The chance is higher if you

- Are age 60 or older
- Have had stomach ulcers or bleeding problems
- Take a blood thinning (anticoagulant) or steroid drug
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed
- Take other drugs containing prescription or nonprescription NSAIDS (aspirin, ibuprofen, naproxen, or others)

Do not use - if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- Stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn
- You have asthma
- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You are taking a diuretic
- You have not been drinking fluids
- You have lost a lot of fluid due to vomiting or diarrhea

Ask a doctor or pharmacist before use if you are

- Taking a prescription drug for diabetes, gout or arthritis
- Taking any other drug
- Under a doctor's care for any serious condition

Warning:

This Unit Dose package is not child resistant and is Intended for Institutional Use Only.

Stop use and ask a doctor if

-you experience any of the following signs of stomach bleeding

- Feel faint
- Have bloody or black stools
- Vomit blood
- have stomach pain that does not get better
- pain get worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Directions

Drink a full glass of water with each dose

Adults and children 12 years and over:

Take 1 to 2 tablets every 4 hours while symptoms last.

Do not take more than 12 tablets in 24 hours unless directed by a doctor

Children under 12 years: consult a doctor

Keep Out of Reach of Children

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Other Information

- Do not use if packet is torn, cut or opened.
- Store at 25C (77F) excursions permitted between 15-30C (59-86F)
- use by expiration date on package

Mfg. by: Time-Cap Labs Inc., Farmingdale, NY 11735

Dist. By: Hospak® Unit Dose Products, Huntley, IL 60142

Rev. 08/17

Distributed by:

Cardinal Health

Dublin, OH 43017

L42652450218A

Principal Display Panel

Aspirin 325 mg

10 Micro-Coated Tablets



NDC 55154-6755-0

R76

ASPIRIN 325mg (5gr)

10 MICRO-COATED TABLETS

Drug Facts

Active Ingredient (in each tablet)

Aspirin 325mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

See product insert for uses, directions, prescribing information, precautions and warnings.

Inactive Ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

STORAGE: Store at 25° C (77° F) excursions permitted between 15° - 30° C (59° - 86° F)

- do not use if packet is torn, cut or opened

- use by expiration date on package

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.

When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Do not use - if you are allergic to aspirin or any other pain reliever/fever reducer

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Mfg. by: Time-Cap Labs Inc., Farmingdale, NY 11735

Dist. By: Hospak® Unit Dose Products, Huntley, IL 60142

Distributed by Cardinal Health

Dublin, OH 43017

L42652450218A

LOT #: XXXXXXXXXX EXP. DATE: XX/XX/XX

ASPIRIN

aspirin tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:55154-6755(NDC:66553-001)

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-6755-0	10 in 1 BAG	06/02/2011	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		06/02/2011	

Labeler - Cardinal Health (603638201)

Revised: 5/2019

Cardinal Health