

VALUMEDS ASPIRIN- aspirin tablet, delayed release
SPIRIT PHARMACEUTICALS LLC

ASPIRIN 325 MG TABLETS

Active ingredient (in each tablet)

Aspirin (NSAID)* 325 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

- temporary relieves minor aches and pains due to:
- headache
- minor arthritis pain
- toothache
- menstrual pain
- colds
- or as recommended by a doctor

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding.

The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDS(aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Do not use

if you have ever had an allergic reaction to aspirin or any other pain reliever/fever

reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- allergic reaction occurs, Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present
 - any new symptom appear
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.
1(800)222-1222

Directions

- adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist. Drink a full glass of water with each dose.
- do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: ask a doctor

Other information

- store below 25 °C (77 °F)
- Tamper Evident Feature: Do not use if printed inner-seal beneath cap is missing or broken

corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Questions or comments?**1(888)333-9792****VALUMEDS™**

Compare to the active ingredient in

ECOTRIN® TABLETS*

REGULAR STRENGTH

ASPIRIN

PAIN RELIEVER (NSAID)

- SAFETY COATED

PROTECT AGAINST

STOMACH UPSET

500 TABLETS



Compare to the active ingredient in
ECOTRIN® TABLETS*

REGULAR STRENGTH

ASPIRIN

PAIN RELIEVER (NSAID)

SAFETY COATED

- PROTECTS AGAINST STOMACH UPSET



500 TABLETS

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Aspirin (NSAID)¹ 325 mg. ¹nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- Temporarily relieves minor aches and pains due to:
 - headache
 - minor arthritis pain
 - toothache
 - menstrual pain
 - colds
- or as recommended by a doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, shock, facial swelling, asthma (wheezing)

Stomach bleeding warning:

- This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

Do not use

If you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis

(continued under label)



ITEM#60531-4

0 40232 60531 4

Distributed By:
Spirit Pharmaceuticals, LLC
Ronkonkoma, NY 11779
ORIG 06/17

Lot No.

Exp. Date



Drug Facts (continued)

Stop use and ask a doctor if

- allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - belly pain
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1 (800) 222-1222

Directions

- adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist. Drink a full glass of water with each dose.
- do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: ask a doctor

Other information

- store below 25°C (77°F)
- Tamper Evident Feature: Do not use if printed inner-seal beneath cap is missing or broken.

Inactive ingredients

corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1(888)333-3792

*This product is not manufactured or distributed by Prestige Brands, Inc. owner of the registered trademark Ecotrin®

VALUMEDS ASPIRIN

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68210-0025
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3S)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

HYPROMELLOSES (UNII: 3NXW29V3WO)
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
MINERAL OIL (UNII: T5L8T28FGP)
POLYSORBATE 80 (UNII: 6OZP39ZG8H)
DIMETHICONE (UNII: 92RU3N3Y1O)
SODIUM HYDROXIDE (UNII: 55X04QC32I)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	T
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-0025-0	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/17/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/17/2018	

Labeler - SPIRIT PHARMACEUTICALS LLC (179621011)

Revised: 12/2023

SPIRIT PHARMACEUTICALS LLC