

FREEGELLS COUGH DROPS MENTHOL CHERRY 80CT- menthol lozenge
RL Albert & Son Inc.

Drug Facts

Active ingredient (in each drop)

Menthol 5.8 mg

Purpose

Cough suppressant/Oral anesthetic

Uses

- temporarily relieves:
- cough as may occur with a cold
- occasional minor irritation and sore throat

Warnings

Sore throat warning:if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly. These symptoms may be serious.

Ask a doctor before use if you have:

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough accompanied by excessive phlegm (mucus)

Stop use and ask a doctor if

- cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache, These could be signs of a serious condition.
- sore throat is severe, or irritation, pain or redness lasts or worsens
- sore mouth does not improve in 7 days

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children.

Directions

adults and children 5 years and over dissolve 1 lozenge slowly in mouth. Repeat every 2 hours as needed or as directed by a doctor.

children under 5 years ask a doctor

Other information

- 11 calories per drop
- do not use if bags is torn or open
- **Contains: Soy**

Inactive ingredients

Corn syrup, eucalyptus oil, FD&C red no.40, natural & artificial flavors, soy lecithin, sucrose and water.

Product label

RESALABLE PAK

TEAR HERE ▶

FREEGELLS COUGH DROPS

Menthol 5mg per drop

SOOTHES SORE THROATS
TEMPORARILY RELIEVES COUGHS

COUGH SUPPRESSANT
ORAL ANESTHETIC

MENTHOL CHERRY

80 DROPS

0 1 2 3 4 5 6 7 8 9

Made in Brazil by Aldin S.A.

Distributed by:
M.J. Hunt & Son, Inc.,
Stamford, CT 06902
Sole Importers U.S.A.

Drug Facts

Warnings
Do not use if you are allergic to any of the ingredients listed above or to any of the following: Menthol, Cherry, or other ingredients. Stop use and ask doctor if you have a history of asthma or other breathing problems. Stop use and ask doctor if you have a history of heart disease or other heart problems. Stop use and ask doctor if you have a history of low blood pressure or other blood pressure problems. Stop use and ask doctor if you have a history of dizziness or other dizziness problems. Stop use and ask doctor if you have a history of fainting or other fainting problems. Stop use and ask doctor if you have a history of seizures or other seizure problems. Stop use and ask doctor if you have a history of diabetes or other diabetes problems. Stop use and ask doctor if you have a history of kidney disease or other kidney disease problems. Stop use and ask doctor if you have a history of liver disease or other liver disease problems. Stop use and ask doctor if you have a history of stomach or intestinal disease or other stomach or intestinal disease problems. Stop use and ask doctor if you have a history of pancreas disease or other pancreas disease problems. Stop use and ask doctor if you have a history of gallbladder disease or other gallbladder disease problems. Stop use and ask doctor if you have a history of thyroid disease or other thyroid disease problems. Stop use and ask doctor if you have a history of parathyroid disease or other parathyroid disease problems. Stop use and ask doctor if you have a history of adrenal disease or other adrenal disease problems. Stop use and ask doctor if you have a history of pituitary disease or other pituitary disease problems. Stop use and ask doctor if you have a history of hypothalamic disease or other hypothalamic disease problems. Stop use and ask doctor if you have a history of pineal disease or other pineal disease problems. Stop use and ask doctor if you have a history of brain disease or other brain disease problems. Stop use and ask doctor if you have a history of spinal cord disease or other spinal cord disease problems. Stop use and ask doctor if you have a history of nerve disease or other nerve disease problems. Stop use and ask doctor if you have a history of muscle disease or other muscle disease problems. Stop use and ask doctor if you have a history of bone disease or other bone disease problems. Stop use and ask doctor if you have a history of joint disease or other joint disease problems. Stop use and ask doctor if you have a history of skin disease or other skin disease problems. Stop use and ask doctor if you have a history of eye disease or other eye disease problems. Stop use and ask doctor if you have a history of ear disease or other ear disease problems. Stop use and ask doctor if you have a history of nose disease or other nose disease problems. Stop use and ask doctor if you have a history of throat disease or other throat disease problems. Stop use and ask doctor if you have a history of mouth disease or other mouth disease problems. Stop use and ask doctor if you have a history of tongue disease or other tongue disease problems. Stop use and ask doctor if you have a history of salivary gland disease or other salivary gland disease problems. Stop use and ask doctor if you have a history of lymphatic system disease or other lymphatic system disease problems. Stop use and ask doctor if you have a history of circulatory system disease or other circulatory system disease problems. Stop use and ask doctor if you have a history of respiratory system disease or other respiratory system disease problems. Stop use and ask doctor if you have a history of digestive system disease or other digestive system disease problems. Stop use and ask doctor if you have a history of urinary system disease or other urinary system disease problems. Stop use and ask doctor if you have a history of reproductive system disease or other reproductive system disease problems. Stop use and ask doctor if you have a history of endocrine system disease or other endocrine system disease problems. Stop use and ask doctor if you have a history of immune system disease or other immune system disease problems. Stop use and ask doctor if you have a history of nervous system disease or other nervous system disease problems. Stop use and ask doctor if you have a history of muscular system disease or other muscular system disease problems. Stop use and ask doctor if you have a history of skeletal system disease or other skeletal system disease problems. Stop use and ask doctor if you have a history of integumentary system disease or other integumentary system disease problems. Stop use and ask doctor if you have a history of sensory system disease or other sensory system disease problems. Stop use and ask doctor if you have a history of motor system disease or other motor system disease problems. Stop use and ask doctor if you have a history of reproductive system disease or other reproductive system disease problems. Stop use and ask doctor if you have a history of endocrine system disease or other endocrine system disease problems. Stop use and ask doctor if you have a history of immune system disease or other immune system disease problems. Stop use and ask doctor if you have a history of nervous system disease or other nervous system disease problems. Stop use and ask doctor if you have a history of muscular system disease or other muscular system disease problems. Stop use and ask doctor if you have a history of skeletal system disease or other skeletal system disease problems. Stop use and ask doctor if you have a history of integumentary system disease or other integumentary system disease problems. Stop use and ask doctor if you have a history of sensory system disease or other sensory system disease problems. Stop use and ask doctor if you have a history of motor system disease or other motor system disease problems.

Directions
Adults: 1 lozenge, 4 to 6 times a day, as needed for cough or throat irritation. Children: 1/2 lozenge, 4 to 6 times a day, as needed for cough or throat irritation. Do not use for more than 7 days. If symptoms persist, consult your doctor.

Other Information
Keep out of reach of children. Do not use if the seal is broken. Do not use if the product is expired. Do not use if the product is damaged. Do not use if the product is contaminated. Do not use if the product is recalled.

Contains 80 Drops

FREEGELLS COUGH DROPS

FREEGELLS COUGH DROPS MENTHOL CHERRY 80CT

menthol lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83002-037
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEVOMENTHOL (UNII: BZ1R15MTK7) (LEVOMENTHOL - UNII:BZ1R15MTK7)	LEVOMENTHOL	5.8 mg

Inactive Ingredients

Ingredient Name	Strength
SUCROSE (UNII: C151H8M554)	
CORN SYRUP (UNII: 9G5L16BK6N)	
EUCALYPTOL (UNII: RV6J6604TK)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor	CHERRY	Imprint Code	F
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83002-037-01	6 in 1 CARTON	03/15/2024	
1		80 in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/15/2024	

Labeler - RL Albert & Son Inc. (005900899)

Revised: 3/2024

RL Albert & Son Inc.