

**SUNMARK LOPERAMIDE HYDROCHLORIDE- loperamide hcl suspension**  
**McKesson**

-----  
**McKesson Loperamide Hydrochloride Oral Suspension Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease

**Ask a doctor or pharmacist before use if you are**

taking antibiotics

**When using this product**

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours
children 6-8 years (48-59 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

## Other information

- each 30 mL (6 tsp) contains: sodium 15 mg
- store between 20-25°C (68-77°F)
- see side panel for lot number and expiration date

## Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

## Questions or comments?

1-800-719-9260

## Package/Label Principal Display Panel

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT

loperamide hydrochloride oral suspension

Anti-Diarrheal

CONTROLS THE SYMPTOMS OF DIARRHEA

GLUTEN FREE

MINT FLAVOR

4 FL OZ (120 mL)

1 mg Loperamide Hydrochloride per 7.5 mL

**Drug Facts**

**Active ingredient Purpose (in each 7.5 mL)**  
Loperamide HCl  
1 mg.....Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use** if you have bloody or black stool

**PEEL BACK HERE**

**sunmark®**

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT\*  
NDC 49348-999-34

**loperamide hydrochloride oral suspension**

**Anti-Diarrheal**

**CONTROLS THE SYMPTOMS OF DIARRHEA**

**GLUTEN FREE**

**MINT FLAVOR**

**4 FL OZ (120 mL)**

1 mg Loperamide Hydrochloride per 7.5 mL

**Do not use if printed plastic neckband is broken or missing.**

**McKesson**

Another Quality Product  
Distributed By McKesson  
One Post Street,  
San Francisco, CA 94104  
Money Back Guarantee  
Please visit us at  
www.sunmarkbrand.com

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.



: 64526 S1 F2

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool ■ a history of liver disease

**Ask a doctor or pharmacist before use if you are taking antibiotics**

**When using this product** ■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

■ symptoms get worse  
■ diarrhea lasts for more than 2 days  
■ you get abdominal swelling or bulging.  
These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison

**Drug Facts (continued)**

Control Center right away. (1-800-222-1222)

**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ shake well before using ■ only use attached measuring cup to dose product

adults and children 12 years and over 30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours

children 9-11 years (60-95 lbs) 15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours

**Drug Facts (continued)**

children 6-8 years (48-59 lbs) 15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours

children under 6 years (up to 47 lbs) ask a doctor

**Other information**

■ each 30 mL (6 tsp) contains: sodium 15 mg  
■ store between 20-25°C (68-77°F)

**Drug Facts (continued)**

■ see side panel for lot number and expiration date

**Inactive ingredients** anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

**Questions or comments?**  
1-800-719-9260

# SUNMARK LOPERAMIDE HYDROCHLORIDE

loperamide hcl suspension

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-999
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	

## Product Characteristics

Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-999-34	120 mL in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091292	02/18/2014	

