

**DAYTIME SEVERE COLD AND COUGH- acetaminophen, dextromethorphan hbr,  
phenylephrine hcl liquid  
H E B**

-----

**Drug Facts**

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves these common cold due to
  - minor aches and pains
  - headache
  - nasal and sinus congestion
  - sore throat
  - cough due to minor throat and bronchial irritation
  - temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blister
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if user is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

## Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses (180 mL) in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL every 4 hours
- children under 12 years of age: do not use

## Other information

- each 30 mL contains: sodium 16 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1 FD&C red #40, , flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate sodium citrate

## Questions or comments?

Call toll free **1-877-753-3935** Monday-Friday 9AM-5PM EST

## Principal Display Panel

### **Compare to Theraflu® ExpressMax® Daytime Severe Cold & Cough active ingredients\***

DAYTIME

#### **Severe Cold & Cough**

Acetaminophen 650 mg /

Pain Reliever/Fever Reducer

Dextromethorphan HBr 20 mg /

Cough Suppressant

Phenylephrine HCl 10 mg/

Nasal Decongestant

Flu & Severe Cold

- Nasal Congestion
- Fever
- Cough

- Sore Throat
- Headache
- Body Ache

Alcohol 10%

For Ages 12 Years & Over

Berry Flavor

FL OZ (mL)

\*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® ExpressMax® Daytime Severe Cold & Cough.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

**Product Label**

Compare to **Theraflu® ExpressMax® Daytime Severe Cold & Cough** active ingredients\*

NDC 37808-718-83

**H-E-B**

DAYTIME  
**Severe Cold & Cough**

**Acetaminophen 650 mg /**  
Pain Reliever/Fever Reducer  
**Dextromethorphan HBr 20 mg /**  
Cough Suppressant  
**Phenylephrine HCl 10 mg /**  
Nasal Decongestant

**Flu & Severe Cold**

- Nasal Congestion • Fever • Cough
- Sore Throat • Headache • Body Ache

Alcohol 10%  
For Ages 12 Years & Over

**Berry Flavor**

PLD-A398B LB009106  
**8.3 FL OZ (245 mL)**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® ExpressMax® Daytime Severe Cold & Cough.

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

**H-E-B**

**100% GUARANTEE promise**

If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

PLD-A398B  
LB009107  
515225



**Drug Facts**

**Active ingredients (in each 30 mL)**      **Purposes**

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Phenylephrine HCl 10 mg.....Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Uses**

- temporarily relieves these symptoms due to a cold
  - minor aches and pains
  - nasal and sinus congestion
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever
- headache
- sore throat

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Drug Facts (continued)**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see **Overdose warning**)
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 30 mL contains: sodium 16 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

**Questions or comments?** Call toll free 1-877-753-3935 Monday-Friday 9AM-5PM EST

**H-E-B Daytime Severe Cold & Cough Berry Flavor**

**DAYTIME SEVERE COLD AND COUGH**

acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-718
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

Product Characteristics			
<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-718-83	245 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2023	

**Labeler - H E B (007924756)**