

**GUAIFENESIN AND PSEUDOEPHEDRINE HCL- guaifenesin and pseudoephedrine hcl tablet, extended release**  
**Rite Aid Corporation**

-----  
**Drug Facts**

**Active ingredient(s)**

Guaifenesin USP, 600  
mg.....Expectorant

Pseudoephedrine HCl USP, 60 mg.....Nasal  
Decongestant

Guaifenesin USP, 1200  
mg.....Expectorant

Pseudoephedrine HCl USP, 120 mg.....Nasal  
Decongestant

**Purpose**

Expectorant and Nasal Decongestant

**Use(s)**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bother some mucus and make coughs more productive
- temporarily relieves nasal congestion due to:
  - common cold
  - hay fever
  - upper respiratory allergies
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- temporarily relieves sinus congestion and pressure\_

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

### **When using this product**

- do not use more than directed

### **Stop use and ask doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

### **Pregnancy/Breastfeeding**

ask a health professional before use.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- children under 12 years of age: do not use
- adults and children 12 years and older: 2 tablets every 12 hours; not more than 4 tablets in 24 hours

### **Other information**

- tamper evident: do not use if carton is open or if printed seal on blister is broken or missing.

### **Storage**

■ store between 20°-25°C (68°-77°F)

### **Inactive ingredients**

FD & C yellow #6, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone, silicon dioxide.

### **Questions**

call **1-888-375-3784**

You may also report side effect to this phone number.

**Keep the carton.**

**It contains important information.**

**See end panel for expiration date.**

Manufactured by:

**Dr. Reddy's Laboratories Limited**

Srikakulam - 532 409 INDIA

### **Principal Display Panel**

**MUCUS RELIEF D**  
GUAIFENESIN 600 mg & PSEUDOEPHEDRINE HCl 60 mg  
Extended-Release Tablets  
EXPECTORANT & NASAL DECONGESTANT

**MUCUS RELIEF D**  
GUAIFENESIN 600 mg & PSEUDOEPHEDRINE HCl 60 mg  
Extended-Release Tablets  
EXPECTORANT & NASAL DECONGESTANT

18  
EXTENDED-RELEASE  
TABLETS

12  
HOUR

Compare to the active ingredients in Mucinex<sup>®</sup> D Extended-Release Tablets.\*

NDC 11822-0046-1

Keep the carton. It contains important information. See end panel for expiration date.

\*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Mucinex<sup>®</sup>.

0 11822 69042 3

150087835

150087835

150087835

EXP

LOT

REV 10/21

MADE IN INDIA

DISTRIBUTED BY: RITE AID  
30 HUNTER LANE, CAMP HILL, PA 17011  
www.riteaid.com

DO NOT USE IF CARTON IS OPEN OR IF PRINTED SEAL ON BLISTER IS BROKEN OR MISSING

**Drug Facts** (continued)

**When using this product**

- do not use more than directed

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 2 tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

**Other information**

- do not use if carton is open or if printed seal on blister is broken or missing
- store at 20° - 25°C (68° - 77°F)

**Inactive ingredients**

FD & C yellow #6, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polydioxane, silicon dioxide.

**Questions? call 1-888-375-3784**

You may also report side effects to this phone number

## GUAIFENESIN AND PSEUDOEPHEDRINE HCL

guaifenesin and pseudoephedrine hcl tablet, extended release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0046(NDC:55111-798)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Guaifenesin</b> (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	600 mg
<b>Pseudoephedrine Hydrochloride</b> (UNII: 6V9V2RYJ8N) (Pseudoephedrine - UNII:7CUC9DDI9F)	Pseudoephedrine Hydrochloride	60 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	RDY;798
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0046-1	1 in 1 CARTON	02/01/2022	
1		18 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0046-2	2 in 1 CARTON	02/01/2022	
2		18 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208369	02/01/2022	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 8/2022

Rite Aid Corporation