

**ACID REDUCER- ranitidine hydrochloride tablet, film coated**  
**SAFEWAY INC.**

-----  
**Drug Facts**

**ACTIVE INGREDIENT (IN EACH TABLET)**

Ranitidine 75 mg (as ranitidine hydrochloride, USP 84 mg)

**PURPOSE**

Acid reducer

**USES**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**WARNINGS**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do Not Use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

Ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**DIRECTIONS**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

#### **OTHER INFORMATION**

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

#### **INACTIVE INGREDIENTS**

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

#### **QUESTIONS?**

Call 1-800-406-7984

#### **PRINCIPAL DISPLAY PANEL**

**Regular Strength**

**NDC 21130-003-30**

**Compare to Zantac 75<sup>®</sup> active ingredient<sup>†</sup>**

**Acid Reducer**

**Ranitidine Tablets, USP 75 mg Acid Reducer**

- **Prevents and Relieves Heartburn Associated with Acid Indigestion and Sour Stomach**



**30 TABLETS**

**(30 DOSES)**

**SAFEWAY<sup>®</sup>**

**DISTRIBUTED BY SAFEWAY INC.**

**5104762/0813**



NDC 21130-003-30

Compare to Zantac 75® active ingredient†

Regular Strength

# Acid Reducer

Ranitidine Tablets, USP 75mg  
Acid Reducer

- Prevents and Relieves Heartburn Associated with Acid Indigestion and Sour Stomach

30 TABLETS (30 DOSES)

SAFEWAY®

ACTUAL SIZE

**Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.**

†This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Zantac 75® Tablets. Zantac 75® is a registered trademark of Boehringer Ingelheim Pharmaceuticals, Inc.

0813  
DISTRIBUTED BY SAFEWAY INC.  
P.O. BOX 99, PLEASANTON, CA 94566-0099  
1-888-SAFEWAY www.safeway.com

MADE IN INDIA  
DRUGS/PV05 13 1523




# Acid Reducer

Ranitidine Tablets, USP 75mg/Acid Reducer

NON VARNISH AREA

020002466

5104762



5104762

<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>your heartburn continues or worsens</li> <li>you need to take this product for more than 14 days</li> <li>if pregnant or breast-feeding, ask a health professional before use.</li> </ul> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	
<p><b>Directions</b></p> <ul style="list-style-type: none"> <li>adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water</li> <li>to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn</li> <li>can be used up to twice daily (do not take more than 2 tablets in 24 hours)</li> <li>children under 12 years: ask a doctor</li> </ul>	
<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.</li> <li>store at 20° - 25° C (68° - 77° F)</li> <li>avoid excessive heat or humidity</li> <li>this product is sodium and sugar free</li> </ul>	
<p><b>Inactive ingredients</b> colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide</p> <p><b>Questions?</b> call 1-800-406-7984</p>	
<p><b>Drug Facts</b></p> <p><b>Active ingredient</b> (in each tablet) Ranitidine 75 mg.....Acid reducer (as ranitidine hydrochloride, USP 84 mg)</p>	<p><b>Uses</b></p> <ul style="list-style-type: none"> <li>relieves heartburn associated with acid indigestion and sour stomach</li> <li>prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages</li> </ul> <p><b>Warnings</b></p> <p><b>Allergy alert:</b> Do not use if you are allergic to ranitidine or other acid reducers</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</li> <li>with other acid reducers</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>chest pain</li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> </ul> <p><b>heartburn over 3 months.</b> This may be a sign of a more serious condition.</p> <p><b>heartburn with lightheadedness, sweating or dizziness</b></p> <ul style="list-style-type: none"> <li>chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness</li> </ul>

## ACID REDUCER

ranitidine hydrochloride tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-003
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDRO CHLO RIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	75 mg

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

SILICON DIO XIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	OR;606
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-003-30	1 in 1 CARTON		
1		30 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201745	12/13/2013	

**Labeler** - SAFEWAY INC. (009137209)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

### Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(21130-003)