


CORDRAN- flurandrenolide cream
CORDRAN- flurandrenolide ointment
Almirall, LLC

**Cordran® Cream
and
Cordran® Ointment
Flurandrenolide, USP**

DESCRIPTION

Cordran® (flurandrenolide, USP) is a potent corticosteroid intended for topical use. Flurandrenolide occurs as white to off-white, fluffy, crystalline powder and is odorless. Flurandrenolide is practically insoluble in water and in ether. One gram of flurandrenolide dissolves in 72 mL of alcohol and in 10 mL of chloroform. The molecular weight of flurandrenolide is 436.52.

The chemical name of flurandrenolide is Pregn-4-ene-3,20-dione, 6-fluoro-11,21-dihydroxy-16,17-[(1-methylethylidene)bis (oxy)]-, (6 α , 11 β , 16 α)-; its empirical formula is C₂₄H₃₃FO₆. The structure is as follows:



Each gram of Cordran® Cream (flurandrenolide Cream, USP) contains 0.5 mg (1.145 μ mol; 0.05%) or 0.25 mg (0.57 μ mol; 0.025%) flurandrenolide in an emulsified base composed of cetyl alcohol, citric acid, mineral oil, polyoxyl 40 stearate, propylene glycol, sodium citrate, stearic acid, and purified water.

Each gram of Cordran® Ointment (flurandrenolide Ointment, USP) contains 0.5 mg (1.145 μ mol; 0.05%) or 0.25 mg (0.57 μ mol; 0.025%) flurandrenolide in a base composed of white wax, cetyl alcohol, sorbitan sesquioleate, and white petrolatum.

CLINICAL PHARMACOLOGY

Cordran is primarily effective because of its anti-inflammatory, antipruritic, and vasoconstrictive actions.

The mechanism of the anti-inflammatory effect of topical corticosteroids is not completely understood. Corticosteroids with anti-inflammatory activity may stabilize cellular and lysosomal membranes. There is also the suggestion that the effect on the membranes of lysosomes prevents the release of proteolytic enzymes and, thus, plays a part in reducing inflammation.

Pharmacokinetics— The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings.

Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption.

Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. They are metabolized primarily in the liver and then excreted in the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

Cordran® (flurandrenolide, USP) is indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.

CONTRAINDICATIONS

Topical corticosteroids are contraindicated in patients with a history of hypersensitivity to any of the components of these preparations.

PRECAUTIONS

General—Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

Conditions that augment systemic absorption include application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings.

Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression using urinary-free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid.

Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, so that supplemental systemic corticosteroids are required.

Pediatric patients may absorb proportionately larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (see Pediatric Use under

PRECAUTIONS).

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted.

In the presence of dermatologic infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, Cordran should be discontinued until the infection has been adequately controlled.

Information for the Patient—Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped in order to be occlusive unless the patient is directed to do so by the physician.
4. Patients should report any signs of local adverse reactions, especially under occlusive dressing.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a patient being treated in the diaper area, because these garments may constitute occlusive dressings.
6. Do not use Cordran on the face, underarms, or groin areas unless directed by your physician.
7. If no improvement is seen within 2 weeks, contact your physician.
8. Do not use other corticosteroid-containing products while using Cordran without first consulting your physician.

Laboratory Tests—The following tests may be helpful in evaluating the HPA axis suppression:

Urinary-free cortisol test
ACTH stimulation test

Carcinogenesis, Mutagenesis, and Impairment of Fertility—Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids.

Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Usage in Pregnancy — Pregnancy Category C—Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively for pregnant patients or in large amounts or for prolonged periods of time.

Nursing Mothers—It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities *not likely* to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use—Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than do mature patients because of a larger skin surface area to body weight ratio.

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in pediatric patients receiving topical corticosteroids. Manifestations of adrenal suppression in pediatric patients include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to pediatric patients should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of pediatric patients.

ADVERSE REACTIONS

The following local adverse reactions are reported infrequently with topical corticosteroids but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence:

- Burning
- Itching
- Irritation
- Dryness
- Folliculitis
- Hypertrichosis
- Acneform eruptions
- Hypopigmentation
- Perioral dermatitis
- Allergic contact dermatitis

The following may occur more frequently with occlusive dressings:

- Maceration of the skin
- Secondary infection
- Skin atrophy
- Striae
- Miliaria

Postmarketing Adverse Reactions

The following adverse reactions have been identified during post approval use of flurandrenolide, USP. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Skin: skin striae, hypersensitivity, skin atrophy, contact dermatitis and skin discoloration.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

DOSAGE AND ADMINISTRATION

For moist lesions, a small quantity of the cream should be rubbed gently into the affected areas 2 or 3 times a day. For dry, scaly lesions, the ointment is applied as a thin film to affected areas 2 or 3 times daily.

Therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, reassessment of the diagnosis may be necessary.

Cordran® (flurandrenolide, USP) should not be used with occlusive dressings unless directed by a physician. Tight-fitting diapers or plastic pants may constitute occlusive dressings.

HOW SUPPLIED

Cordran® Cream is supplied in aluminum tubes as follows:

Cordran® Cream, 0.025%:

120 g (NDC 16110-034-12)

Cordran® Cream, 0.05%:

120 g (NDC 16110-035-12)

Cordran® Ointment is supplied in aluminum tube as follows:

Cordran® Ointment, 0.05%:

60 g (NDC 16110-026-60)

120 g (NDC 16110-026-12)

Keep out of reach of children.

Storage

Keep tightly closed.

Protect from light.

Store at 20° to 25°C (68° to 77°F) with excursions permitted to 15° to 30°C (59° to 86°F) [See USP controlled room temperature.].


Rx Only

Revision: May 2017

ALMIRALL, LLC

Manufactured by DPT Laboratories, San Antonio, TX 78215
For Almirall, LLC, 101 Lindenwood Drive, Suite 400, Malvern, PA 19355

PRINCIPAL DISPLAY PANEL - NDC: 16110-035-12 - 0.05% Cream 120 g Tube Label



PRINCIPAL DISPLAY PANEL - NDC: 16110-035-12 - 0.05% Cream 120 g Carton Label



PRINCIPAL DISPLAY PANEL - NDC: 16110-034-12 - 0.025% Cream 120 g Tube Label

Usual Dosage: Apply to affected area 2 or 3 times a day. See package insert.

Each gram contains: flurandrenolide 0.25 mg (0.025%), cetyl alcohol, citric acid, mineral oil, polyoxyxl 40 stearate, propylene glycol, sodium citrate, stearic acid and purified water.

Store at 20° to 25°C (68° to 77°F) with excursions permitted to 15° to 30°C (59° to 86°F) [See USP controlled room temperature].

Keep Tightly Closed and Protect from Light.

IMPORTANT: Do not use if seal has been punctured or is not visible.

Cap is Not Child Resistant. Keep Away from Children.

See Crimp for Expiration Date and Lot No.



Manufactured by DPT Laboratories, San Antonio, TX 78215
For Aqua Pharmaceuticals, an Almirall Company, Exton, PA 19341

N 106776 3 16110 03412 6



120g

NDC 16110-034-12

0.025%

cordran

Flurandrenolide Cream, USP

FOR EXTERNAL USE ONLY - NOT FOR USE IN THE EYES

Rx only

**PRINCIPAL DISPLAY PANEL - NDC: 16110-034-12 - 0.025% Cream 120 g
Carton Label**



PRINCIPAL DISPLAY PANEL - NDC: 16110-026-60 - 0.05% Ointment 60 g Tube Label

60 g

NDC 16110-026-60

Cordran®

Flurandrenolide Ointment, USP

0.05%

Rx only

FOR EXTERNAL USE ONLY - NOT FOR USE IN THE EYES

USUAL DOSAGE: Apply to affected area 2 or 3 times a day. See package insert.

Each gram contains: flurandrenolide 0.5 mg (0.05%), white wax, cetyl alcohol, sorbitan sesquioleate and white petrolatum.

Store at 20° to 25°C (68° to 77°F) with excursions permitted to 15° to 30°C (59° to 86°F)
[See USP controlled room temperature].

Keep Tightly Closed and Protect from Light.

IMPORTANT: Do not use if seal has been punctured or is not visible.

Cap is Not Child Resistant. Keep Away from Children.

See Crimp for Expiration Date and Lot No.

Manufactured by DPT Laboratories, San Antonio, TX 78215

For Aqua Pharmaceuticals, an Almirall Company, Eddon, PA 19341

N
3 16110 02660 2
105674

AQUA
PHARMACEUTICALS
an Almirall company



**PRINCIPAL DISPLAY PANEL - NDC: 16110-026-60 - 0.05% Ointment 60 g
Carton Label**



CORDRAN

flurandrenolide cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16110-035
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FLURANDRENOLIDE (UNII: 8EUL29XUQT) (FLURANDRENOLIDE - UNII:8EUL29XUQT)	FLURANDRENOLIDE	0.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CETYL ALCOHOL (UNII: 936JST6JCN)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
MINERAL OIL (UNII: T5L8T28FGP)	

POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

SODIUM CITRATE (UNII: 1Q73Q2JULR)

STEARIC ACID (UNII: 4ELV7Z65AP)

WATER (UNII: 059QF0KO0R)

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16110-035-60	1 in 1 CARTON	10/18/1965	
1		60 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:16110-035-12	1 in 1 CARTON	10/18/1965	
2		120 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA012806	10/18/1965	

CORDRAN

flurandrenolide cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16110-034
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FLURANDRENOLIDE (UNII: 8EUL29XUQT) (FLURANDRENOLIDE - UNII:8EUL29XUQT)	FLURANDRENOLIDE	0.25 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CETYL ALCOHOL (UNII: 936JST6JCN)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
WATER (UNII: 059QF0KOOR)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16110-034-12	1 in 1 CARTON	10/12/2018	07/30/2021
1		120 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA012806	10/12/2018	07/30/2021

CORDRAN

flurandrenolide ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16110-026
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FLURANDRENOLIDE (UNII: 8EUL29XUQT) (FLURANDRENOLIDE - UNII:8EUL29XUQT)	FLURANDRENOLIDE	0.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WHITE WAX (UNII: 7G1J5DA97F)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SORBITAN SESQUIOLEATE (UNII: 0W8RRI5W5A)	
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16110-026-60	1 in 1 CARTON	10/18/1965	01/30/2021
1		60 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA012806	10/18/1965	01/30/2021

Labeler - Almirall, LLC (605425912)

Establishment

Name	Address	ID/FEI	Business Operations
DPT Laboratories, Ltd.		832224526	ANALYSIS(16110-035, 16110-034, 16110-026) , LABEL(16110-035, 16110-034, 16110-026) , MANUFACTURE(16110-035, 16110-034, 16110-026) , PACK(16110-035, 16110-034, 16110-026)

Revised: 1/2023

Almirall, LLC