TRIAMCINOLONE ACETONIDE- triamcinolone acetonide spray, metered Apotex Corp.

Drug Facts

Active ingredient (in each spray)

Triamcinolone acetonide 55 mcg (glucocorticoid)

Purpose

Allergy symptom reliever

Uses

Temporarily relieves these symptoms of hey fever or other respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings

For external use only

Do not use

- In children under 2 years of age
- If you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if

- you have, or come in contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as persist fever
- you have any change in vision
- you have severe or frequent nosebleeds

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Read insert (inside package) on how to:

- get a new bottle ready (primed) before first use
- prime bottle again if not used for more than 2 weeks
- use the spray
- clean the spray nozzle

| ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER | | | |
|--|---|--|--|
| adults and children 12 years | once daily, spray 2 times into each nostril while sniffing gently | | |
| of age and older | once your allergy symptoms improve, reduce to 1 spray in each nostril | | |
| CHILDREN 2 TO UNDER 12 YEARS OF AGE | | | |
| • the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray longer than two months a year. | | | |
| | an adult should supervise use | | |
| children 6 to | once daily, spray 1 time into each nostril while sniffing gently | | |
| under 12 years of age | if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. | | |
| children 2 to | an adult should supervise use | | |
| under 6 years of age | once daily, spray 1 time into each nostril while sniffing gently | | |
| children under 2 years of age | ∘ do not use | | |

- do not use more than directed
- if you forget a dose, do **not** double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after one week, stop using and talk to a doctor
- do not use for the common cold

shake well before each use

Other information

- TAMPER EVIDENT: DO NOT USE IF SEALS ON CARTON ARE BROKEN OR MISSING
- Store between 20°C to 25°C (68°F to 77°F)

Inactive ingredients

benzalkonium chloride solution (50% W/V) dextrose monohydrate, edetate disodium dihydrate, hydrochloric acid, microcrystalline cellulose, polysorbate 80 purified water & sodium hydroxide

Questions or comments?

Call toll-free 1-800-706-5575

Principal Display Panel

Triamcinolone Acetonide Nasal Spray, USP 55 mcg per spray allergy symptom reliever(glucocorticoid) 120 MD per spray

NDC 60505-6275-7





TRIAMCINOLONE ACETONIDE

triamcinolone acetonide spray, metered

| Product Information | | | | |
|-------------------------|----------------|--------------------|----------------|--|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:60505-6275 | |
| Route of Administration | NASAL | | | |

| Active Ingredient/Active Moiety | | | |
|---|----------------------------|----------|--|
| Ingredient Name | Basis of Strength | Strength | |
| TRIAMCINOLONE ACETONIDE (UNII: F446C597KA) (Triamcinolone Acetonide - UNII: F446C597KA) | TRIAMCINOLONE ACETONIDE | 55 ug | |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) | |
| DEXTROSE MONOHYDRATE (UNII: LX22YL083G) | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |
| HYDROCHLORIC ACID (UNII: QTT17582CB) | |
| SODIUM HYDROXIDE (UNII: 55X04QC32I) | |
| WATER (UNII: 059QF0KO0R) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYSORBATE 80 (UNII: 60ZP39ZG8H) | |
| CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311) | |

| Packaging | | | | |
|-----------|----------------------|--|-------------------------|-----------------------|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:60505- 6275-7 | 1 in 1 CARTON | 01/26/2024 | |
| 1 | | 120 in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.) | | |

| Marketing Information | | | |
|-----------------------|---|-------------------------|-----------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA214615 | 01/26/2024 | |
| | | | |

Labeler - Apotex Corp. (845263701)

Revised: 4/2024 Apotex Corp.