

**COOL N HEAT LIDOCAINE DRY- lidocaine and menthol, unspecified form spray**  
**TOPCO ASSOCIATES LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Cool N' Heat Lidocaine Dry Spray**

**Drug Facts**

<b>Active ingredients</b>	<b>Purposes</b>
Lidocaine 4%	Topical anesthetic
Menthol 1%	Topical analgesic

**Use**

temporarily relieves minor pain

**Warnings**

**For external use only**

**Do not use**

- on large areas of the body or on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor

**When using this product**

- use only as directed. Read and follow all directions and warnings on this label
- do not allow contact with the eyes and mucous membranes
- rare cases of serious burns have been reported with products of this type
- do not apply to wounds or damaged, broken or irritated skin
- avoid spraying on face
- avoid inhalation of spray
- do not bandage or apply local heat (such as heating pads) or a medicated patch to area of use
- do not use at the same time as other topical analgesics
- a transient burning sensation may occur upon application but generally disappears in several days
- avoid applying into skin folds

**Stop use and ask a doctor if**

- condition worsens
- redness is present
- irritation develops

- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

### **Extremely Flammable**

- do not use near heat or flame or while smoking
- avoid long term storage above 104°F (40°C)
- do not puncture or incinerate. Contents under pressure
- do not store at temperatures above 120°F (49°C)

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

**adults and children over 12 years :**

- spray affected area every 6 to 8 hours, not to exceed 3 applications in a 24 hour period
- product will dry quickly on its own, and does not need to be rubbed in

**children 12 years or younger:** ask a doctor

### **Inactive ingredients**

caprylic/capric triglyceride, isobutane, propylene glycol, SD alcohol 40B (30 %)

### **Questions or comments?**

**1-888-577-8033**

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007

### **PRINCIPAL DISPLAY PANEL - 113 g Can Label**

NDC 36800-908-35

TopCare  
health®

COMPARE TO ICY HOT® LIDOCAINE PLUS  
MENTHOL DRY SPRAY ACTIVE INGREDIENTS\*

MAXIMUM STRENGTH

Cool n' Heat  
Dry Spray  
Pain Relief  
with Lidocaine + Menthol

LIDOCAINE 4% / TOPICAL ANESTHETIC  
MENTHOL 1% / TOPICAL ANALGESIC

•OUR PHARMACISTS•  
RECOMMEND

CONTINUOUS SPRAY

- Temporarily Numbs Pain Away
- Dries Instantly
- Non-Greasy

✓QUALITY  
GUARANTEED

NET WT 4 OZ (113 g)

Scan here for more information or call 1-888-423-0139

NDC 36800-908-35



COMPARE TO ICY HOT® LIDOCAINE PLUS MENTHOL DRY SPRAY ACTIVE INGREDIENTS\*

MAXIMUM STRENGTH

# Cool n' Heat Dry Spray

Pain Relief

with Lidocaine + Menthol

LIDOCAINE 4% / TOPICAL ANESTHETIC  
MENTHOL 1% / TOPICAL ANALGESIC

CONTINUOUS SPRAY

- Temporarily Numbs Pain Away
- Dries Instantly
- Non-Greasy

NET WT 4 OZ (113 g)



✓QUALITY  
GUARANTEED



Scan here for more information or call 1-888-423-0139

### Drug Facts

#### Active ingredients

Lidocaine 4%.....Topical anesthetic  
Menthol 1%.....Topical analgesic

#### Purposes

**Use** temporarily relieves minor pain

#### Warnings

For external use only

Do not use ■ on large areas of the body or on cut, irritated or swollen skin  
■ on puncture wounds ■ for more than one week without consulting a doctor

When using this product

■ use only as directed. Read and follow all directions and warnings on this label  
■ do not allow contact with the eyes and mucous membranes  
■ rare cases of serious burns have been reported with products of this type  
■ do not apply to wounds or damaged, broken or irritated skin  
■ avoid spraying on face ■ avoid inhalation of spray  
■ do not bandage or apply local heat (such as heating pads) or a medicated patch to area of use  
■ do not use at the same time as other topical analgesics  
■ a transient burning sensation may occur upon application but generally disappears in several days  
■ avoid applying into skin folds

Stop use and ask a doctor if ■ condition worsens ■ redness is present ■ irritation develops  
■ symptoms persist for more than 7 days or clear up and occur again within a few days  
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

Extremely Flammable ■ do not use near heat or flame or while smoking  
■ avoid long term storage above 104°F (40°C) ■ do not puncture or incinerate. Contents under pressure  
■ do not store at temperatures above 120°F (49°C)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

adults and children over 12 years :

■ spray affected area every 6 to 8 hours, not to exceed 3 applications  
in a 24 hour period  
■ product will dry quickly on its own, and does not need to be rubbed in  
children 12 years or younger: ask a doctor

#### Inactive ingredients

caprylyl/capric triglyceride, isobutane, propylene glycol, SD alcohol 40B (30%)

Questions or comments? 1-888-577-8033

DISTRIBUTED BY TOPOCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007  
©TOPOCO STVA1120 QUESTIONS? 1-888-423-0139 topcare@topoco.com www.topcarebrand.com

\*This product is not manufactured or distributed by Chatterm, Inc., owner of the registered trademark Icy Hot®.



DOT 2Q M5655

COOL N HEAT LIDOCAINE DRY

lidocaine and menthol, unspecified form spray

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-850
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	40 mg in 1 g
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A)	MENTHOL, UNSPECIFIED FORM	10 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>ISOBUTANE</b> (UNII: BXR49TP611)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-850-04	113 g in 1 CAN; Type 0: Not a Combination Product	05/01/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part348	05/01/2021	

**Labeler** - TOPCO ASSOCIATES LLC (006935977)

Revised: 3/2021

TOPCO ASSOCIATES LLC