

**ACETAMINOPHEN- acetaminophen tablet, film coated, extended release  
RETAIL BUSINESS SERVICES,LLC.**

-----  
**ARTHRITIS PAIN RELIEF**

**Acetaminophen Extended-release Tablets USP, 650 mg**

**Pain Reliever/Fever Reducer**

**For the temporary relief of minor arthritis pain**

**ArthritisPain**

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**Active ingredient (in each caplet)**

Acetaminophen USP, 650 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
  
- temporarily reduces fever

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

### **Ask a doctor before use if you have**

liver disease

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding**

ask a health professional before use.

### **Keep out of reach of children**

Keep out of reach of children

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see overdose warning)**

adults

- take 2 caplets every 8 hours with water

- swallow whole; do not crush, chew, split or dissolve
  - do not take more than 6 caplets in 24 hours
  - do not use for more than 10 days unless directed by a doctor
- under 18 years of age
- ask a doctor

### **Other information**

- store between 20-25°C (68-77°F)
- **do not use if foil inner seal is broken or missing**

### **Inactive ingredients**

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

### **Questions or comments?**

Contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.

### **PDP**

code # :		view
size :	1-13/4 X 1-13/4 X 3-3/8	date
ref # :	FP280604B	
material :	.016 SBS	

COATING  
FREE AREA

R-00  
551000015



**Drug Facts (continued)**

Acetaminophen, ask a doctor or pharmacist  
if you are allergic to acetaminophen or any of the inactive  
ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the  
blood thinning drug warfarin.

Always use with caution if:  
 • you push, pull, strain or back more than 10 days  
 • in lower back strain or back more than 3 days  
 • in new symptoms occur  
 • in redness or swelling is present.  
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.  
Stop use if results of studies.

**Overdose warning:** In case of overdose, get medical help or  
contact a Poison Control Center right away. (1-800-222-1222)  
Quick medical attention is critical for adults as well as for children  
even if you do not notice any signs or symptoms.

**Directions**

If the next dose more than directed (see overdose warning)

adults	take 2 capsules every 8 hours with water do not exceed 6 capsules in 24 hours do not use for more than 10 days unless directed by a doctor
under 18 years of age	ask a doctor

**Other Information**

Store between 20°-25°C (68°-77°F)  
Do not use if full timer used to break or subling

**Drug Facts (continued)**

**Active Ingredients**  
 hydroxy ethyl cellulose, hydroxyzine, magnesium  
 stearate, microcrystalline cellulose, polyethylene glycol,  
 polydioxane, pregelatinized starch, sodium starch glycolate,  
 stearic acid.

**Quantity or contents**  
 50 capsules in a white plastic bottle with a tamper-resistant cap.  
 Contains No Aspirin

\*This product is not manufactured or distributed by  
McNeil Consumer Healthcare, distributor of Tylenol®  
& HR Arthritis Pain.

DISTRIBUTED BY: FOODHOLD U.S.A., LLC

LANDOVER, MD 21766

MADE IN INDIA

© 2012 Food Hold, LLC

Quantity guaranteed at your nearest bank.

This product has been tested by an  
independent laboratory to ensure  
its quality and effectiveness.



For the Temporary Relief  
of Minor Arthritis Pain

Arthritis Pain

50 CAPSULES 650 mg each

Compare to the active ingredient of Tylenol®  
& HR Arthritis Pain—  
See New Warning

NO. 2249-118-05



3 4 1520 3 1695 1

Lot  
Exp.

COATING  
FREE AREA

CAREONE®  
ARTHRITIS  
PAIN RELIEF

Acetaminophen Extended-Release  
Tablets USP, 650 mg  
Pain Reliever/Fever Reducer

Arthritis Pain

CAREONE®  
ARTHRITIS PAIN RELIEF

Acetaminophen Extended-Release  
Tablets USP, 650 mg  
Pain Reliever/Fever Reducer

For the Temporary Relief  
of Minor Arthritis Pain

Arthritis Pain

50 CAPSULES 650 mg each

Compare to the active ingredient of Tylenol®  
& HR Arthritis Pain—  
See New Warning

Important: Read all product information before using.  
Keep the carton for important information.

Do not use if full timer used to break or subling

**Drug Facts**

**Active ingredient (in each capsule):** Purpose  
Acetaminophen USP, 650 mg. **7-in release/fever reducer**

**Uses**

- to temporarily relieve minor aches and pains due to:  
 • minor pain of arthritis  
 • muscular aches  
 • headache  
 • sinusitis and nasal congestion  
 • the common cold  
 • toothache  
 • to temporarily reduce fever

**Warnings**

Use as directed. This product contains acetaminophen. Seven  
overdoses may occur if you take  
 • more than 6 capsules in 24 hours, which is the maximum daily  
 amount.

Do not take other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product.  
 Always drink acetaminophen may cause severe liver damage.  
 Symptoms may include:  
 • loss of appetite  
 • nausea  
 • vomiting  
 • loss of energy  
 • loss of weight  
 • loss of interest in food  
 • loss of interest in activities  
 • loss of interest in sex  
 • loss of interest in work  
 • loss of interest in school  
 • loss of interest in sports  
 • loss of interest in hobbies  
 • loss of interest in social activities  
 • loss of interest in family activities  
 • loss of interest in friends  
 • loss of interest in pets  
 • loss of interest in travel  
 • loss of interest in shopping  
 • loss of interest in reading  
 • loss of interest in television  
 • loss of interest in movies  
 • loss of interest in music  
 • loss of interest in games  
 • loss of interest in puzzles  
 • loss of interest in board games  
 • loss of interest in card games  
 • loss of interest in dice games  
 • loss of interest in board games  
 • loss of interest in card games  
 • loss of interest in dice games





<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	

### Product Characteristics

<b>Color</b>	white (White to off white colored)	<b>Score</b>	no score
<b>Shape</b>	OVAL (Capsule shaped, biconvex intact film coated tablets)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	G;650
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-185-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/19/2020	
2	NDC:72476-185-21	225 in 1 BOTTLE; Type 0: Not a Combination Product	08/19/2020	
3	NDC:72476-185-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/19/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	08/19/2020	

**Labeler** - RETAIL BUSINESS SERVICES,LLC. (967989935)

Revised: 12/2023

RETAIL BUSINESS SERVICES,LLC.