

**DIABETIC TUSSIN DM MAXIMUM STRENGTH- dextromethorphan hydrobromide and guaifenesin liquid**  
**MEDTECH PRODUCTS INC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**DiabeticTussin DM Max Strength 61787-515**

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan hydrobromide 20 mg

Guaifenesin 400 mg

**Purposes**

Cough Suppressant

Expectorant

**Uses**

- temporarily relieves cough caused by the common cold or inhaled irritants.
- helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus.

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a cough that occurs with too much phlegm (mucus)
- a chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor if**

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- take every 4 hours
- do not exceed 6 doses in 24 hours

Adults	10 mL (2 teaspoons)
Children under 12 years	ask a doctor

**Other information**

- store at room temperature 20-25°C (68-77°F)
- keep tightly closed

**Inactive ingredients**

Acesulfame K, artificial raspberry flavors, hypromellose, menthol, methylparaben, polyethylene glycol, potassium sorbate, purified water and sucralose. Citric acid may be used to adjust pH.

**Questions or comments?**

Call **1-800-579-8327**, serious side effects associated with use of this product may be reported to this number.

**Package Principal Display Panel**

**SUGAR & ALCOHOL FREE!**

**Specifically Formulated for Diabetics**

**Diabetic Tussin®**

**MAXIMUM STRENGTH**

**COUGH & CHEST CONGESTION**

**DM**

Dextromethorphan HBr (Cough Suppressant)

Guaifenesin (Expectorant)

**Relieves:**

- Mucus
- Cough
- Sore throat
- Chest congestion

Improved Berry Flavor

4 FL OZ (118 mL)

**Diabetic Tussin. DM**  
**MAXIMUM STRENGTH**  
**Cough & Chest Congestion DM**

**SUGAR FREE!**  
 FOR PEOPLE WITH DIABETES

**THE Diabetic Tussin. DIFFERENCE**

**Specifically Formulated FOR PEOPLE WITH DIABETES**

**MAXIMUM STRENGTH**

**Cough & Chest Congestion DM**

**EVERYTHING YOU WANT AND NOTHING YOU DON'T**

**RELIEVES**

- cough
- chest congestion
- mucus
- sore throat

**MADE WITHOUT:**

- SUGAR - ALCOHOL
- SODIUM - SORBITOL
- FRUCTOSE - GLUTEN
- DYES OR ASPARTAME

**#1 PHARMACIST RECOMMENDATION**

**BERRY**  
 4 FL OZ (118 mL)

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan hydrobromide 20 mg Cough Suppressant  
 Guaifenesin 600 mg Expectorant

**Purposes**

**Uses**

- temporarily relieves cough caused by the common cold or inhaled irritants
- helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts as these could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**

- take every 4 hours
- do not exceed 6 doses in 24 hours

**Adults** 10 mL (2 teaspoons)  
**Children under 12 years** ask a doctor

**Other information**

- store at room temperature 20° to 25°C (68° to 77° F)
- keep tightly closed

**Drug Facts (continued)**

**Inactive ingredients**

acesulfame potassium, artificial raspberry flavor, hydroxytoluene, menthol, methylparaben, polyethylene glycol, potassium sorbate, purified water, and sucralose. Citric acid may be used to adjust pH.

**Questions or comments?**  
 Call 1-800-579-8327. Serious side effects associated with use of this product may be reported to this number.

**PERSONS WITH DIABETES - CONSULT YOUR DOCTOR OR PHARMACIST BEFORE USING ANY MEDICATION OR DIETARY SUPPLEMENT.**

**THIS PRODUCT DOES NOT CONTAIN ANY MEDICATION TO IMPROVE OR RELIEVE DIABETES.**

**SAVE CARTON FOR COMPLETE DRUG FACTS.**

**TAMPER EVIDENT: FOR YOUR PROTECTION THIS BOTTLE HAS A SAFETY SEAL AROUND THE CAP OF THE BOTTLE NECK AND UNDER THE CAP.**

Dist. by Medach Products Inc.  
 Tonawanda, NY 10581  
 A Praxair Consumer Healthcare company  
 ©2022 Trade dress is owned by Medach Products Inc.  
 All rights reserved.  
 Made in USA.  
 DTUS02101

\*U.S. News & World Report & Pharmacy Times, 2021-2022



**DIABETIC TUSSIN DM MAXIMUM STRENGTH**

dextromethorphan hydrobromide and guaifenesin liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:61787-515
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 10 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>RASPBERRY</b> (UNII: 4N14V5R27W)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	RASPBERRY (artificial raspberry flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:61787-515-04	1 in 1 BOX	02/01/2020	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:61787-515-08	1 in 1 BOX	02/01/2020	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
---------------------------	---	-----------------------------	---------------------------

OTC monograph final	part341	02/01/2020	
---------------------	---------	------------	--

**Labeler** - MEDTECH PRODUCTS INC (114707784)

### **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Akorn Operating Company LLC (dba Akorn)		117696873	manufacture(61787-515)

Revised: 3/2022

MEDTECH PRODUCTS INC