

CURIST PAIN RELIEF- acetaminophen tablet
Little Pharma, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Curist Pain Relief (Acetaminophen 500 mg) Tablets

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever / Fever reducer

Uses

- temporarily relieves minor aches and pains
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**
- **adults and children 12 years and over:** take 2 tablets every 6 hours while symptoms last. Do not take more than 6 tablets in 24 hours.
- **children under 12 years of age:** do not use

Other information

store between 20-25°C (68-77°F)

Inactive ingredients

polyvinylpyrrolidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

curisrelief.com

DO NOT USE IF INNER FOIL SEAL UNDER BOTTLE CAP IS BROKEN OR MISSING.

CONTAINS NO ASPIRIN

Distributed by: Little Pharma, Inc., New York, NY 10023

1-844-243-1241

curist

Pain Relief

Acetaminophen

Tablets, 500 mg

Pain Reliever / Fever Reducer

Extra Strength

90 Tablets

curist

Pain Relief

Acetaminophen
Tablets, 500 mg
Pain Reliever / Fever Reducer
Extra Strength

90 Tablets

curistrelief.com

DO NOT USE IF INNER FOIL SEAL UNDER BOTTLE CAP IS BROKEN OR MISSING.

Drug Facts

Active Ingredient (in each tablet)
Acetaminophen 500 mg, Pain reliever/Fever reducer

Purpose
• temporarily reduces fever

Uses
• temporarily relieves minor aches and pains

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 4000 mg of acetaminophen in 24 hours
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin redness • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away

CONTAINS NO ASPHIN (CONTINUED ON BACK OF LABEL)
Distributed by: Little Pharma, Inc. New York, NY 10023
1-844-243-1241

3 72559 00805 0

PEEL HERE

Glue Area

STOP PEELING

Drug Facts (continued)

Do not use - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if - pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present - new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not take more than directed (see overdose warning)
• adults and children 12 years and over: take 2 tablets every 6 hours while symptoms last; do not take more than 6 tablets in 24 hours, unless directed by a doctor
• children under 12 years of age: do not use

Other information
store between 20-25°C (68-77°F)

Inactive Ingredients
polyvinylpyrrolidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

CURIST PAIN RELIEF

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72559-008
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
--------------	-------	--------------	----------

Shape	ROUND	Size	12mm
Flavor		Imprint Code	AP;013
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72559-008-05	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/07/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/07/2020	

Labeler - Little Pharma, Inc. (074328189)

Revised: 4/2020

Little Pharma, Inc.