

**ALL DAY PAIN RELIEF- naproxen sodium capsule, liquid filled
Family Dollar (FAMILY WELLNESS)**

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

***nonsteroidal anti-inflammatory drug**

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed

Heart attack and stroke: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222)

Directions

- **do not take more than directed**

- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work
- adults and children 12 years and older:
 - take 1 capsule every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 capsules within the first hour
 - do not exceed 2 capsules in any 8- to 12-hour period
 - do not exceed 3 capsules in a 24-hour period
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- read all directions and warnings before use.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol 600, povidone K30, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

**COMPARE TO THE ACTIVE INGREDIENT IN ALEVE®†
STRENGTH TO LAST 12 HOURS**

All Day Pain Relief

Naproxen Sodium Capsules 220 mg

Pain Reliever / Fever Reducer (**NSAID**)

CAPSULES

(**LIQUID-FILLED CAPSULES)

†This product is not manufactured or distributed by Bayer HealthCare, LLC distributor of Aleve®

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC

10611 MONROE., MATTHEWS, NC 28105

Product Label

Lot No.:
Exp. Date:

Rev # 02/17

3225 1 38050 1

PLD-A4337E F0004599 SKU099788

Drug Facts (continued)

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ drink a full glass of water with each dose
■ if taken with food, this product may take longer to work
■ adults and children 12 years and older:
■ take 1 capsule every 8 to 12 hours while symptoms last
■ for the first dose you may take 2 capsules within the first hour
■ do not exceed 2 capsules in any 8- to 12-hour period
■ do not exceed 3 capsules in a 24-hour period
■ children under 12 years: ask a doctor

Other information
■ each capsule contains: sodium 20 mg
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
■ read all directions and warnings before use. Keep carton.

inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol 600, polyvidone K30, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Drug Facts (continued)

Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if
■ the stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking any other drug

When using this product, take with food or milk if stomach upset occurs.

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ slurred speech
■ leg swelling ■ trouble breathing
■ weakness in one part or side of body
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear
■ you have difficulty swallowing
■ it feels like the capsule is stuck in your throat

This product is not manufactured or distributed by Bayer HealthCare, LLC, distributor of Aleve®.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
10511 MONROE RD., MATTHEWS, NC 28105

NOT 100% SATISFIED?
Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg (NSAID)*, nonsteroidal anti-inflammatory drug)

Purposes
Pain reliever/fever reducer

Uses
■ temporarily relieves minor aches and pains due to:
■ minor pain of arthritis ■ muscular aches
■ backache ■ menstrual cramps
■ headache ■ toothache
■ temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed
■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

COMPARE TO THE ACTIVE INGREDIENT IN ALEVE®†

NDC 55319-174-20

FAMILY wellness

STRENGTH TO LAST 12 HOURS

All Day Pain Relief

Naproxen Sodium Capsules 220 mg
Pain Reliever/Fever Reducer (NSAID) ACTUAL SIZE

20 CAPSULES**
(**LIQUID-FILLED CAPSULES)

100% SATISFACTION OR YOUR MONEY BACK GUARANTEED

OTC network

FAMILY WELLNESS All Day Pain Relief

ALL DAY PAIN RELIEF			
naproxen sodium capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-174
Route of Administration	ORAL		
Active Ingredient/Active Moiety			

Ingredient Name		Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM	220 mg	
Inactive Ingredients				
Ingredient Name		Strength		
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)				
GELATIN (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
LACTIC ACID (UNII: 33X04XA5AT)				
MANNITOL (UNII: 3OWL53L36A)				
POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9)				
POVIDONE K30 (UNII: U725QWY32X)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SORBITAN (UNII: 6O92ICV9RU)				
SORBITOL (UNII: 506T60A25R)				
Product Characteristics				
Color	BLUE	Score	no score	
Shape	CAPSULE	Size	21mm	
Flavor		Imprint Code	NPX	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-174-20	1 in 1 BOX	12/31/2017	12/30/2022
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA021920	12/31/2017	12/30/2022	

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)

Revised: 12/2020

Family Dollar (FAMILY WELLNESS)