

**ANTACID- aluminum hydroxide, magnesium hydroxide, simethicone liquid**  
**Rij Pharmaceutical Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 5 mL) (One teaspoonful)**

Aluminum Hydroxide (Equi. to Dried Gel, USP) 225 mg

Magnesium Hydroxide 200 mg

Simethicone 25 mg

**Purpose**

Antacid

Antigas

**Uses**

relieves

- heartburn
- acid indigestion
- sour stomach
- associated symptoms of gas and upset stomach
- pressure and bloating commonly referred to as gas

**Warnings**

**Do not take** more than 12 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks. Except under the advice and supervision of a physician.

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if**

symptoms last for more than 2 weeks

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

### **Directions**

- **shake well before each use**
- Adults and children 12 years and older, take 2 to 4 teaspoonsful between meals, at bedtime or as directed by a physician
- do not take more than 12 teaspoonsful in 24 or use the maximum dosage for more than 2 weeks
- Children under 12 years: consult a physician

### **Other information**

- TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING
- Store at room temperature 15° - 30°C (59° - 86°F)
- keep tightly closed
- protect from freezing
- each teaspoon (5 mL) contains: magnesium 83 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

### **Inactive ingredients**

flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, water and xanthan gum

### **Principal Display Panel**

NDC 53807-151-12

Compare to the Active  
Ingredient in Maalox® Plus†



**PLUS**

# Antacid & Antigas

Magnesia and Alumina

Oral Suspension  
with Simethicone

Non-Constipating  
Pleasant Tasting



**Lemon Creme Flavored**

**12 FL. OZ (355 mL)**

# Antacid & Antigas Plus

## Drug Facts

### Active Ingredients

(In each 5 ml) (One teaspoonful)

### Purpose

Aluminum Hydroxide (Equi. to Dried Gel, USP) 225 mg .....	Antacid
Magnesium Hydroxide 200 mg .....	Antacid
Simethicone 25 mg .....	Antigas

**Uses** relieves ■ heartburn ■ acid indigestion ■ sour stomach  
■ associated symptoms of gas and upset stomach ■ pressure and bloating commonly referred to as gas

## Warnings

**Do not take** more than 12 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks. Except under the advise and supervision of a physician.

### Ask a doctor before use if you have

- kidney disease ■ a magnesium-restricted diet

### Ask a doctor or pharmacist before use if you are

taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ symptoms last more than 2 weeks

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

## Directions ■ shake well before each use

- Adults and children 12 years and older, take 2 to 4 teaspoons between meals, at bedtime or as directed by a physician
- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- Children under 12 years: consult a physician

## Other Information

- **TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.**
- store at room temperature 15°-30°C (59°-86°F) ■ keep tightly closed
- protect from freezing
- each teaspoon (5 mL) contains: magnesium 83 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

**Inactive ingredients:** flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, water and xanthan gum.

**Questions? 845-692-5799**

†This product is not manufactured or distributed by the owner of the registered trademark Maalox® Plus All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from retailer where purchased.

Manufactured By:  
RIJ PHARMACEUTICAL CORP.  
40 Commercial Avenue  
Middletown, NY 10941  
Distributed By:  
Kinray Inc.,  
Whitestone, NY 11357



## ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone liquid

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53807-151
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	225 mg in 5 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	25 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
BUTYL PARABEN (UNII: 3QP11U3FV8)	
SACCHARIN (UNII: FST467XS7D)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	
SORBITOL (UNII: 506T60A25R)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
LEMON OIL (UNII: I9GRO824LL)	

## Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53807-151-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part332	03/16/1999	

---

## Establishment

Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-151)

Revised: 4/2018

Rij Pharmaceutical Corporation