MINOXIDIL EXTRA STRENGTH FOR MEN- minoxidil solution Chain Drug Consortium, LLC

Minoxidil Topical Solution for Men

Drug Facts

Active ingredient

Minoxidil, USP 5% w/v

Purpose

Hair regrowth treatment for men

Use

to regrow hair on the top of the scalp (vertex only, see pictures on side of carton)

Warnings

For external use only. For use by men only.

Flammable: Keep away from fire or flame

Do not use if

- you are a woman
- your amount of hair loss is different than that shown on the side of this carton or your hair loss is on the front of the scalp. Minoxidil topical solution 5% is not intended for frontal baldness or receding hairline.
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have

heart disease

When using this product

- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 2 months with twice a day usage.

For some men, you may need to use this product for at least 4 months before you see results.

• the amount of hair regrowth is different for each person. This product will not work for all men.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply one mL with dropper 2 times a day directly onto the scalp in the hair loss area
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- hair regrowth has not been shown to last longer than 48 weeks in large clinical trials with continuous treatment with minoxidil topical solution 5% for men
- In clinical studies with mostly white men aged 18 to 49 years with moderate degrees of hair loss, minoxidil topical solution 5% for men provided more hair regrowth than minoxidil topical solution 2%
- store at controlled room temperature 20° to 25°C (68° to 77°F)

Inactive ingredients

alcohol (30% v/v), propylene glycol (50% v/v), purified water

Questions?

1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Principal Display Panel

Premier Value®

COMPARE TO THE ACTIVE INGREDIENT IN MEN'S ROGAINE® EXTRA STRENGTH*

Extra Strength For Men Minoxidil TOPICAL SOLUTION, USP 5% HAIR REGROWTH TREATMENT

Reactivates Hair Follicles to Stimulate Regrowth

Clinically Proven to Help Regrow Hair

Unscented

NOT FOR USE BY WOMEN

FOR BEST RESULTS
APPLY DIRECTLY TO SCALP
TWICE DAILY

Two Month Supply
Two 60mL (2 fl oz) Bottles

INDEPENDENTLY TESTED SATISFACTION GUARANTEED





COMPARE TO THE ACTIVE INGREDIENT IN MEN'S ROGAINE® EXTRA STRENGTH*

Extra Strength For Men

Minoxidil

TOPICAL SOLUTION, USP 5% HAIR REGROWTH TREATMENT

Reactivates Hair Follicles to Stimulate Regrowth

Clinically Proven to Help Regrow Hair

Unscented


NOT FOR USE BY WOMEN


FOR BEST RESULTS

APPLY DIRECTLY TO SCALP
TWICE DAILY

Two Month Supply Two 60mL (2 fl oz) Bottles







Drug Facts

Purpose

Use to regrow hair on the top of the scalp (vertex only, see pictures on side of carton)

For external use only. For use by men only. Flammable: Keep away from fire or flame

Do not use if

- your amount of hair loss is different than that shown on the side of this carton or your hair loss is on the front of the scalp. Minoxidil topical solution 5% is not intended for frontal baldness or receding hairline.
- you have no family history of hair loss
 your hair loss is sudden and/or patchy
- you do not know the reason for your hair loss
 you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
 you use other medicines on the scalp

Ask a doctor before use if you have heart disease

When using this product

- do not apply on other parts of the body
- · avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 2 months with twice a day usage. For some men, you may need to use this product for at least 4 months before you see results. • the amount of hair regrowth is different for each person. This product will not work for all men.
- Stop use and ask a doctor if
- chest pain, rapid heartbeat, faintness, or dizziness occurs scalp irritation or redness occurs
- sudden, unexplained weight gain occurs your hands or feet swell
- unwanted facial hair growth occurs
 you do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply one mL with dropper 2 times a day directly onto the scalp in the hair loss area
 using more or more often will not improve results
- · continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
 keep the carton. It contains important information.
- hair regrowth has not been shown to last longer than 48 weeks in large clinical trials with continuous treatment with minoxidil topical solution 5% for men
- In clinical studies with mostly white men aged 18 to 49 years with moderate degrees of hair loss, minoxidil topical solution 5% for men provided more hair regrowth than minoxidil tonical solution 2%
- store at controlled room temperature 20° to 25°C (68° to 77°F)

Inactive ingredients alcohol (30% v/v), propylene glycol (50% v/v), purified water

Questions? 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Not For Use By Women:

May grow facial hair. May be harmful if used during pregnancy or breast-feeding.

Minoxidil Topical Solution 5% is for men who have a general thinning of hair on the top of the scalp (vertex only, as shown below). Not intended for frontal baldness or a receding hairline. Minoxidil Topical Solution 5% for men has been shown to regrow hair in men with the following degrees of thinning hair or hair loss.







If men have more hair loss or hair loss in a place different than shown above, Minoxidil Topical Solution 5% for men may not work.

This Package Contains:

- Two 60 mL (2 fl oz) Bottles of Minoxidil Topical Solution 5% (two month supply).
- One child-resistant dropper applicator
- Information leaflet with complete directions on how to use and obtain best results
- *This product is not affiliated with, manufactured by, or produced by the makers or owners of Rogaine®

Distributed by: Pharmacy Value Alliance, LLC 407 East Lancaster Avenue, Wayne, PA 19087



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

Made in Canada 00940916 R1116 VC114246

MINOXIDIL EXTRA STRENGTH FOR MEN

minoxidil solution

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68016-382

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

Inactive Ingredients

Ingredient Name Strength

ALCOHOL (UNII: 3K9958V90M)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)

Packaging

	rackaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:68016- 382-00	2 in 1 CARTON	09/11/2003			
1		60 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product				

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
ANDA	ANDA075518	09/11/2003	

Revised: 11/2022 Chain Drug Consortium, LLC