

**MEDIQUE MEDI-LAX ENTERIC COATED- bisacodyl 5 mg tablet, sugar coated**  
**Unifirst First Aid Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Medique Medi-lax Enteric Coated**

***Drug Facts***

***Active ingredient (in each tablet)***

Bisacodyl USP 5 mg

***Purpose***

Stimulant laxative

***Uses***

- for the temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

***Warnings***

**Do not use** if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness or cramps

**Stop use and ask a doctor if you**

- have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- need to use a laxative for more than 1 week

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## ***Directions***

**Adults and children 12 years and older:** 1 to 3 tablets in a single daily dose

**Children under 12 years:** do not give to children under 12 years of age

## ***Other information***

- store between 68° to 77°F (20° to 25°C)
- tamper evident sealed blister pack(s)
- do not use if a blister unit is torn, broken or shows any signs of tampering

## ***Inactive ingredients***

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C Yellow #10 aluminum lake, FD&C Yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

## ***Questions or comments***

Call 1-800-634-7680

## **Medique Medi-lax Label**

Medique®

Medi-lax

Stimulant Laxative

Bisacodyl Delayed-Release Tablets USP, 5 mg

\*Compare to the active ingredients in Ducolax®

Enteric Coated Tablets

Tamper Evident Blister Pack

25 Tablets

## Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
Bisacodyl USP 5 mg.....Stimulant laxative

**Uses** ■ for temporary relief of occasional constipation and irregularity  
■ this product generally produces bowel movement in 6 to 12 hours

### Warnings

Do not use if you cannot swallow without chewing.

Ask a doctor before use if you have ■ stomach pain, nausea or vomiting  
■ a sudden change in bowel habits that lasts more than 2 weeks

When using this product ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk ■ it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if you ■ have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

■ need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

■ take with a glass of water

Adults and children 12 years of age and older: 1 to 3 tablets in a single daily dose

Children 6 to under 12 years of age: 1 tablet in a single daily dose

Children under 6 years of age: ask a doctor

### Other information

■ store between 68° to 77°F (20° to 25°C) ■ tamper evident sealed blister pack(s)

■ do not use if a blister unit is torn, broken or shows any signs of tampering

**Inactive ingredients** acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments?** Call 1-800-634-7680

**RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION**

**Medi-lax**  
Stimulant Laxative

**Medique**

**Medi-lax**

Stimulant Laxative

**Medi-lax**  
Stimulant Laxative

Bisacodyl Delayed-Release Tablets USP, 5 mg

\* Compare to the active ingredient in Dulcolax®

25  
Tablets

Enteric Coated Tablets  
Tamper Evident Blister Pack



Reorder #85889  
Mfd. for Medique Products  
Fort Myers, FL 33967  
www.medicqueproducts.com  
\*This product is not manufactured or distributed by  
Boehringer Ingelheim Pharmaceuticals, Inc.,  
owner of the registered trademark Dulcolax®.

## MEDIQUE MEDI-LAX ENTERIC COATED

bisacodyl 5 mg tablet, sugar coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-859
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISACODYL</b> (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL 1000</b> (UNII: U076Q6Q621)	
<b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	

<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>ACACIA</b> (UNII: 5C5403N26O)
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SODIUM ALGINATE</b> (UNII: C269C4G2ZQ)
<b>AMMONIA</b> (UNII: 5138Q19F1X)
<b>POVIDONE</b> (UNII: FZ989GH94E)
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	5
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-859-05	1 in 1 BOX	07/01/2020	
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	07/01/2020	

**Labeler** - Unifirst First Aid Corporation (832947092)