

ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ ORANGE ZEST- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent Bayer HealthCare LLC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

AS Severe Cold PowerFast Fizz Orange Zest UI1614457

Drug Facts

Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)*

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Antihistamine

Nasal decongestant

Uses

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains · headache · runny nose
- nasal and sinus congestion · sneezing · sore throat
- temporarily reduces fever

Directions

Directions

● adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.

● children under 12 years: do not use

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if

changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have

- asthma ● diabetes
- thyroid disease ● glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for
- gout · diabetes · arthritis
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint ● vomit blood ● have bloody or black stools
- have stomach pain that does not get better
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness, or sleeplessness occur

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information

Other information

- each tablet contains: potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

Inactive ingredients

Inactive ingredients anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavor, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

Questions or Comments

Questions or comments? **1-800-986-0369** (Mon - Fri 9AM - 5PM EST)

Carton 24 count



Alka-Seltzer PLUS® Severe Cold Orange Zest
PowerFast Fizz™

NEW Improved Flavor

ASPIRIN (NSAID) / Pain Reliever-Fever Reducer

Chlorpheniramine Maleate / Antihistamine

Phenylephrine Bitartrate / Nasal Decongestant

- Nasal Congestion
- Runny Nose
- Sore Throat
- Headache + Body Ache
- Sinus Pressure

24 EFFERVESCENT TABLETS

ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ ORANGE ZEST

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-0020
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1W5297W6MV)	PHENYLEPHRINE BITARTRATE	7 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
MANNITOL (UNII: 3OWL53L36A)	
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
POVIDONE (UNII: FZ989GH94E)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CALCIUM SILICATE (UNII: S4255P4G5M)	
DIMETHICONE (UNII: 92RU3N3Y1O)	

Product Characteristics

Color	white (SPOTTED)	Score	no score
Shape	ROUND	Size	25mm
Flavor	ORANGE	Imprint Code	ASP;ORANGE
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0020-01	10 in 1 CARTON	07/01/2020	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:0280-0020-02	12 in 1 CARTON	03/30/2023	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/01/2020	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 8/2023

Bayer HealthCare LLC.