

# **NOREPINEPHRINE BITARTRATE - norepinephrine bitartrate injection**


## **Baxter Healthcare Corporation**

-----

### **DESCRIPTION**

Norepinephrine (sometimes referred to as *l-arterenol/Levarterenol* or *l-norepinephrine*) is a sympathomimetic amine which differs from epinephrine by the absence of a methyl group on the nitrogen atom.

Norepinephrine bitartrate is (-)- $\alpha$ -(aminomethyl)-3,4-dihydroxybenzyl alcohol tartrate (1:1) (salt) monohydrate and has the following structural formula:



Norepinephrine is supplied in sterile aqueous solution in the form of the bitartrate salt to be administered by intravenous infusion following dilution. Norepinephrine is sparingly soluble in water, very slightly soluble in alcohol and ether, and readily soluble in acids. Each mL contains the equivalent of 1 mg base of norepinephrine, sodium chloride for isotonicity. It has a pH of 3 to 4.5. The air in the ampules has been displaced by nitrogen gas.

### **CLINICAL PHARMACOLOGY**

Norepinephrine functions as a peripheral vasoconstrictor (alpha-adrenergic action) and as an inotropic stimulator of the heart and dilator of coronary arteries (beta-adrenergic action).

### **INDICATIONS AND USAGE**

For blood pressure control in certain acute hypotensive states (e.g., pheochromocytectomy, sympathectomy, poliomyelitis, spinal anesthesia, myocardial infarction, septicemia, blood transfusion, and drug reactions).

As an adjunct in the treatment of cardiac arrest and profound hypotension.

### **CONTRAINDICATIONS**

Norepinephrine Bitartrate Injection, USP should not be given to patients who are

hypotensive from blood volume deficits except as an emergency measure to maintain coronary and cerebral artery perfusion until blood volume replacement therapy can be completed. If Norepinephrine Bitartrate Injection, USP is continuously administered to maintain blood pressure in the absence of blood volume replacement, the following may occur: severe peripheral and visceral vasoconstriction, decreased renal perfusion and urine output, poor systemic blood flow despite "normal" blood pressure, tissue hypoxia, and lactate acidosis.

Norepinephrine Bitartrate Injection, USP should also not be given to patients with mesenteric or peripheral vascular thrombosis (because of the risk of increasing ischemia and extending the area of infarction) unless, in the opinion of the attending physician, the administration of Norepinephrine Bitartrate Injection, USP is necessary as a life-saving procedure.

Cyclopropane and halothane anesthetics increase cardiac autonomic irritability and therefore seem to sensitize the myocardium to the action of intravenously administered epinephrine or norepinephrine. Hence, the use of Norepinephrine Bitartrate Injection, USP during cyclopropane and halothane anesthesia is generally considered contraindicated because of the risk of producing ventricular tachycardia or fibrillation.

The same type of cardiac arrhythmias may result from the use of Norepinephrine Bitartrate Injection, USP in patients with profound hypoxia or hypercarbia.

## **WARNINGS**

Norepinephrine should be used with extreme caution in patients receiving monoamine oxidase inhibitors (MAOI) or antidepressants of the triptyline or imipramine types, because severe, prolonged hypertension may result.

## **PRECAUTIONS**

### **General**

**Avoid Hypertension:** Because of the potency of norepinephrine and because of varying response to pressor substances, the possibility always exists that dangerously high blood pressure may be produced with overdoses of this pressor agent. It is desirable, therefore, to record the blood pressure every two minutes from the time administration is started until the desired blood pressure is obtained, then every five minutes if administration is to be continued.

The rate of flow must be watched constantly, and the patient should never be left unattended while receiving norepinephrine. Headache may be a symptom of hypertension due to overdosage.

**Site of Infusion:** Whenever possible, infusions of norepinephrine should be given into a large vein, particularly an antecubital vein because, when administered into this vein, the risk of necrosis of the overlying skin from prolonged vasoconstriction is apparently very slight. Some authors have indicated that the femoral vein is also an acceptable route of administration. A catheter tie-in technique should be avoided, if possible, since the obstruction to blood flow around the tubing may cause stasis and increased local concentration of the drug. Occlusive vascular diseases (for example, atherosclerosis,

arteriosclerosis, diabetic endarteritis, Buerger's disease) are more likely to occur in the lower than in the upper extremity. Therefore, one should avoid the veins of the leg in elderly patients or in those suffering from such disorders. Gangrene has been reported in the lower extremity when infusions of norepinephrine were given in an ankle vein.

**Extravasation:** The infusion site should be checked frequently for free flow. Care should be taken to avoid extravasation of norepinephrine into the tissues, as local necrosis might ensue due to the vasoconstrictive action of the drug. Blanching along the course of the infused vein, sometimes without obvious extravasation, has been attributed to vasa vasorum constriction with increased permeability of the vein wall, permitting some leakage.

This also may progress on rare occasions to superficial slough, particularly during infusion into leg veins in elderly patients or in those suffering from obliterative vascular disease. Hence, if blanching occurs, consideration should be given to the advisability of changing the infusion site at intervals to allow the effects of local vasoconstriction to subside.

**IMPORTANT- Antidote for Extravasation Ischemia:** To prevent sloughing and necrosis in areas in which extravasation has taken place, the area should be infiltrated as soon as possible with 10 mL to 15 mL of saline solution containing from 5 mg to 10 mg of phentolamine, an adrenergic blocking agent. A syringe with a fine hypodermic needle should be used, with the solution being infiltrated liberally throughout the area, which is easily identified by its cold, hard, and pallid appearance. Sympathetic blockade with phentolamine causes immediate and conspicuous local hyperemic changes if the area is infiltrated within 12 hours. Therefore, phentolamine should be given as soon as possible after the extravasation is noted.

### **Drug Interactions:**

Cyclopropane and halothane anesthetics increase cardiac autonomic irritability and therefore seem to sensitize the myocardium to the action of intravenously administered epinephrine or norepinephrine. Hence, the use of norepinephrine during cyclopropane and halothane anesthesia is generally considered contraindicated because of the risk of producing ventricular tachycardia or fibrillation. The same type of cardiac arrhythmias may result from the use of norepinephrine in patients with profound hypoxia or hypercarbia.

Norepinephrine should be used with extreme caution in patients receiving monoamine oxidase inhibitors (MAOI) or antidepressants of the triptyline or imipramine types, because severe, prolonged hypertension may result.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility:**

Studies have not been performed.

### **Pregnancy Category C:**

Animal reproduction studies have not been conducted with norepinephrine. It is also not known whether norepinephrine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Norepinephrine should be given to a pregnant woman only if clearly needed.

### **Nursing Mothers:**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when norepinephrine is administered to a nursing woman.

### **Pediatric Use:**

Safety and effectiveness in pediatric patients has not been established.

### **Geriatric Use:**

Clinical studies of norepinephrine bitartrate injection did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Norepinephrine infusions should not be administered into the veins in the leg in elderly patients (see **PRECAUTIONS**, General).

## **ADVERSE REACTIONS**

The following reactions can occur:

*Body As A Whole:* Ischemic injury due to potent vasoconstrictor action and tissue hypoxia.

*Cardiovascular System:* Bradycardia, probably as a reflex result of a rise in blood pressure, arrhythmias.

*Nervous System:* Anxiety, transient headache.

*Respiratory System:* Respiratory difficulty.

*Skin and Appendages:* Extravasation necrosis at injection site.

Prolonged administration of any potent vasopressor may result in plasma volume depletion which should be continuously corrected by appropriate fluid and electrolyte replacement therapy. If plasma volumes are not corrected, hypotension may recur when norepinephrine is discontinued, or blood pressure may be maintained at the risk of severe peripheral and visceral vasoconstriction (e.g., decreased renal perfusion) with diminution in blood flow and tissue perfusion with subsequent tissue hypoxia and lactic acidosis and possible ischemic injury. Gangrene of extremities has been rarely reported.

Overdoses or conventional doses in hypersensitive persons (e.g., hyperthyroid patients) cause severe hypertension with violent headache, photophobia, stabbing retrosternal pain, pallor, intense sweating, and vomiting.

## **OVERDOSAGE**

Overdosage with norepinephrine may result in headache, severe hypertension, reflex

bradycardia, marked increase in peripheral resistance, and decreased cardiac output. In case of accidental overdosage, as evidenced by excessive blood pressure elevation, discontinue norepinephrine until the condition of the patient stabilizes.

## **DOSAGE AND ADMINISTRATION**

**Norepinephrine Bitartrate Injection is a concentrated, potent drug which must be diluted in dextrose containing solutions prior to infusion. An infusion of norepinephrine should be given into a large vein (see PRECAUTIONS).**

### **Restoration of Blood Pressure in Acute Hypotensive States**

Blood volume depletion should always be corrected as fully as possible before any vasopressor is administered. When, as an emergency measure, intraaortic pressures must be maintained to prevent cerebral or coronary artery ischemia, norepinephrine can be administered before and concurrently with blood volume replacement.

**Diluent:** Norepinephrine should be diluted in 5 percent dextrose injection or 5 percent dextrose and sodium chloride injections. These dextrose containing fluids are protection against significant loss of potency due to oxidation. **Administration in saline solution alone is not recommended.** Whole blood or plasma, if indicated to increase blood volume, should be administered separately (for example, by use of a Y-tube and individual containers if given simultaneously).

*Average Dosage:* Add a 4 mL ampul (4 mg) of norepinephrine to 1,000 mL of a 5 percent dextrose containing solution. Each mL of this dilution contains 4 mcg of the base of norepinephrine. Give this solution by intravenous infusion. Insert a plastic intravenous catheter through a suitable bore needle well advanced centrally into the vein and securely fixed with adhesive tape, avoiding, if possible, a catheter tie-in technique as this promotes stasis. An IV drip chamber or other suitable metering device is essential to permit an accurate estimation of the rate of flow in drops per minute. After observing the response to an initial dose of 2 mL to 3 mL (from 8 mcg to 12 mcg of base) per minute, adjust the rate of flow to establish and maintain a low normal blood pressure (usually 80 mm Hg to 100 mm Hg systolic) sufficient to maintain the circulation to vital organs. In previously hypertensive patients, it is recommended that the blood pressure should be raised no higher than 40 mm Hg below the preexisting systolic pressure. The average maintenance dose ranges from 0.5 mL to 1 mL per minute (from 2 mcg to 4 mcg of base).

*High Dosage:* Great individual variation occurs in the dose required to attain and maintain an adequate blood pressure. In all cases, dosage of norepinephrine should be titrated according to the response of the patient. Occasionally much larger or even enormous daily doses (as high as 68 mg base or 17 ampules) may be necessary if the patient remains hypotensive, but occult blood volume depletion should always be suspected and corrected when present. Central venous pressure monitoring is usually helpful in detecting and treating this situation.

*Fluid Intake:* The degree of dilution depends on clinical fluid volume requirements. If large volumes of fluid (dextrose) are needed at a flow rate that would involve an excessive dose of the pressor agent per unit of time, a solution more dilute than 4 mcg per mL should be used. On the other hand, when large volumes of fluid are clinically undesirable, a concentration greater than 4 mcg per mL may be necessary.

*Duration of Therapy:* The infusion should be continued until adequate blood pressure and tissue perfusion are maintained without therapy. Infusions of norepinephrine should be reduced gradually, avoiding abrupt withdrawal. In some of the reported cases of vascular collapse due to acute myocardial infarction, treatment was required for up to six days.

### **Adjunctive Treatment in Cardiac Arrest**

Infusions of norepinephrine are usually administered intravenously during cardiac resuscitation to restore and maintain an adequate blood pressure after an effective heartbeat and ventilation have been established by other means. [Norepinephrine's powerful beta-adrenergic stimulating action is also thought to increase the strength and effectiveness of systolic contractions once they occur.]

*Average Dosage:* To maintain systemic blood pressure during the management of cardiac arrest, norepinephrine is used in the same manner as described under Restoration of Blood Pressure in Acute Hypotensive States.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to use, whenever solution and container permit.

Do not use the solution if its color is pinkish or darker than slightly yellow or if it contains a precipitate.

Avoid contact with iron salts, alkalis, or oxidizing agents.

### **HOW SUPPLIED**

Norepinephrine Bitartrate Injection, USP, contains the equivalent of 4 mg base of norepinephrine per each 4 mL ampule (1 mg/mL), and is available as follows:

|          |                  |                                       |
|----------|------------------|---------------------------------------|
| AIN00610 | NDC 36000-162-10 | Ampules of 4 mL in shelf carton of 10 |
|----------|------------------|---------------------------------------|

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Retain in carton until time of use.

Protect from light.

Manufactured for:

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

Manufactured by:

Sintetica S.A., Switzerland

67797 03

Issue: 2018-10-31

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 4mg/mL**

AIN00610 NDC: 36000-162-01

**Norepinephrine Bitartrate Injection, USP**

**4 mg/4 mL (1 mg per mL)** Rx Only

**CONTAINS NO SULFITES - PRESERVATIVE FREE**

Each mL contains: norepinephrine bitartrate equal to 1mg norepinephrine base.

**FOR IV INFUSION ONLY - DILUTE BEFORE USE.  
PROTECT FROM LIGHT.**

Directions: See Insert.

**BAXTER**

Deerfield, IL 60015 USA

68797 02

2018-10-31

AIN00610

NDC: 36000-162-01

**Norepinephrine Bitartrate Injection, USP**

**4 mg/4 mL (1 mg per mL)**

RX only

**CONTAINS NO SULFITES – PRESERVATIVE FREE**

Each mL contains: norepinephrine bitartrate equal to 1mg norepinephrine base.

**FOR IV INFUSION ONLY - DILUTE BEFORE USE.  
PROTECT FROM LIGHT.**

Directions: See Insert.



68797 02

***Baxter***

LOT:

EXP.:

Deerfield, IL 60015 USA

2018-10-31

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 4mg/mL**

STERILE INJECTION AIN00610 NDC : 36000-162-10 10 Ampules 4 mL Each

**Norepinephrine Bitartrate Injection, USP** Rx only

4 mg/4 mL (1 mg per mL)

CONTAINS NO SULFITES – PRESERVATIVE FREE

**Warning:** This is a potent drug. Dosage should be controlled by frequent determination of blood pressure. Do not leave patient unattended during administration. Avoid extravasation. Read package insert carefully.

**FOR INTRAVENOUS INFUSION ONLY. DILUTE BEFORE USE. DISCARD UNUSED PORTION.**

Usual Dosage and Dilution Information - See package insert.

Do not use the solution if its color is pinkish or darker than slightly yellow or if it contains a precipitate.

Avoid contact with iron salts, alkalis, or oxidizing agents.

Each mL contains: norepinephrine bitartrate, equivalent to 1 mg norepinephrine base, sodium chloride for isotonicity.

The air in the ampules has been displaced by nitrogen gas.

**These ampules are Easy Break Ampules.**

**Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F)**

**[see USP Controlled Room Temperature].**

**Retain in carton until time of use. PROTECT FROM LIGHT.**

Manufactured by:

**Sintetica S.A.**

Switzerland

Manufactured for:

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

66797 02

2018-10-31

66797 02  
2018-10-31

Manufactured by:  
Sintetica S.A.  
Switzerland

Manufactured for:  
Baxter Healthcare Corporation  
Deerfield, IL 60015 USA

STERILE INJECTION

AIN00610

NDC : 36000-162-10

10 Ampules 4 mL Each

# Norepinephrine Bitartrate Injection, USP

Rx only

4 mg/4 mL (1 mg per mL)

CONTAINS NO SULFITES – PRESERVATIVE FREE

**Warning:** This is a potent drug. Dosage should be controlled by frequent determination of blood pressure. Do not leave patient unattended during administration. Avoid extravasation. Read package insert carefully.

**FOR INTRAVENOUS INFUSION ONLY. DILUTE BEFORE USE. DISCARD UNUSED PORTION.**

Usual Dosage and Dilution Information - See package insert.

Do not use the solution if its color is pinkish or darker than slightly yellow or if it contains a precipitate.

Avoid contact with iron salts, alkalis, or oxidizing agents.

Each mL contains: norepinephrine bitartrate, equivalent to 1 mg norepinephrine base, sodium chloride for isotonicity.

The air in the ampules has been displaced by nitrogen gas.

**These ampules are Easy Break Ampules.**

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F)

[see USP Controlled Room Temperature].

Retain in carton until time of use. PROTECT FROM LIGHT.

GTIN (01)  
LOT (10)  
EXPIRY (17)  
SERIAL (21)

00000000000000  
LLLLL  
YYMMDD  
0000000000000000



**Baxter**

## Norepinephrine Bitartrate Injection, USP

10 Ampules 4 mL Each

CONTAINS NO SULFITES – PRESERVATIVE FREE



3 36000 16210 4

STERILE INJECTION AIN00610 NDC : 36000-162-10 10 Ampules 4 mL Each

## Norepinephrine Bitartrate Injection, USP Rx only

4 mg/4 mL (1 mg per mL)

CONTAINS NO SULFITES – PRESERVATIVE FREE

**Warning:** This is a potent drug. Dosage should be controlled by frequent determination of blood pressure. Do not leave patient unattended during administration. Avoid extravasation. Read package insert carefully.

**FOR INTRAVENOUS INFUSION ONLY. DILUTE BEFORE USE. DISCARD UNUSED PORTION.**

Usual Dosage and Dilution Information - See package insert.

Do not use the solution if its color is pinkish or darker than slightly yellow or if it contains a precipitate.

Avoid contact with iron salts, alkalis, or oxidizing agents.

Each mL contains: norepinephrine bitartrate, equivalent to 1 mg norepinephrine base, sodium chloride for isotonicity.

The air in the ampules has been displaced by nitrogen gas.

**These ampules are Easy Break Ampules.**

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F)

[see USP Controlled Room Temperature].

**Retain in carton until time of use. PROTECT FROM LIGHT.**

Manufactured by:

**Sintetica S.A.**

Switzerland

Manufactured for:

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

420100139-01

2023-04-14



**NOREPINEPHRINE BITARTRATE**  
norepinephrine bitartrate injection

**Product Information**

|                                |                         |                           |               |
|--------------------------------|-------------------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN PRESCRIPTION DRUG | <b>Item Code (Source)</b> | NDC:36000-162 |
| <b>Route of Administration</b> | INTRAVENOUS             |                           |               |

**Active Ingredient/Active Moiety**

| <b>Ingredient Name</b>   | <b>Basis of Strength</b> | <b>Strength</b> |
|--|--------------------------|-----------------|
| <b>NOREPINEPHRINE BITARTRATE</b> (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV) | NOREPINEPHRINE           | 1 mg<br>in 1 mL |

**Inactive Ingredients**

| <b>Ingredient Name</b>                    | <b>Strength</b> |
|---|-----------------|
| <b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X) |                 |
| <b>NITROGEN</b> (UNII: N762921K75)        |                 |

**Packaging**

| <b>#</b> | <b>Item Code</b> | <b>Package Description</b>                           | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|----------|------------------|--|-----------------------------|---------------------------|
| 1        | NDC:36000-162-10 | 10 in 1 CARTON                                       | 04/01/2012                  |                           |
| 1        | NDC:36000-162-01 | 04 mL in 1 AMPULE; Type 0: Not a Combination Product |                             |                           |

**Marketing Information**

| <b>Marketing Category</b> | <b>Application Number or Monograph Citation</b> | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|---------------------------|---|-----------------------------|---------------------------|
| ANDA                      | ANDA040859                                      | 04/01/2012                  |                           |

**Labeler** - Baxter Healthcare Corporation (005083209)**Registrant** - Baxter Healthcare Corporation (005083209)**Establishment**

| <b>Name</b>    | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b>  |
|----------------|----------------|---------------|---|
| SINTETICA S.A. |                | 480895478     | ANALYSIS(36000-162) , LABEL(36000-162) , MANUFACTURE(36000-162) , PACK(36000-162) |

**Establishment**

| <b>Name</b>  | <b>Address</b> | <b>ID/FEI</b> | <b>Business Operations</b>                   |
|--------------|----------------|---------------|--|
| SINTETICA SA |                | 480601586     | ANALYSIS(36000-162) , MANUFACTURE(36000-162) |

