

**EFFERVESCENT COLD RELIEF- aspirin, chlorpheniramine maleate, phylephrine bitartrate tablet, effervescent  
Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**TopCare Effervescent Cold Relief**

**Active ingredients**

**Active ingredients**

**(in each effervescent tablet) . . . . . Purpose**

- Aspirin 325 mg (NSAID)\*. . . . . Pain reliever/fever reducer
- Chlorpheniramine maleate 2 mg . . . . . Antihistamine
- Phenylephrine Bitartrate 7.8 mg . . . . . Nasal decongestant

\*Nonsteroidal anti-inflammatory drug

**Purpose**

**(in each effervescent tablet) . . . . . Purpose**

- Aspirin 325 mg (NSAID)\*. . . . . Pain reliever/fever reducer
- Chlorpheniramine maleate 2 mg . . . . . Antihistamine
- Phenylephrine Bitartrate 7.8 mg . . . . . Nasal decongestant

\*Nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves these symptoms due to the common cold
- sneezing ■ nasal congestion ■ sore throat
- headache ■ minor aches and pains ■ runny nose
- sinus congestion and pressure
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do Not Use**

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients.

### **Ask a doctor before use if**

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

### **Ask a doctor before use if you have**

**Ask a doctor before use if you have** ■ asthma ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** ■ you are taking sedatives or tranquilizers ■ presently taking a prescription drug

■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

**When using this product** ■ do not exceed recommended dosage

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

### **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Keep out of reach of children.**

### **In case of overdose**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours
- do not take more than 8 tablets in 24 hours

■ *children under 12 years: ask a doctor*

### **Other information**

**each tablet contains:** sodium 464 mg

■ phenylketonurics: contains phenylalanine 9 mg per tablet

■ store at room temperature (59°-86°F)

### **Inactive ingredients**

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

### **Questions or Comments?**

**Call 1-888-423-0139**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**



**Drug Facts (continued)**

for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer or if you are allergic to aspirin. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients, ask a doctor before use. If stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn or you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, ask a doctor before use. If you have asthma, glaucoma, diabetes, thyroid disease, or trouble breathing due to an enlarged prostate gland, ask a doctor before use. If you have asthma, glaucoma, diabetes, or a bleeding problem such as emphysema or chronic bronchitis, been placed on a sodium-restricted diet, ask a doctor or pharmacist before use. If you are taking sedatives or tranquilizers, or presently taking a prescription drug, ask a doctor or pharmacist before use. If you are taking sedatives or tranquilizers, or presently taking a prescription drug, do not exceed recommended dosage.

**When using this product** do not exceed recommended dosage. You may get drowsy or avoid alcoholic drinks. Excess may occur, especially in children. Alcohol, sedatives and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery.

**STOP use and ask a doctor** if you experience any of the following signs of stomach bleeding: feel faint, have bloody or black stools, have stomach pain that does not get better.

**Drug Facts**

**Active Ingredients**  
(in each effervescent tablet)  
Aspirin 325 mg (NSAID)  
Chlorpheniramine maleate 2 mg  
Phenylephrine bitartrate 7.5 mg  
Mometasone furoate nasal spray 0.05 mg

**Purpose**  
Pain reliever/fever reducer  
Allergy medicine

**Uses**  
temporarily relieves these symptoms due to the common cold:  
sore throat  
runny nose  
nasal congestion  
fever  
temporarily reduces fever

**Warnings**  
Hayfe's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor before these symptoms could be an early sign of Hayfe's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives or facial swelling.  
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The stomach bleeding problems may increase drowsiness. Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product. Take more or

**TopCare** health

# Effervescent Cold Relief

ASPIRIN (NSAID)/PAIN RELIEVER-FEVER REDUCER  
CHLORPHENIRAMINE MALEATE/ANTIHISTAMINE  
PHENYLEPHRINE BITARTRATE/NASAL DECONGESTANT

20 EFFERVESENT TABLETS ORIGINAL FLAVOR

**TopCare** health

NDC 36800-768-20

COMPARE TO ALKA-SELTZER PLUS® COLD ACTIVE INGREDIENTS\*

# Effervescent Cold Relief

ASPIRIN (NSAID)/PAIN RELIEVER-FEVER REDUCER  
CHLORPHENIRAMINE MALEATE/ANTIHISTAMINE  
PHENYLEPHRINE BITARTRATE/NASAL DECONGESTANT

**FAST RELIEF OF:**

- Nasal Congestion
- Runny Nose
- Headache & Body Aches
- Sore Throat
- Sinus Pressure

20 EFFERVESENT TABLETS ORIGINAL FLAVOR

**Drug Facts (continued)**

an allergic reaction occurs. Seek medical help right away

pain or nasal congestion gets worse or lasts more than 7 days

fever gets worse or lasts more than 3 days

new symptoms occur

redness or swelling is present

ringing in the ears or a loss of hearing occurs

nervousness, dizziness or sleeplessness occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**

do not use more than directed (see overdose warning)

adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours

do not take more than 8 tablets in 24 hours

children under 12 years: ask a doctor

**Drug Facts (continued)**

**Other information**  
each tablet contains: sodium 464 mg

phenylketonurics: contains phenylalanine 9 mg per tablet

store at room temperature (59°-86°F)

**Inactive ingredients**  
acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments?**  
Call 1-888-423-0139

\*This product is not manufactured, distributed or licensed by Bayer, the maker of Alka-Seltzer Plus® Cold.

DISTRIBUTED BY TOPCOO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007  
© TOPCOO TOWARZ21 QUESTIONS? 1-888-423-0139  
topcare@topcoo.com www.topcarebrand.com

Visit here for more information:  
<http://topbrnds.com/48zqf>

This Top Care® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

QUALITY GUARANTEED

PN: 87360 TC C3  
REV: 05/21

0 36800 06711 0

**DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED**

# EFFERVESENT COLD RELIEF

aspirin, chlorpheniramine maleate, phylephrine bitartrate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-768
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE BITARTRATE</b> (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-768-20	10 in 1 CARTON	07/01/2010	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/01/2010	

---

**Labeler** - Topco Associates LLC (006935977)

Revised: 11/2021

Topco Associates LLC