

**MUCINEX FAST-MAX COLD, FLU AND SORE THROAT- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride capsule, liquid filled**  
**RB Health (US) LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Mucinex® Fast-Max®**

**Cold, Flu & Sore Throat**

**Drug Facts**

<b>Active ingredients (in each liquid gel)</b>	<b>Purposes</b>
<b>Acetaminophen 325 mg</b>	<b>Pain reliever/fever reducer</b>
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - headache
  - cough
  - minor aches and pains
  - sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not use more than directed**

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick

medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

### **Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat

### **Inactive ingredients**

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, shellac, sorbitol sorbitan solution, titanium dioxide, water

### **Questions?**

**1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224  
Made in China

**PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton**

***Fast Dissolving Liquid Gels!***

**MAXIMUM STRENGTH**

NDC 63824-591-16

**Mucinex®**

***FAST-MAX*®**

**COLD, FLU**

**& SORE THROAT**

**Acetaminophen** – Pain Reliever/Fever Reducer

Dextromethorphan HBr – Cough Suppressant

Guaifenesin – Expectorant • Phenylephrine HCl – Nasal Decongestant

- Controls Cough, Thins & Loosens Mucus
- Relieves Nasal & Chest Congestion
- Relieves Headache & Fever

16

**LIQUID GELS**

**(Liquid Filled Capsules)**

DAY TIME

FOR AGES 12+

**MAXIMUM STRENGTH**



**COLD, FLU & SORE THROAT**

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

**16 LIQUID GELS**

*Fast Dissolving Liquid Gels!*

**MAXIMUM STRENGTH** NDC 63824-591-16



**COLD, FLU & SORE THROAT**

Maximum Strength per 4-hour dose

Do not take more than a total of 12 liquid gels in a 24-hour period.

Take only as directed.

Keep carton for complete product information.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Acetaminophen** – Pain Reliever/Fever Reducer  
**Dextromethorphan HBr** – Cough Suppressant  
**Guaifenesin** – Expectorant • **Phenylephrine HCl** – Nasal Decongestant

- ✓ Controls Cough, Thins & Loosens Mucus
- ✓ Relieves Nasal & Chest Congestion
- ✓ Relieves Headache & Fever



Also try the Max Strength, Multi-Symptom Mucinex® Fast-Max® Night Time products for night time cold & flu symptom relief

**16 LIQUID GELS**  
**(Liquid Filled Capsules)**

HEALTH • HYGIENE • HOME

[www.mucinex.com](http://www.mucinex.com)  
 Patents: [www.rb.com/patents](http://www.rb.com/patents)  
 Dist. by: RB Health (US)  
 Parsippany, NJ 07054-0224  
 Made in China  
 ©2019 RB Health  
 021819 8357060

**DAY TIME**  
**FOR AGES 12+**



### Drug Facts

#### Active ingredients (in each liquid gel)

Acetaminophen 325 mg	..... Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	..... Cough suppressant
Guaifenesin 200 mg	..... Expectorant
Phenylephrine HCl 5 mg	..... Nasal decongestant

#### Purposes

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - headache
  - minor aches and pains
  - temporarily reduces fever
  - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- cough
- sore throat

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- high blood pressure

### Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

#### Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

#### Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, shellac, sorbitol sorbitan solution, titanium dioxide, water

#### Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.





## MUCINEX FAST-MAX COLD, FLU AND SORE THROAT

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63824-591
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	26mm
<b>Flavor</b>		<b>Imprint Code</b>	PC26
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:63824-591-16	2 in 1 CARTON	07/15/2015	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/15/2015	

**Labeler** - RB Health (US) LLC (081049410)

Revised: 7/2022

RB Health (US) LLC