

NAPROXEN SODIUM- naproxen sodium tablet

Walgreens

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

***nonsteroidal anti-inflammatory drug**

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular ache
 - backache
 - menstrual cramp
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

take with food or milk if stomach upset occurs.

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing

- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work
- adults and children 12 years and older:
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** sodium 20 mg
- store at 20° to 25°C (68° to 77°F). excursions permitted between 15°C and 30°C (59°F to 86°F)[See USP Controlled Room Temperature]. Avoid high humidity and excessive heat above 40°C (104°F). Protect from Light.
- read all directions and warnings before use. Keep carton.
- swallow whole: do not crush, chew or dissolve

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, polyethylene glycol, sorbitan, sorbitol, white pharmaceutical ink

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Aleve® Liquid Gels††

All Day Pain Relief

NAPROXEN SODIUM

CAPSULES, 220 mg (NSAID) /

PAIN RELIEVER / FEVER REDUCER

12 HOUR

LIQUID GELS**

(**Liquid-filled capsules)

††This product is not manufactured or distributed by Bayer Healthcare, LLC, distributor of Aleve® Liquid Gels.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**


**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT
INFORMATION**

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Package Label

NDC 0363-7510-50



Compare to the active ingredient in ALEVE® Liquid Gels*


All Day Pain Relief

NAPROXEN SODIUM CAPSULES, 220 mg (NSAID) / PAIN RELIEVER / FEVER REDUCER

12 Hour

50

LIQUID GELS**
(**liquid-filled capsules)



ACTUAL SIZE

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*
*nonsteroidal anti-inflammatory drug

Purposes
Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to:
 ■ minor pain of arthritis ■ muscular aches
 ■ backache ■ menstrual cramps
 ■ headache ■ toothache
 ■ the common cold
 ■ temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if
 ■ the stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic
 ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product, take with food or milk. If stomach upset occurs.

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ you have symptoms of heart problems or stroke:
 ■ chest pain ■ trouble breathing
 ■ weakness in one part or side of body
 ■ slurred speech ■ leg swelling
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
 ■ you have difficulty swallowing
 ■ it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ drink a full glass of water with each dose
 ■ if taken with food, this product may take longer to work
 ■ adults and children 12 years and older
 ■ take 1 capsule every 8 to 12 hours while symptoms last
 ■ for the first dose you may take 2 capsules within the first hour
 ■ do not exceed 2 capsules in any 8- to 12-hour period
 ■ do not exceed 3 capsules in a 24-hour period
 ■ children under 12 years: ask a doctor


Other information
 ■ each capsule contains: sodium 20 mg
 ■ store at 20 to 25°C (68 to 77°F), excursions permitted between 15°C and 30°C (59°F to 86°F) [See USP Controlled Room Temperature]. Avoid high humidity and excessive heat above 40°C (104°F). Protect from light.
 ■ read all directions and warnings before use. Keep carton.
 ■ swallow whole; do not crush, chew, or dissolve

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, polyethylene glycol, sorbitol, sorbitol, white pharmaceutical ink (ammonium hydroxide, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, titanium dioxide)

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.


KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
 **This product is not manufactured or distributed by Bayer HealthCare, LLC, distributor of Aleve® Liquid Gels.

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com ©2022 Walgreen Co.

ITEM 917441 W10024-0322-L



PLD-A530B FC008176

WALGREENS All Day Pain Relief

NAPROXEN SODIUM

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-7510
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII: 57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTIC ACID (UNII: 33X04XA5AT)	

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics			
Color	blue	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	NP220
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-7510-40	1 in 1 BOX	01/31/2020	
1		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:0363-7510-80	1 in 1 BOX	01/31/2020	
2		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:0363-7510-20	1 in 1 BOX	01/31/2020	
3		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:0363-7510-50	1 in 1 BOX	01/31/2020	
4		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202807	01/31/2020	

Labeler - Walgreens (008965063)