

**RANITIDINE- ranitidine tablet**  
**CARDINAL HEALTH**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Ranitidine 75 mg (as ranitidine hydrochloride USP, 84 mg)

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
  - do not chew tablet
- children under 12 years: ask a doctor

## **Other information**

- do not use if carton or printed foil under cap is open or torn
- avoid excessive heat or humidity
- store at 20°-25°C (68°-77°F)
- this product is sodium and sugar free

## **Inactive ingredients**

FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

## **Questions?**

Call **1-888-375-3784**

**Read the directions, consumer information leaflet and warnings before use.  
Keep the carton. It contains important information.**

## **Bottle label**

LEADER<sup>2</sup><sub>TM</sub>

NDC 70000-0033-1

Regular Strength

# Acid Control

Ranitidine Tablets  
75 mg | Acid Reducer

Prevents & Relieves  
Heartburn Associated  
with Acid Indigestion  
and Sour Stomach

30 TABLETS  
(30 DOSES)

**TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

**IMPORTANT:** This label does not contain full product information. See carton for complete information. Read the directions, consumer information leaflet and warnings before use. Retain carton and leaflet for reference.

### Drug Facts

#### Active Ingredient (in each tablet) Purpose

Ranitidine 75 mg (as ranitidine

hydrochloride USP, 84 mg) ..... Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings** See warnings on carton

#### Directions

■ adults and children 12 years and over:

■ to relieve symptoms, swallow 1 tablet with a glass of water

■ to prevent symptoms, swallow 1 tablet with a glass of

water 30 to 60 minutes before eating food or drinking

beverages that cause heartburn

■ can be used up to twice daily (do not take more than 2

tablets in 24 hours)

■ do not chew tablet

■ children under 12 years: ask a doctor

**Other information** ■ this product is sodium and sugar free

■ do not use if carton or printed foil under cap is open or torn

■ avoid excessive heat or humidity

■ store at 20°-25°C (68°-77°F).

#### Inactive Ingredients

FD&C red #40 aluminum lake, hydroxypropylcellulose, iron oxide

black, magnesium stearate, microcrystalline cellulose,

polyethylene glycol, titanium dioxide.

**Questions?** Call 1-888-375-3784

Dist. by CAH, Dublin, OH 43017 © 2019 Cardinal Health

**100% Money Back Guarantee**

Return to place of purchase if not satisfied.

**CIN 5524665**

**REV. 6/19**

150076090

LOT

EXP

Carton label



## RANITIDINE

ranitidine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70000-0033
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ranitidine hydrochloride (UNII: BK76465IHM) (ranitidine - UNII:884KT10YB7)	ranitidine	75 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>Hypromelloses</b> (UNII: 3NXW29V3WO)	
<b>magnesium stearate</b> (UNII: 70097M6I3O)	
<b>cellulose, microcrystalline</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>ferrosoferric oxide</b> (UNII: XM0M87F357)	
<b>titanium dioxide</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	P75
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0033-1	1 in 1 CARTON	05/28/2019	09/30/2019
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:70000-0033-2	1 in 1 CARTON	05/28/2019	09/30/2019
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075294	05/28/2019	

**Labeler** - CARDINAL HEALTH (063997360)

Revised: 6/2019

CARDINAL HEALTH