

POLYSACCHARIDE-IRON- polysaccharide-iron capsule
Advance Pharmaceutical Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Polysaccharide-Iron Complex 150 mg

Drug Facts

.

Active Ingredient
(in each capsule)

Iron (as Polysaccharide Iron) 150 mg

Purpose

Iron Supplement

Uses

- Iron Supplement

WARNINGS

Do not exceed recommended dosage. The treatment of any anemic condition should be under the advice and supervision of a doctor. Occasional gastrointestinal discomfort (such as nausea) may be minimized by taking with meals. Iron-containing medication may occasionally cause black stools, constipation or diarrhea. **If pregnant or breast feeding**, ask a health professional before use.

Accidental overdose of Iron containing products is a leading cause of fatal poisoning in children under 6.

DRUG INTERACTION PRECAUTION: Since oral iron products interfere with absorption of tetracycline antibiotics, these products should not be taken within two hours of each other.

Keep out of reach of children.

In case of accidental overdose, call a doctor or poison control center immediately.

Directions

Adults only: One capsule daily or as directed by a doctor. **Do not exceed recommended dosage.**

Other Information

- store at 15-30 °C (59-86 °F). Do not expose to excessive heat or moisture.

Inactive Ingredients

Microcrystalline Cellulose, geltain, corscarmellose sodium, citric acid anhydrous, stearic acid, titanium

dioxide, D&C Yellow #10, sodium lauryl sulfate, FD&C Yellow #6, FD&C Red#40, FD&C Blue #1and white ink

Questions or Comments

**TAMPER RESISTENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

**Advance
Pharmaceutical Inc.**

100 CAPSULES

IRON SUPPLEMENT

**Polysaccharide-Iron
Complex 150 mg**

Poly-Fe^x

**TAMPER RESISTANT - DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

DIRECTIONS: Adults only: One capsule daily or as directed
by a doctor. **Do not exceed recommended dosage.**

Supplement Facts

Serving Size 1 Capsule

Amount Per Serving	% Daily Value
Iron (as Polysaccharide Iron) 150 mg	833%

Other ingredients: Microcrystalline cellulose, gelatin,
croscarmellose sodium, citric acid anhydrous, stearic acid,
titanium dioxide, D&C Yellow # 10, sodium lauryl sulfate,
FD&C Yellow # 6, FD&C Red # 40, FD&C Blue # 1 and white ink

WARNING: Accidental overdose of iron-containing products
is a leading cause of fatal poisoning in children under 6. **Keep
this product out of reach of children.** In case of accidental
overdose, call a doctor or poison control center immediately.

WARNINGS: Do not exceed recommended dosage. The
treatment of any anemic condition should be under the advice
and supervision of a doctor. Occasional gastrointestinal
discomfort (such as nausea) may be minimized by taking with
meals. Iron-containing medication may occasionally cause
black stools, constipation or diarrhea.

If pregnant or breast-feeding, ask a health professional before use.

DRUG INTERACTION PRECAUTION: Since oral iron products
interfere with absorption of tetracycline antibiotics, these
products should not be taken within two hours of each other.

Questions or comments?

Call **631-981-4600**, 8:30 am – 4:30 pm ET, Monday – Friday

Store between 15°-30°C (59°-86°F). Do not expose to
excessive heat or moisture.

Manufactured by: **Advance Pharmaceutical Inc.**
Holtsville, NY 11742, USA



LA0415

Lot No.:

Exp. Date:

NDC

POLYSACCHARIDE-IRON

polysaccharide-iron capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:17714-077
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IRON (UNII: E1UOL152H7) (IRON - UNII:E1UOL152H7)	IRON	150 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
GELATIN (UNII: 2G86QN327L)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	

Product Characteristics

Color	orange	Score	score with uneven pieces
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	077
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17714-077-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/26/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/26/2015	

Labeler - Advance Pharmaceutical Inc. (078301063)

