

**DOXERCALCIFEROL- doxercalciferol capsule**  
**Rising Pharmaceuticals, Inc.**

-----  
**Doxercalciferol Capsules**

**DESCRIPTION**


Doxercalciferol, the active ingredient in doxercalciferol capsule, is a synthetic vitamin D<sub>2</sub> analog that undergoes metabolic activation *in vivo* to form 1 $\alpha$ ,25-dihydroxyvitamin D<sub>2</sub> (1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub>), a naturally occurring, biologically active form of vitamin D<sub>2</sub>. Doxercalciferol Capsules is available as soft gelatin capsules containing 0.5 mcg, 1 mcg or 2.5 mcg doxercalciferol. Each capsule also contains triglyceridemedium chain (MCT), ammonium hydroxide, butylated hydroxyanisole (BHA), ethanol, propylene glycol, SDA 35A alcohol, black iron oxide, polyvinyl acetate phthalate and propylene glycol. The capsule shells contain gelatin, glycerin and titanium dioxide. In addition, the 0.5 mcg capsule shells contain D&C Yellow# 10 and FD&C Red# 40, the 1 mcg capsule shells contain FD&C Red # 40, and the 2.5 mcg capsule shells contain D&C Yellow# 10.

Doxercalciferol is a colorless crystalline compound with a calculated molecular weight of 412.66 and a molecular formula of C<sub>28</sub>H<sub>44</sub>O<sub>2</sub>. It is soluble in oils and organic solvents, but is relatively insoluble in water. Chemically, doxercalciferol is (1 $\alpha$ ,3 $\beta$ ,5Z,7E,22E)-9,10-secoergosta-5,7,10(19),22-tetraene-1,3-diol. The structural formula is presented in Figure 1 below:

---

**Figure 1: Chemical Structure of Doxercalciferol**

---



Other names frequently used for doxercalciferol are 1 $\alpha$ -hydroxyvitamin D<sub>2</sub>, 1 $\alpha$ -OH-D<sub>2</sub>, and 1 $\alpha$ -hydroxyergocalciferol.

**CLINICAL PHARMACOLOGY**

Vitamin D levels in humans depend on two sources: (1) exposure to the ultraviolet rays of the sun for conversion of 7-dehydrocholesterol in the skin to vitamin D<sub>3</sub> (cholecalciferol) and (2) dietary intake of either vitamin D<sub>2</sub> (ergocalciferol) or vitamin D<sub>3</sub>. Vitamin D<sub>2</sub> and vitamin D<sub>3</sub> must be metabolically activated in the liver and the kidney before becoming fully active on target tissues. The initial step in the activation process is the introduction of a hydroxyl group in the side chain at C-25 by the hepatic enzyme, CYP 27 (a vitamin D-25-hydroxylase). The products of this reaction are 25-(OH)D<sub>2</sub> and 25-(OH)D<sub>3</sub>, respectively. Further hydroxylation of these metabolites occurs in the mitochondria of kidney tissue, catalyzed by renal 25-hydroxyvitamin D-1- $\alpha$ -hydroxylase to produce 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub>, the primary biologically active form of vitamin D<sub>2</sub>, and 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>3</sub> (calcitriol), the biologically active form of

vitamin D<sub>3</sub>.

## **Mechanism of Action**

Calcitriol (1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>3</sub>) and 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub> regulate blood calcium at levels required for essential body functions. Specifically, the biologically active vitamin D metabolites control the intestinal absorption of dietary calcium, the tubular reabsorption of calcium by the kidney and, in conjunction with parathyroid hormone (PTH), the mobilization of calcium from the skeleton. They act directly on bone cells (osteoblasts) to stimulate skeletal growth, and on the parathyroid glands to suppress PTH synthesis and secretion. These functions are mediated by the interaction of these biologically active metabolites with specific receptor proteins in the various target tissues. In patients with chronic kidney disease (CKD), deficient production of biologically active vitamin D metabolites (due to lack of or insufficient 25-hydroxyvitamin D-1-alpha-hydroxylase activity) leads to secondary hyperparathyroidism, which contributes to the development of metabolic bone disease.

## **Pharmacokinetics and Metabolism**

Doxercalciferol is absorbed from the gastrointestinal tract and activated by CYP 27 in the liver to form 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub> (major metabolite) and 1 $\alpha$ ,24-dihydroxyvitamin D<sub>2</sub> (minor metabolite). Activation of doxercalciferol does not require the involvement of the kidneys.

In healthy volunteers, peak blood levels of 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub>, the major metabolite of doxercalciferol, are attained at 11–12 hours after repeated oral doses of 5 to 15 mcg of doxercalciferol and the mean elimination half-life of 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub> is approximately 32 to 37 hours with a range of up to 96 hours. The mean elimination half-life in patients with end-stage renal disease (ESRD) on dialysis appears to be similar. Hemodialysis causes a temporary increase in 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub> mean concentrations, presumably due to volume contraction. 1 $\alpha$ ,25-(OH)<sub>2</sub>D<sub>2</sub> is not removed from blood during hemodialysis.

## **Clinical Studies**

### **Dialysis**

The safety and effectiveness of Doxercalciferol Capsules were evaluated in two double-blind, placebo-controlled, multicenter clinical studies (Study A and Study B) in a total of 138 patients with chronic kidney disease on hemodialysis (Stage 5 CKD). Patients in Study A were an average age of 52 years (range: 22–75), were 55% male, and were 58% African-American, 31% Caucasian, and 11% Hispanic, and had been on hemodialysis for an average of 53 months. Patients in Study B were an average of 52 years (range: 27–75), were 45% male, and 99% African-American, and 1% Caucasian, and had been on hemodialysis for an average of 56 months. After randomization to two groups, eligible patients underwent an 8-week washout period during which no vitamin D derivatives were administered to either group. Subsequently, all patients received Doxercalciferol Capsules in an open-label fashion for 16 weeks followed by a double-blind period of 8 weeks during which patients received either Doxercalciferol Capsules or placebo. The initial dose of doxercalciferol during the open-label phase was 10 micrograms after each dialysis session (3 times weekly) for a total of 30 mcg per week. The dosage of doxercalciferol was adjusted as necessary by the investigator in an attempt to achieve intact parathyroid hormone (iPTH) levels within a targeted range of 150 to 300 pg/mL. The maximum dosage was limited to 20 mcg after each dialysis session (60 mcg/week). If at any time during the trial iPTH fell below 150 pg/mL, doxercalciferol was immediately suspended and restarted at a lower dosage the following week.

### **Results**

One hundred and six of the 138 patients who were treated with Doxercalciferol Capsules during the 16-week open-label phase achieved iPTH levels  $\leq$ 300 pg/mL. Ninety-four of these patients exhibited plasma iPTH levels  $\leq$ 300 pg/mL on at least 3 occasions. Eighty-seven patients had plasma iPTH levels  $<$ 150 pg/mL on at least one occasion during the open-label phase of study participation.

Mean weekly doses during the 16-week open-label period in Study A ranged from 14.8 mcg to 28.7 mcg. In Study B, the mean weekly doses during the 16-week open-label period ranged from 19.2 mcg to 28 mcg.

Decreases in plasma iPTH from baseline values were calculated using as baseline the average of the last 3 values obtained during the 8-week washout phase and are displayed in **Table 1** below:

**Table 1: iPTH Summary Data for Dialysis Patients Receiving Doxercalciferol Capsules**

		<b>iPTH (pg/mL)</b>	
		<b>means ± s.d.(n*)</b>	
		<b>p Value v. Baseline</b>	
		<b>p Value v. Placebo</b>	
		<b>Doxercalciferol Capsules</b>	<b>Placebo</b>
Study A	Baseline	797.2 ± 443.8 (30)	847.1 ± 765.5 (32)
		n.a.	
		0.97	
	Week 16 (open-label)	384.3 ± 397.8 (24)	526.5 ± 872.2 (29)
	<0.001	<0.001	
	0.72		
	Week 24 (double-blind)	404.4 ± 262.9 (21)	672.6 ± 356.9 (24)
		<0.001	0.70
		0.008	
Study B	Baseline	973.9 ± 567.0 (41)	990.4 ± 488.3 (35)
		n.a.	
		0.81	
	Week 16 (open-label)	476.1 ± 444.5 (37)	485.9 ± 443.4 (32)
	<0.001	<0.001	
	0.91		
	Week 24 (double-blind)	459.8 ± 443.0 (35)	871.9 ± 623.6 (30)
		<0.001	<0.065
		<0.001	

\*all subjects; last value to discontinuation

In both studies, iPTH levels increased progressively and significantly in 65.9% of the patients during the 8-week washout (control) period during which no vitamin D derivatives were administered. In contrast, Doxercalciferol Capsules treatment resulted in a statistically significant reduction from baseline in mean iPTH levels during the 16-week open-label treatment period in more than 93.5% of the 138 treated patients. During the double-blind period (weeks 17 to 24), the reduction in mean iPTH levels was maintained in the Doxercalciferol Capsules treatment group compared to a return to near baseline in the placebo group.

In the clinical trials, the values for iPTH varied widely from patient to patient and from week to week for individual patients. **Table 2** shows the numbers of patients within each group who achieved and maintained iPTH levels below 300 pg/mL during the open-label and double-blind phases. Seventy-four of 138 patients (53.6%) had plasma iPTH levels within the target range (150–300 pg/mL) during Weeks 14–16.

**Table 2: Number of Times iPTH ≤300 pg/mL**

	1		2		≥3	
	Doxercalciferol Capsules	Placebo	Doxercalciferol Capsules	Placebo	Doxercalciferol Capsules	Placebo

Study A	Weeks 1–16 (open-label)	2/30	2/32	0/30	0/32	22/30	23/32
	Weeks 17–24 (double-blind)	0/24	9/29	3/24	1/29	17/24	5/29
Study B	Weeks 1–16 (open-label)	2/41	4/35	1/41	0/35	29/41	21/35
	Weeks 17–24 (double-blind)	2/37	6/32	1/37	4/32	26/37	4/32

During the 8-week double-blind phase, more patients achieved and maintained the target range of values for iPTH with Doxercalciferol Capsules than with placebo.

#### Pre-dialysis

The safety and effectiveness of Doxercalciferol Capsules were evaluated in two clinical studies in 55 patients with Stage 3 or Stage 4 chronic kidney disease. Eighty-two percent of the patients were male, the average age was 64.6 years, 51% were Caucasian, 40% African-American, and the average serum iPTH level at baseline was 194.6 pg/mL. While levels of 25-(OH) vitamin D were not evaluated at baseline, retrospective assessments of stored serum revealed that the mean  $\pm$  SD serum 25-(OH) vitamin D was  $18.5 \pm 8.1$  ng/mL (range: <5 to 54 ng/mL) in the study population.

After randomization to two groups, eligible patients underwent an 8-week washout period during which no vitamin D derivatives were administered to either group. Subsequently, one group received Doxercalciferol Capsules and the other placebo during a double-blind period of 24 weeks. The initial dose of doxercalciferol was 1 mcg per day. The dosage of doxercalciferol was adjusted as necessary by the investigator in order to reduce intact parathyroid hormone (iPTH) levels to a target of  $\geq 30\%$  below post-washout baseline. The maximum dosage was limited to 3.5 mcg per day. If at any time during the trial iPTH fell below 15 pg/mL, doxercalciferol was immediately suspended and restarted at a lower dosage the following week.

#### Results

Decreases in the mean plasma iPTH from baseline values were calculated using as baseline the average of the last 2 values obtained during the 8-week washout phase. In analyses of pooled data from the two studies, iPTH levels decreased from baseline by an average of 101.4 pg/mL in the Doxercalciferol Capsules group and by 4.4 pg/mL in the placebo group ( $p < 0.001$ ). Greater reductions of iPTH with Doxercalciferol Capsules compared to placebo were observed in each study. Twenty (74%) of 27 subjects in the Doxercalciferol Capsules group achieved mean plasma iPTH suppression of  $\geq 30\%$  from baseline for the last four weeks of treatment, whereas two (7%) of the 28 subjects treated with placebo achieved this level of iPTH suppression. In the doxercalciferol-treated patients, the reductions in plasma iPTH were associated with a reduction in serum bone-specific alkaline phosphatase.

## INDICATIONS AND USAGE

### Dialysis Patients

Doxercalciferol Capsules is indicated for the treatment of secondary hyperparathyroidism in patients with chronic kidney disease on dialysis.

### Pre-Dialysis Patients

Doxercalciferol Capsules is indicated for the treatment of secondary hyperparathyroidism in patients with Stage 3 or Stage 4 chronic kidney disease.

## CONTRAINDICATIONS

Doxercalciferol Capsules should not be given to patients with a tendency towards hypercalcemia or current evidence of vitamin D toxicity.

## WARNINGS

Overdosage of any form of vitamin D, including doxercalciferol, is dangerous (see **OVERDOSAGE**). Progressive hypercalcemia due to overdosage of vitamin D and its metabolites may be so severe as to require emergency attention. Acute hypercalcemia may exacerbate tendencies for cardiac arrhythmias and seizures and may potentiate the action of digitalis drugs. Chronic hypercalcemia can lead to generalized vascular calcification and other soft-tissue calcification. The serum calcium times serum phosphorus (Ca X P) product should be maintained at  $<55 \text{ mg}^2/\text{dL}^2$  in patients with chronic kidney disease. Radiographic evaluation of suspect anatomical regions may be useful in the early detection of this condition.

Since doxercalciferol is a precursor for  $1\alpha,25\text{-(OH)}_2\text{D}_2$ , a potent metabolite of vitamin  $\text{D}_2$ , pharmacologic doses of vitamin D and its derivatives should be withheld during doxercalciferol treatment to avoid possible additive effects and hypercalcemia.

Oral calcium-based or other non-aluminum-containing phosphate binders and a low phosphate diet should be used to control serum phosphorus levels in patients with chronic kidney disease. Uncontrolled serum phosphorus exacerbates secondary hyperparathyroidism and can lessen the effectiveness of doxercalciferol in reducing blood PTH levels. If hypercalcemia occurs after initiating doxercalciferol therapy, the dose of doxercalciferol and/or calcium-containing phosphate binders should be decreased. If hyperphosphatemia occurs after initiating doxercalciferol, the dose of doxercalciferol should be decreased and/or the dose of phosphate binders increased. (See dosing recommendations for doxercalciferol under **DOSAGE AND ADMINISTRATION** section.)

Magnesium-containing antacids and doxercalciferol should not be used concomitantly in patients on chronic renal dialysis because such use may lead to the development of hypermagnesemia.

## PRECAUTIONS

### General

Active vitamin D sterols should not be used as initial treatment of nutritional vitamin D deficiency (as defined by low 25-hydroxyvitamin D). Patients should be checked and treated for nutritional vitamin D deficiency prior to initiating treatment with Doxercalciferol capsule. The principal adverse effects of treatment with doxercalciferol are hypercalcemia, hyperphosphatemia, hypercalciuria, and oversuppression of PTH (iPTH less than 150 pg/mL). Prolonged hypercalcemia can lead to calcification of soft tissues, including the heart and arteries, and hyperphosphatemia can exacerbate hyperparathyroidism. Hypercalciuria can accelerate the onset of renal failure through nephrocalcinosis. Oversuppression of PTH may lead to adynamic bone syndrome. All of these potential adverse effects should be managed by regular patient monitoring and appropriate dosage adjustments. During treatment with doxercalciferol, patients usually require dose titration, as well as adjustment in co-therapy (i.e., dietary phosphate binders) in order to maximize PTH suppression while maintaining serum calcium and phosphorus within prescribed ranges.

### Dialysis

In four adequate and well-controlled studies, the incidence of hypercalcemia and hyperphosphatemia increased during therapy with Doxercalciferol Capsules. The observed increases during doxercalciferol treatment, although occurring at a low rate, underscore the importance of regular safety monitoring of serum calcium and phosphorus levels throughout treatment. Patients with higher pre-

treatment serum levels of calcium (>10.5 mg/dL) or phosphorus (>6.9 mg/dL) were more likely to experience hypercalcemia or hyperphosphatemia. Therefore, doxercalciferol should not be given to patients with a recent history of hypercalcemia or hyperphosphatemia, or evidence of vitamin D toxicity.

### Pre-dialysis

In two clinical studies, the incidences of hypercalcemia and hyperphosphatemia during therapy with Doxercalciferol Capsules were similar to placebo therapy, and no episodes of hypercalciuria were observed. The baseline median 25-(OH) vitamin D levels of patients enrolled in these studies was 17.2 ng/mL. Ninety-three percent of patients had 25-(OH) vitamin D levels less than 30 ng/mL; 26% had 25-(OH) vitamin D levels  $\geq 20$  to <30 ng/mL; 58% had levels >10 to <20 ng/mL; 7% had levels >5 to <10 ng/mL; and 2% had levels <5 ng/mL. The incidences of hypercalcemia, hyperphosphatemia, and hypercalciuria in patients treated with doxercalciferol for hyperparathyroidism related to pre-dialysis renal insufficiency has not been fully studied when 25-(OH) vitamin D levels are greater than or equal to 30 ng/mL.

### Information for the Patient

The patient, spouse, or guardian should be informed about compliance with dosage instructions, adherence to instructions about diet, calcium supplementation, and avoidance of the use of nonprescription drugs without prior approval from their physician. Patients should also be carefully informed about the symptoms of hypercalcemia (see **ADVERSE REACTIONS** section). Patients' total combined elemental calcium intake (dietary and phosphate binder) should not exceed 2 g daily.

### Laboratory Tests

Serum or plasma iPTH and serum calcium, phosphorus, and alkaline phosphatase should be determined periodically. In the early phase of treatment for dialysis patients, iPTH, serum calcium, and serum phosphorus should be determined prior to initiation of doxercalciferol treatment and weekly thereafter. For pre-dialysis patients, serum levels of calcium and phosphorus and plasma levels of iPTH should be monitored at least every two weeks for 3 months after initiation of doxercalciferol therapy or following dose-adjustments in doxercalciferol therapy, then monthly for 3 months, and every 3 months thereafter.

### Drug Interactions

Specific drug interaction studies have not been conducted. Cholestyramine has been reported to reduce intestinal absorption of fat-soluble vitamins; therefore, it may impair intestinal absorption of doxercalciferol. Magnesium-containing antacids and doxercalciferol should not be used concomitantly because such use may lead to the development of hypermagnesemia (see **WARNINGS**). The use of mineral oil or other substances that may affect absorption of fat may influence the absorption and availability of doxercalciferol. Although not examined specifically, enzyme inducers (such as glutethimide and phenobarbital) may affect the 25-hydroxylation of doxercalciferol and may necessitate dosage adjustments. Cytochrome P450 inhibitors (such as ketoconazole and erythromycin) may inhibit the 25-hydroxylation of doxercalciferol. Hence, formation of the active doxercalciferol moiety may be hindered.

### Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 104-week carcinogenicity study in rats, there was an increased incidence of benign and malignant adrenal pheochromocytomas in both males and females at oral doses of 0.04, 0.13 and 0.39 mcg/kg/day ( $\leq 1$  times the human exposure in pre-dialysis patients with a maximum recommended dose of 3.5 mcg/day or 24.5 mcg/week). This increased incidence of pheochromocytomas in rats may be due to altered calcium homeostasis by doxercalciferol. No evidence of genetic toxicity was observed in an *in vitro* bacterial mutagenicity assay (Ames test) or a mouse lymphoma gene mutation assay. Doxercalciferol caused structural chromatid and chromosome aberrations in an *in vitro* human

lymphocyte clastogenicity assay with metabolic activation. However, doxercalciferol was negative in an *in vivo* mouse micronucleus clastogenicity assay. Doxercalciferol had no effect on male or female fertility in rats at oral doses up to 2.5 mcg/kg/day (approximately 3 times the maximum recommended human dose of 60 mcg/week based on mcg/m<sup>2</sup> body surface area).

### Use in Pregnancy

#### Pregnancy Category B

Reproduction studies in rats and rabbits, at doses up to 20 mcg/kg/day and 0.1 mcg/kg/day (approximately 25 times and less than the maximum recommended human dose of 60 mcg/week based on mcg/m<sup>2</sup> body surface area, respectively) have revealed no teratogenic or fetotoxic effects due to doxercalciferol. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### Nursing Mothers

It is not known whether doxercalciferol is excreted in human milk. Because other vitamin D derivatives are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from doxercalciferol, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

### Pediatric Use

Safety and efficacy of doxercalciferol in pediatric patients have not been established.

### Geriatric Use

Of the 138 patients treated with Doxercalciferol Capsules in two Phase 3 clinical studies, 30 patients were 65 years or over. In these studies, no overall differences in efficacy or safety were observed between patients 65 years or older and younger patients.

### Hepatic Insufficiency

Since patients with hepatic insufficiency may not metabolize doxercalciferol appropriately, the drug should be used with caution in patients with impaired hepatic function. More frequent monitoring of iPTH, calcium, and phosphorus levels should be done in such individuals.

## ADVERSE REACTIONS

### Dialysis

Doxercalciferol Capsules have been evaluated for safety in clinical studies in 165 patients with chronic kidney disease on hemodialysis. In two placebo-controlled, double-blind, multicenter studies, discontinuation of therapy due to any adverse event occurred in 2.9% of 138 patients treated with Doxercalciferol Capsules for four to six months (dosage titrated to achieve target iPTH levels, see **CLINICAL PHARMACOLOGY/Clinical Studies**) and in 3.3% of 61 patients treated with placebo for two months. Adverse events occurring in the Doxercalciferol Capsules group at a frequency of 2% or greater and more frequently than in the placebo group are presented in **Table 3** below:

**Table 3: Adverse Events Reported by ≥2% of Doxercalciferol Capsules Treated Patients and More Frequently Than Placebo During the Double-blind Phase of Two Clinical Studies**

Adverse Event	Doxercalciferol Capsules (n=61) %	Placebo (n=61) %
<b>Body as a Whole</b>		

Abscess	3.3	0.0
Headache	27.9	18.0
Malaise	27.9	19.7
<b>Cardiovascular System</b>		
Bradycardia	6.6	4.9
<b>Digestive System</b>		
Anorexia	4.9	3.3
Constipation	3.3	3.3
Dyspepsia	4.9	1.6
Nausea/Vomiting	21.3	19.7
<b>Musculoskeletal System</b>		
Arthralgia	4.9	0.0
<b>Metabolic and Nutritional</b>		
Edema	34.4	21.3
Weight increase	4.9	0.0
<b>Nervous System</b>		
Dizziness	11.5	9.8
Sleep disorder	3.3	0.0
<b>Respiratory System</b>		
Dyspnea	11.5	6.6
<b>Skin</b>		
Pruritus	8.2	6.6

A patient who reported the same medical term more than once was counted only once for that medical term.

### Pre-dialysis

Doxercalciferol Capsules have been evaluated for safety in clinical studies in 55 patients (27 active and 28 placebo) with chronic kidney disease, Stages 3 or 4. In two placebo-controlled, double-blind, multicenter studies, discontinuation of therapy due to any adverse event occurred in one (3.7%) of 27 patients treated with Doxercalciferol Capsules for 24 weeks (dosage titrated to achieve target iPTH levels, see **CLINICAL PHARMACOLOGY/Clinical Studies**) and in three (10.7%) of 28 patients treated with placebo for 24 weeks. Adverse events occurring in the Doxercalciferol Capsules group at a frequency of 5% or greater and more frequently than in the placebo group are as follows: **Body as a Whole** – Infection, Chest Pain; **Digestive System** – Constipation, Dyspepsia; **Hematologic and Lymphatic** – Anemia; **Metabolic and Nutritional** – Dehydration; **Nervous System** – Depression, Hypertonia, Insomnia, Paresthesia; **Respiratory System** – Cough increased, Dyspnea, Rhinitis.

Potential adverse effects of doxercalciferol are, in general, similar to those encountered with excessive vitamin D intake. The early and late signs and symptoms of vitamin D intoxication associated with hypercalcemia include:

#### Early

Weakness, headache, somnolence, nausea, vomiting, dry mouth, constipation, muscle pain, bone pain, metallic taste, and anorexia.

#### Late

Polyuria, polydipsia, anorexia, weight loss, nocturia, conjunctivitis (calcific), pancreatitis, photophobia, rhinorrhea, pruritus, hyperthermia, decreased libido, elevated blood urea nitrogen (BUN), albuminuria, hypercholesterolemia, elevated serum aspartate transaminase (AST) and alanine transaminase (ALT), ectopic calcification, hypertension, cardiac arrhythmias, sensory disturbances, dehydration, apathy,

arrested growth, urinary tract infections, and, rarely, overt psychosis.

**To report SUSPECTED ADVERSE REACTIONS, contact Rising Pharmaceuticals, Inc. at 1-866-562-4597 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

## OVERDOSAGE

Administration of doxercalciferol to patients in excess doses can cause hypercalcemia, hypercalciuria, hyperphosphatemia, and oversuppression of PTH secretion leading in certain cases to adynamic bone disease. High intake of calcium and phosphate concomitant with doxercalciferol may lead to similar abnormalities. High levels of calcium in the dialysate bath may contribute to hypercalcemia.

### Treatment of Hypercalcemia and Overdosage

General treatment of hypercalcemia (greater than 1 mg/dL above the upper limit of the normal range in dialysis patients; >10.7 mg/dL in pre-dialysis patients) consists of immediate suspension of doxercalciferol therapy, institution of a low calcium diet, and withdrawal of calcium supplements. Serum calcium levels should be determined at least weekly until normocalcemia ensues. Hypercalcemia usually resolves in 2 to 7 days. When serum calcium levels have returned to within normal limits, doxercalciferol therapy may be reinstated at a dose that is lower (at least 2.5 mcg in dialysis patients and 0.5 mcg in pre-dialysis patients) than prior therapy. In dialysis patients, serum calcium levels should be obtained weekly after all dosage changes and during subsequent dosage titration. Persistent or markedly elevated serum calcium levels may be corrected by dialysis against a reduced calcium or calcium-free dialysate.

### Treatment of Accidental Overdosage of Doxercalciferol

The treatment of acute accidental overdosage of doxercalciferol should consist of general supportive measures. If drug ingestion is discovered within a relatively short time (10 minutes), induction of emesis or gastric lavage may be of benefit in preventing further absorption. If drug ingestion is discovered later than 10 minutes post-ingestion, the administration of mineral oil may promote its fecal elimination. Serial serum electrolyte determinations (especially calcium), rate of urinary calcium excretion, and assessment of electrocardiographic abnormalities due to hypercalcemia should be obtained. Such monitoring is critical in patients receiving digitalis. Discontinuation of supplemental calcium and institution of a low calcium diet are also indicated in accidental overdosage. If persistent and markedly elevated serum calcium levels occur, treatment with standard medical care should be followed, as needed. Based on similarities between doxercalciferol and its active metabolite,  $1\alpha,25\text{-(OH)}_2\text{D}_2$ , it is expected that doxercalciferol is not removed from the blood by dialysis.

## DOSAGE AND ADMINISTRATION

### Adult Administration

The optimal dose of doxercalciferol must be carefully determined for each patient. **Table 4** provides the current recommended therapeutic target levels for iPTH in patients with chronic kidney disease:

**Table 4: Target Range of Intact Plasma PTH by Stage of CKD**

CKD Stage	GFR (mL/min/1.73m <sup>2</sup> )	Target "intact" PTH (pg/mL)
3	30 to 59	35 to 70
4	15 to 29	70 to 110
5	<15 (or dialysis)	150 to 300

From Table 15 of National Kidney Foundation, *K/DOQI Clinical Practice Guidelines for Bone Metabolism and Disease in Chronic Kidney Disease*. Am J Kidney Dis 42:S1-S202, 2003 (suppl 3)

## Dialysis

The recommended initial dose of Doxercalciferol Capsules is 10 mcg administered three times weekly at dialysis (approximately every other day). The initial dose should be adjusted, as needed, in order to lower blood iPTH into the range of 150 to 300 pg/mL. The dose may be increased at 8-week intervals by 2.5 mcg if iPTH is not lowered by 50% and fails to reach the target range. The maximum recommended dose of Doxercalciferol Capsules is 20 mcg administered three times a week at dialysis for a total of 60 mcg per week. Drug administration should be suspended if iPTH falls below 100 pg/mL and restarted one week later at a dose that is at least 2.5 mcg lower than the last administered dose. During titration, iPTH, serum calcium, and serum phosphorus levels should be obtained weekly. If hypercalcemia, hyperphosphatemia, or a serum calcium times serum phosphorus product greater than  $55 \text{ mg}^2/\text{dL}^2$  is noted, the dose of doxercalciferol should be decreased or suspended and/or the dose of phosphate binders should be appropriately adjusted. If suspended, the drug should be restarted at a dose that is at least 2.5 mcg lower.

Dosing must be individualized and based on iPTH levels with monitoring of serum calcium and serum phosphorus levels. The following is a suggested approach in dose titration:

**Table 5: Dialysis Dosing Recommendations**

<b>Initial Dosing</b>	
<u>iPTH Level</u>	<u>Doxercalciferol Capsules Dose</u>
>400 pg/mL	10 mcg three times per week at dialysis
<b>Dose Titration</b>	
<u>iPTH Level</u>	<u>Doxercalciferol Capsules Dose</u>
Above 300 pg/mL	Increase by 2.5 mcg at eight-week intervals as necessary
150 – 300 pg/mL	Maintain
<100 pg/mL	Suspend for one week, then resume at a dose that is at least 2.5 mcg lower

## Pre-dialysis

The recommended initial dose of Doxercalciferol Capsules is 1 mcg administered once daily. The initial dose should be adjusted, as needed, in order to lower blood iPTH to within target ranges (see table below). The dose may be increased at 2-week intervals by 0.5 mcg to achieve the target range of iPTH. The maximum recommended dose of Doxercalciferol Capsules is 3.5 mcg administered once per day.

Serum levels of calcium and phosphorus and plasma levels of iPTH should be monitored at least every two weeks for 3 months after initiation of doxercalciferol therapy or following dose adjustments in doxercalciferol therapy, then monthly for 3 months, and every 3 months thereafter. If hypercalcemia, hyperphosphatemia, or a serum calcium times phosphorus product greater than  $55 \text{ mg}^2/\text{dL}^2$  is noted, the dose of doxercalciferol should be decreased or suspended and/or the dose of phosphate binders should be appropriately adjusted. If suspended, the drug should be restarted at a dose that is at least 0.5 mcg lower.

Dosing must be individualized and based on iPTH levels with monitoring of serum calcium and serum phosphorus levels. **Table 6** presents a suggested approach in dose titration:

**Table 6: Pre-dialysis Dosing Recommendations**

<b>Initial Dosing</b>	
<u>iPTH Level</u>	<u>Doxercalciferol Capsules Dose</u>
>70 pg/mL (Stage 3)	1 mcg once per day

>110 pg/mL (Stage 4)

### Dose Titration

<u>iPTH Level</u>	<u>Doxercalciferol Capsules Dose</u>
Above 70 pg/mL (Stage 3) 110 pg/mL (Stage 4)	Increase by 0.5 mcg at two-week intervals as necessary
35 to 70 pg/mL (Stage 3) 70 to 110 pg/mL (Stage 4)	Maintain
<35 pg/mL (Stage 3) <70 pg/mL (Stage 4)	Suspend for one week, then resume at a dose that is at least 0.5 mcg lower

### HOW SUPPLIED

Doxercalciferol capsules 0.5 mcg are available in opaque red oval soft gel capsules with "L" imprinted on one side and plain on the other side. Bottle of 50 count NDC 64980-228-50

Doxercalciferol capsules 1 mcg are available in opaque pink oval soft gel capsules with "M" imprinted on one side and plain on the other side. Bottle of 50 count NDC 64980-229-50

Doxercalciferol capsules 2.5 mcg are available in opaque yellow oval soft gel capsules with "H" imprinted on one side and plain on the other side. Bottle of 50 count NDC 64980-230-50

Store at 20 to 25°C (68 to 77°F)

[see USP Controlled Room Temperature]

### Manufactured for:

Rising Pharmaceuticals, Inc.

Allendale, NJ 07401

Made in France

I06/16 OS248A-01-82-01

### PRINCIPAL DISPLAY PANEL - 0.5 mcg Capsule

Rising® NDC 64980-228-50

Doxercalciferol

Capsules

0.5 mcg

Rx ONLY

50 Capsules

**Rising®** NDC 64980-228-50  
**Doxercalciferol**  
**Capsules**  
**0.5 mcg**  
**50 Capsules**  
**Rx only**

**Each capsule contains:**  
0.5 mcg of doxercalciferol.

**Usual Dosage:**  
See package insert for dosage and use information.

**Keep this and all drugs out of reach of children.**

**Store at 20° to 25°C (68° to 77°F) (see USP controlled room temperature).**

**Manufactured for:**  
Rising Pharmaceuticals, Inc.  
Allendale, NJ 07401

**Made in France** 03/16  
CLB-P-751-800-16961

**N 3** 64980 22850 3

**Overprint Area**  
18 x 13 mm

**LOT NO.:**  
**EXP.:**

### PRINCIPAL DISPLAY PANEL - 1 mcg Capsule

Rising® NDC 64980-229-50

**Doxercalciferol**  
**Capsules**  
**1 mcg**  
**Rx ONLY**  
**50 Capsules**

**Rising®** NDC 64980-229-50  
**Doxercalciferol**  
**Capsules**  
**1 mcg**  
**50 Capsules** **Rx only**

**Each capsule contains:**  
1 mcg of doxercalciferol.

**Usual Dosage:**  
See package insert for dosage and use information.

**Keep this and all drugs out of reach of children.**

**Store at 20° to 25°C (68° to 77°F) [see USP controlled room temperature].**

**Manufactured for:**  
Rising Pharmaceuticals, Inc.  
Allendale, NJ 07401

**Made in France** 03/16  
CLB-P-751-800-16962

**Overprint Area**  
18 x 13 mm

LOT NO.:  
EXP.:

**PRINCIPAL DISPLAY PANEL - 2.5 mcg Capsule**

**Rising®** NDC 64980-230-50  
**Doxercalciferol**  
**Capsules**  
**2.5 mcg**  
**Rx ONLY**  
**50 Capsules**

**Rising®** NDC 64980-230-50  
**Doxercalciferol**  
**Capsules**  
**2.5 mcg**  
**50 Capsules** **Rx only**

**Each capsule contains:**  
2.5 mcg of doxercalciferol.

**Usual Dosage:**  
See package insert for dosage and use information.

**Keep this and all drugs out of reach of children.**

**Store at 20° to 25°C (68° to 77°F) [see USP controlled room temperature].**

**Manufactured for:**  
Rising Pharmaceuticals, Inc.  
Allendale, NJ 07401

**Made in France** 03/16  
CLB-P-751-800-16963

**Overprint Area**  
18 x 13 mm

LOT NO.:  
EXP.:

**DOXERCALCIFEROL**

doxercalciferol capsule

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:64980-228
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
Doxercalciferol (UNII: 3DIZ9LF5Y9) (Doxercalciferol - UNII:3DIZ9LF5Y9)	Doxercalciferol	0.5 ug

**Inactive Ingredients**

Ingredient Name	Strength
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

### Product Characteristics

Color	PINK (salmon)	Score	no score
Shape	OVAL	Size	6 mm
Flavor		Imprint Code	g
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64980-228-50	1 in 1 CARTON	09/09/2016	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201518	09/09/2016	

## DOXERCALCIFEROL

doxercalciferol capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:64980-229
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Doxercalciferol (UNII: 3DIZ9LF5Y9) (Doxercalciferol - UNII:3DIZ9LF5Y9)	Doxercalciferol	1 ug

### Inactive Ingredients

Ingredient Name	Strength
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	

GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

### Product Characteristics

Color	PINK (peach)	Score	no score
Shape	OVAL	Size	6mm
Flavor		Imprint Code	g
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64980-229-50	1 in 1 CARTON	09/09/2016	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201518	09/09/2016	

## DOXERCALCIFEROL

doxercalciferol capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:64980-230
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Doxercalciferol (UNII: 3DIZ9LF5Y9) (Doxercalciferol - UNII:3DIZ9LF5Y9)	Doxercalciferol	2.5 ug

### Inactive Ingredients

Ingredient Name	Strength
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

## Product Characteristics

<b>Color</b>	YELLOW (butter yellow)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	g
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64980-230-50	1 in 1 CARTON	09/09/2016	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201518	09/09/2016	

**Labeler** - Rising Pharmaceuticals, Inc. (041241766)

## Establishment

Name	Address	ID/FEI	Business Operations
Catalent Pharma Solutions LLC		051762268	MANUFACTURE(64980-228, 64980-229, 64980-230)

Revised: 12/2017

Rising Pharmaceuticals, Inc.