

**GENUINE FIRST AID INDUSTRIAL FIRST AID - aspirin, acetaminophen, ibuprofen, bacitracin zinc, neomycin sulfate, polymyxin b, benzalkonium chloride, lidocaine, benzalkonium chloride**

**Genuine First Aid, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Genuine First Aid Industrial Kit 122pcs**

**Active Ingredient**

Active Ingredient:

Benzalkonium Chloride 0.40%

**Purpose**

Antiseptic

**Use**

For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

**Warnings**

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

**Directions**

Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

**Inactive Ingredients**

Inactive Ingredient: Purified water

LOT/EXP:                      Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

### **Active Ingredients**

Active Ingredient: .....Bacitracin Zinc 400 units

Neomycin Sulfate 5mg ( equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

### **Purpose**

Triple Antibiotic

Uses: To help prevent infection in:  
minor cuts; scrapes; burns

### **Warnings**

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

### **Directions**

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

### **Inactive Ingredient**

water

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

Triple Antibiotic Ointment 10pcs

Net wt. 0.9g (1/32oz)

100

Triple Antibiotic

### **Active Ingredients**

ACTIVE INGREDIENTS:

Benzalkonium Chloride 0.13%

Lidocaine HCL 0.5%

### **Purpose**

Purpose: First aid antiseptic, external analgesic

### **Uses**

First aid to help prevent infection and for the temporary relief of pain and itching associated with:

Minor Cuts

Scrapes

Burns

### **Warnings**

For external use only

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week

Do not use:

in the eyes or apply over large areas of the body.

longer than 1 week unless directed by a doctor.  
in large quantities, particularly over raw surfaces or blistered areas.

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns.

When using this product, avoid contact with the eyes.

Stop use and ask a doctor if  
condition worsens  
symptoms persist for more than 7 days  
condition clears up and occurs again within a few days

Keep out of reach of Children.

If ingested, contact a Poison Control Center right away.

### **Directions**

Adults and children 2 years of age and older  
clean affected area.

apply a small amount of this product on the area 1 to 3 times daily.

may be covered with a sterile bandage

children under 2 years of age: consult a doctor

Other Information:

Store at room temperature (do not freeze).

Taper evident sealed packets.

Do not use packet if opened or torn.

### **Inactive Ingredients**

Aloe vera, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl monostearate, methylparaben, mineral oil, polyethylene glycol, propylene glycol, propylparaben, purified water, stearic acid, irolamine

LOT/EXP:

Made in CHINA

20130301

GFA Production Xiamen Co., Ltd

No. 20 Huli Industrial Park, Meixi Road, Tong'an, Xiamen, Fujian, China 361100

Tel: 86-592-7269515 Fax: 86-592-7269528 Http: //www.gfaproduction.com

Genuine First Aid Burn Cream

Antiseptic Pain Relief With Aloe

Net Wt 0.9g (1/32 oz)

Manufactured in CHINA for

Genuine First Aid.

**Active Ingredient (in each tablet)**

Ibuprofen USP (NSAID\*) 200mg

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily relieves minor aches and pains due to:

the common cold

headache

toothache

muscular aches

backache

minor pain of arthritis

menstrual cramps temporarily reduces fever

**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed**

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

**Ask a doctor before use if you are:**

taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

**When using this product**

take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

**Other information**

Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

**Inactive Ingredients**

cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

IBUPROFEN 2 Tablets

**Active Ingredient (in each tablet)**

Acetaminophen 325 mg

**Purpose**

Analgesic/antipyretic

**Uses**

temporary relief of minor aches and pains associated with:

common cold; headache; toothache; muscular aches; backache; arthritis; menstrual cramps; and reduction of fever

**Warnings**

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product

**Do not use**

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

**Inactive Ingredients**

Cornstarch, polyethylene glycol, stearic acid, povidone

**Directions**

Adults and Children  
12 years of age  
or older

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

Children 6-11 years  
of age

Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.

Children under 6  
years of age

Do not use this regular strength product.  
This will provide more than the  
recommended dose (overdose) and could  
cause serious health problems.

Store at 59-86 degree F (15-30 degree C)

tamper evident sealed packets; do not use any open or torn packets

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

GENUINE FIRST AID 2 Tablets

NON-ASPIRIN

### **Active Ingredient (in each tablet)**

Aspirin (NSAID\*) 325 mg  
\*nonsteroidal anti-inflammatory drug

### **Purpose**

Pain Reliever / fever reducer

### **Uses**

Temporarily relieves minor aches and pains associated with:  
headache ; muscular aches ; minor arthritis pain ; backache ; common cold ; toothache ; menstrual cramps  
; Temporarily reduces fever

### **Warnings**

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed

### **Do not use**

if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after



heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

under a doctor's care for any serious condition; taking any other drug

When using this product: take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

do not use more than directed

the smallest effective dose should be used

drink a full glass of water with each dose

do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

### **Inactive Ingredients**

hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

ASPIRIN 2 Tablets

# ASPIRIN

2 Tablets

**Active ingredient (in each tablet)** **Purpose**  
Aspirin (NSAID) 325 mg . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** Temporarily relieves minor aches and pains associated with  
■ headache ■ muscular aches ■ minor arthritis pain ■ backache  
■ common cold ■ toothache ■ menstrual cramps  
**Temporarily reduces fever**

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ▶

## Warnings (continued)

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs  
**Stop use and ask a doctor if**

■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing  
■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

**do not use more than directed**  
■ the smallest effective dose should be used  
■ drink a full glass of water with each dose  
■ do not take longer than 10 days, unless directed by a doctor

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information** ■ store at 59°-86°F (15°-30°C) ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets

**Inactive ingredients** hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755



GENUINE FIRST AID. 2 Tablets

# NON-ASPIRIN

**Active ingredient (in each tablet)** **Purpose**  
Acetaminophen 325 mg . . . . . Analgesic/antipyretic

## Uses

temporary relief of minor aches and pains associated with  
■ common cold ■ headache ■ toothache  
■ muscular aches ■ backache ■ arthritis  
■ menstrual cramps ■ and reduction of fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 12 tablets in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor

**Ask a doctor before use if the user has liver disease**

**Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin** ▶

## Warnings (continued)

**Stop use and ask a doctor if** ■ symptoms do not improve ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ a rare sensitivity reaction occurs

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Do not exceed recommended dosage.**

## Directions

Adults and children 12 years of age and older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

**Other information** ■ store at 59°-86°F (15°-30°C) ■ tamper evident sealed packets ■ do not use any open or torn packets

**Inactive ingredients** corn starch, hydroxypropyl methylcellulose, polyethylene glycol, pregelatinized starch, stearic acid. May contain povidone and sodium starch glycolates.

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755



# IBUPROFEN 2 Tablets

**Active ingredient (in each tablet)** Purpose  
 Ibuprofen USP (NSAID)\* 200mg . . . . . Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves minor aches and pains due to:  
 ■ the common cold ■ headache ■ toothache ■ muscular aches  
 ■ backache ■ minor pain of arthritis ■ menstrual cramps  
 temporarily reduces fever

**Warnings Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ shock ■ facial swelling ■ asthma (wheezing) ■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ taking any other drug containing an NSAID (prescription or nonprescription) ▶

**Warnings (continued)**

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information** ■ store at controlled room temperature ■ avoid excessive heat 40° C (104° F) ■ tamper evident sealed packets ■ do not use any opened or torn packets

**Inactive ingredients** cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Distributed by GENUINE FIRST AID  
 600 Cleveland Str Suite 400, Clearwater, FL 33755

Reorder TAO-001

## Genuine Triple Antibiotic

**First Aid Ointment To Help Prevent Infection**

**Each Gram Contains:**  
**Bacitracin Zinc 400 units**  
**Neomycin Sulfate 5 mg**  
**(equivalent to 3.5 mg Neomycin base)**

**Polymyxin B Sulfate 5000 units**  
**Net Wt 0.9g - (1/32 oz)**



GFA Production Xiamen Co., Ltd  
 No. 20 Hui Industrial Park, Meixi Road,  
 Tong'an, Xiamen, Fujian, China 361100  
 Weikang Ltd & Weikang Tech Consulting  
 Suite B, 29 Harley Street  
 LONDON W1G 9QR, England, United Kingdom

**Drug Facts - Triple Antibiotic**

Each Gram Contains:	Purpose:
Bacitracin Zinc.....	400 units
Neomycin Sulfate.....	5 mg (equivalent to 3.5mg Neomycin base)
Polymyxin B Sulfate .....	5000 units

**Uses:** To help prevent infections in minor cuts, scrapes or burns.

**Warnings:** For external use only  
**Do not use:** in eyes, over large areas of the body, if allergic to any of the ingredients, or for more than one week unless directed by a physician.

**Stop use and consult a doctor:** if the condition persists or gets worse, a rash or other allergic reaction develops.

**Keep out of reach of children:** If ingested, contact a Poison Control Center right away.

**Directions:** Clean affected area, apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily. May be covered with a sterile bandage.

**Other information:**  
 Store at room temperature.  
**Inactive ingredient:** Vaseline...96.41%  
 Mineral oil.....2%  
 Purified water.



Reorder BC-001

# Genuine First Aid Burn Cream

**Antiseptic Pain Relief  
With Aloe**

## ACTIVE INGREDIENTS:

**Benzalkonium**

**Chloride 0.13%**

**Lidocaine HCL 0.5%**

**Net Wt 0.9g - (1/32 oz)**



GFA Production Xiamen Co., Ltd  
No. 20 Huli Industrial Park, Meibei Road,  
Tong'an, Xiamen, Fujian, China 361100

Wellkang Ltd t/a Wellkang Tech Consulting  
Suite B, 29 Hatfield Street  
LONDON W1G 9QR, England, United Kingdom

### Drug Facts-Burn Cream

Active Ingredient: Purpose:

Benzalkonium Chloride.....0.13%  
Lidocaine HCL..... 0.5%

Purpose: First aid antiseptic,  
external analgesic.

Uses: First aid to help prevent  
infection and for the temporary relief  
of pain and itching associated with:  
minor cuts, scrapes and burns.

Warnings: For external use only.

Do not use: in eyes, in large  
quantities, over raw blistered areas,  
on deep puncture wounds, animal  
bites or serious burns, for more than  
one week.

Directions: Clean affected area.  
Apply a small amount on area, not  
more than 3 times daily. May be  
covered with a sterile bandage.

Other information:

Store at room temperature.

Keep out of reach of children.

If ingested, contact a Poison Control  
Center right away.

Inactive ingredient:Peregal-O... 2.5%  
Glycerin monostearate...2.5%  
Glycerol.....5%  
Purified water.



LOT

XXXXXXXXXX



XXXXXXXXXX



**DRUG FACTS - Antiseptic Towelette**

**Active Ingredient:** Benzalkonium Chloride 0.13%.. **Purpose:** First Aid Antiseptic

**Use:** For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

**Warnings:** For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

**Do not use:** In the eyes, or over large areas of the body.

**Directions:** Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

**Inactive ingredient:** Purified water.

LOT/EXP: XXXXXXXX

TEAR HERE

REORDER AST-001

**Antiseptic Towelette**  
**Toallitas Antisépticas**



GFA Production Xiamen Co., Ltd  
www.gfaproduction.com



GFA Production Xiamen Co., Ltd  
No. 20 Hull Industrial Park, Meid Road,  
Tongan, Xiamen, Fujian, China 361100  
Wellkang Ltd t/a Wellkang Tech Consulting  
Suite B, 29 Hanley Street  
LONDON W1G 9QR, England, United Kingdom



**Industrial First Aid Kit**  
**Botiquin de Primeros Auxilios Industrial**



• **122 pieces**

• **Exceeds ANSI Z308.1-2009 standard for the workplace**

• **Use for Car, Travel, and Home**

• **Critical Supplies for Wound Care** not found in other kits



• **122 piezas**

• **Excede estándar ANSI Z308.1-2009 para el lugar de trabajo**

• **Use en Carro, de Viaje y en el Hogar**

• **Artículos Críticos para el Cuidado de Heridas que no se Encuentran en Otros Equipos**

**READY AMERICA.**  
THE DISASTER SUPPLY PROFESSIONALS

# Industrial First Aid Kit

## Botiquin de Primeros Auxilios Industrial

**122**  
pieces  
**122** # de piezas

Qty	Item Description
24	Adhesive Bandage 3" x 3/4"
6	Adhesive Bandage 1" x 3"
12	Adhesive Bandage Fabric 1" x 3"
3	X-Large Strip
12	Junior Bandage 3/8" x 1 1/2"
2	Knuckle Fabric Bandage
2	Fingertip Fabric Bandage
2	Butterfly Wound Closure
2	Gauze Pads 2" x 2"
4	Gauze Pads 3" x 3"
1	Trauma Pad 5" x 9"
1	Gauze Roll 2"
1	First aid Tape Roll 1/2" x 5yd.
10	Antiseptic Wipes (sting free)
6	Antibiotic Ointment
6	First Aid/Burn Cream

Qty	Item Description
1	Moleskin 2" x 2"
1	Eye Pad
1	Safety Pin
4	Exam Quality Vinyl Gloves
1	Tweezer Plastic
10	Cotton Tipped Applicators
2	Ibuprofen 200mg
2	Aspirin 325mg
2	Non-Aspirin 325mg
1	Instant Cold Pack
1	Triangle Bandage
1	First Aid Guide
1	Case

**122 # of Pieces**

Qty	Descripción de Artículos
24	Venda Adhesiva 3" x 3/4"
6	Venda Adhesiva 1" x 3"
12	Venda Adhesiva de Tela 1" x 3"
3	Mando Cinta Largo
12	Vendajes Junior 3/8" x 1 1/2"
2	Venda Para Nudillo de Yelo
2	Vendaje para las puntas de los dedos de Yelo
2	Cierre de Herida Mariposa
2	Gasas 2" x 2"
4	Gasas 3" x 3"
1	Almohadilla de Trauma 5" x 9"
1	Roll de Gasas de 2"
1	Roll de Cinta Primeros Auxilios 1/2" x 5yd.
10	Toallitas Antisépticas (sin ardor)
6	Pomada Antibiótica
6	Crema de Primeros Auxilios Para Quemaduras
1	Venda Anti Ampolla 2" x 2"
1	Almohadilla para los Ojos
1	Imperdible
4	Guantes de Examen de Calidad de Vinilo
1	Pinesa de Plástico
10	Aplicadores con punta de Algodón
2	Tabletas de Ibuprofen 200mg
2	Aspirina 325mg
2	Tabletas no-aspirina 325mg
1	Compresas Frías Instantáneas
1	Vendaje Triángulo
1	Guía de Primeros Auxilios
1	Cuadro

**122 # de Piezas**



Manufactured in China  
for Ready America, Inc.  
Escondido, CA 92029  
[www.readyamerica.com](http://www.readyamerica.com)



Caution! This kit meets ANSI/ISEA Z308.1-2009 only when required minimum III is maintained with first aid products marked ANSI/ISEA Z308  
¡Cuidado! Este kit cumple con ANSI/ISEA Z308.1-2009 sólo cuando lo mínimo requerido se mantiene con productos de primeros auxilios marcados ANSI/ISEA Z308.

## GENUINE FIRST AID INDUSTRIAL FIRST AID

aspirin, acetaminophen, ibuprofen, bacitracin zinc, neomycin sulfate, polymyxin b, benzalkonium chloride, lidocaine, benzalkonium chloride kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52124-40 16
--------------	----------------	--------------------	-----------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-40 16-6	1 in 1 KIT		

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	10 PACKAGE	8.0 mL
Part 2	6 PACKET	5.4 g
Part 3	6 TUBE	3.0 g
Part 4	1 PACKET	2
Part 5	1 PACKAGE	2
Part 6	1 PACKAGE	2

### Part 1 of 6

#### ANTISEPTIC

benzalkonium chloride swab

## Product Information

Item Code (Source) NDC:52124-0001

Route of Administration TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	04/23/2010	

## Part 2 of 6

### GENUINE FIRST AID BURN ANTISEPTIC PAIN RELIEF WITH ALOE

benzalkonium chloride, lidocaine cream

## Product Information

Item Code (Source) NDC:52124-0004

Route of Administration TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	0.5 g in 100 g

## Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0004-1	0.9 g in 1 PACKET		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333A	04/13/2011	

## Part 3 of 6

### GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc,neomycin sulfate,polymyxin b sulfate ointment

## Product Information

Item Code (Source)	NDC:52124-0003
Route of Administration	TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

## Inactive Ingredients



Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0003-1	0.5 g in 1 TUBE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	04/13/2011	

### Part 4 of 6

### IBUPROFEN

ibuprofen tablet

### Product Information

Item Code (Source)	NDC:52124-0009
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

**Product Characteristics**

<b>Color</b>	white (White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;352
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0009-1	2 in 1 PACKET		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/17/2010	

**Part 5 of 6****NON-ASPIRIN**

acetaminophen tablet

**Product Information**

<b>Item Code (Source)</b>	NDC:52124-0010
<b>Route of Administration</b>	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

**Product Characteristics**

<b>Color</b>	white (White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;234
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0010-1	2 in 1 PACKAGE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/23/2010	

## Part 6 of 6

### ASPIRIN

aspirin tablet

## Product Information

Item Code (Source)	NDC:52124-0011
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STARCH, CORN (UNII: O8232NY3SJ)	

## Product Characteristics

Color	white (White)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0011-1	2 in 1 PACKAGE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	04/23/2010	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333E	05/03/2011	

**Labeler** - Genuine First Aid, LLC (619609857)

Revised: 7/2012

Genuine First Aid, LLC