

**ANTI-DIARRHEAL/ANTI-GAS MULTI-SYMPTOM RELIEF - loperamide hydrochloride 2mg, simethicone 125mg tablet  
Wal-Mart Stores, Inc.**

-----  
**Equate Anti-diarrheal/ Anti-gas Multi-Symptom Relief**

**ACTIVE INGREDIENTS (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**PURPOSES**

Anti-diarrheal

Anti-gas

**USE(S)**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**WARNINGS**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**DO NOT USE**

- if you have bloody or black stool
- if you have difficulty swallowing

**ASK A DOCTOR BEFORE USE IF YOU HAVE**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

## WHEN USING THIS PRODUCT

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

## STOP USE AND ASK DOCTOR IF

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

## IF PREGNANT OR BREAST FEEDING,

ask a health professional before use.

## KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## DIRECTIONS

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

## OTHER INFORMATION

- **each caplet contains:** calcium 180 mg, sodium 3 mg
- store between 20-25°C (68-77°F). Protect from light.
- **do not use if blister unit is torn or broken**

## **INACTIVE INGREDIENTS**

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

## **QUESTIONS OR COMMENTS**

1-888-287-1915

## **PRINCIPAL DISPLAY PANEL**

NDC 79903-116-33

equate™

Compare to Imodium®

Multi-Symptom Relief active ingredients\*\*

Loperamide Hydrochloride

and Simethicone Tablets, 2 mg/125 mg

**Anti-Diarrheal/**

**Anti-Gas**


Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas

24 Caplets\*

\*Capsule-Shaped Tablets



## ANTI-DIARRHEAL/ANTI-GAS MULTI-SYMPTOM RELIEF

loperamide hydrochloride 2mg, simethicone 125mg tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-116
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4I1)	LOPERAMIDE HYDROCHLORIDE	2 mg
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>ANHYDROUS DIBASIC CALCIUM PHOSPHATE</b> (UNII: L11K75P92J)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

STEARIC ACID (UNII: 4ELV7Z65AP)

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	16mm
<b>Flavor</b>	CARAMEL	<b>Imprint Code</b>	2;63
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-116-33	4 in 1 CARTON	07/15/2022	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214541	07/15/2022	

**Labeler** - Wal-Mart Stores, Inc. (051957769)

**Registrant** - Wal-Mart Stores, Inc. (051957769)

Revised: 7/2022

Wal-Mart Stores, Inc.