

IBUPROFEN- ibuprofen tablet, film coated
COSTCO WHOLESALE CORPORATION

Kirkland 44-291

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- backache
- menstrual cramps
- toothache
- headache
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling
- shock
- asthma (wheezing)
- skin reddening
- hives
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- slurred speech
- leg swelling
- trouble breathing
- weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- use by expiration date on package
- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

KIRKLAND
Signature

COMPARE TO
ADVIL®

*active ingredient***

NDC 63981-291-14
ITM. / ART. 808082

IBUPROFEN
Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

Actual Size

500
Tablets

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed
by PF Consumer Healthcare 1 LLC, owner of the
registered trademark Advil® .

Made in the USA

Manufactured by: LNK INTERNATIONAL, INC.
60 Arkay Drive, Hauppauge, NY 11788 USA
For: Costco Wholesale Corporation
P.O. Box 34535, Seattle, WA 98124-1535 USA
1-800-774-2678 www.costco.com 20V09017a
50844 REV1221F29114

No print/No varnish area
Lot no/Exp date

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil®.

Made in the USA

PEEL HERE FOR MORE DRUG FACTS

KIRKLAND Signature

COMPARE TO ADVIL®
active ingredient**

NDC 63981-291-14
ITM. / ART. 808082

IBUPROFEN
Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

500 Tablets

Actual Size

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each brown tablet) **Purpose**

Ibuprofen USP, 200 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - backache ■ menstrual cramps ■ toothache
 - headache ■ the common cold ■ muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling ■ shock ■ asthma (wheezing)
- skin reddening ■ hives ■ blisters ■ rash

If an allergic reaction occurs, stop use and seek medical help right away.

Manufactured by: LNK INTERNATIONAL, INC.
60 Arkay Drive, Hauppauge, NY 11788 USA
For: Costco Wholesale Corporation
P.O. Box 34535, Seattle, WA 98124-1535 USA
1-800-774-2678 www.costco.com
20V09017a
50844 REV1221F29114

PLASTIC BOTTLE

how2recycle.info

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ have bloody or black stools ■ vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ slurred speech
 - leg swelling ■ trouble breathing
- weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- use by expiration date on package
- store between 20°-25° C (68°-77° F)
- avoid excessive heat 40° C (104° F)

Inactive ingredients canauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
Call 1-800-426-6391 8:30 AM-4:00 PM ET, Monday-Friday

STOP PEELING

Kirkland 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:63981-291 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-----------------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|-----------------|
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK) | |
| MAGNESIUM STEARATE (UNII: 70097M6I3O) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | brown | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | 44;291 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:63981-291-14 | 500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 05/24/1988 | |
| 2 | NDC:63981-291-03 | 1 in 1 PACKAGE | 05/24/1988 | 08/23/2021 |
| 2 | | 10 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing | Application Number or Monograph | Marketing Start | Marketing End |
|------------------|--|------------------------|----------------------|
|------------------|--|------------------------|----------------------|

| Category | Citation | Date | Date |
|----------|------------|------------|------|
| ANDA | ANDA075010 | 05/24/1988 | |

Labeler - COSTCO WHOLESALE CORPORATION (103391843)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 038154464 | pack(63981-291) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867837 | manufacture(63981-291) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867894 | manufacture(63981-291) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|--|
| LNK International, Inc. | | 868734088 | manufacture(63981-291) , pack(63981-291) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 967626305 | pack(63981-291) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 117025878 | manufacture(63981-291) |