OMEPRAZOLE - omeprazole tablet, delayed release A-S Medication Solutions

Omeprazole Delayed-Release Tablets, 20 mg/acid reducer

Active ingredient (in each tablet)

Omeprazole delayed-release tablet, 20 mg (equivalent to 20.6 mg omeprazole magnesium USP)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

Do not use if you are allergic to omeprazole.

- Omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening blisters rash

If an allergy reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

• had heartburn over 3 months. This may be a sign of a more serious condition.

- frequent wheezing, particularly with heartburn
- unexplained weight loss nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25° C (68° to 77° F) and protect from moisture

Inactive ingredients

colloidal silicon dioxide, crospovidone, glyceryl monostearate, hypromellose, hydroxypropyl cellulose, iron oxide red, magnesium stearate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, silicified microcrystalline cellulose, sodium hydroxide, sodium lauryl sulphate, starlac (contains lactose monohydrate and corn starch), sugar spheres (contains sucrose and corn starch), talc, titanium dioxide, triethyl citrate.

Questions?

1-888-588-1418

HOW SUPPLIED

Product: 50090-6888

NDC: 50090-6888-0 14 TABLET, DELAYED RELEASE in a BOTTLE / 2 in a CARTON

Omeprazole



omeprazole tablet, delayed release

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50090-6888(NDC:69230-318)
--------------	----------------	--------------------	-------------------------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE (UNII: KG604840X9) (OMEPRAZOLE - UNII: KG604840X9)	OMEPRAZ OLE	20 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPOVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
GLYCERYL MONOSTEARATE (UNII: 2300U9XXE4)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
SUCROSE (UNII: C151H8M554)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	pink	Score	no score	
Shape	OVAL	Size	14mm	
Flavor		Imprint Code	H;O9	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50090- 6888-0	2 in 1 CARTON	12/06/2023		
1		14 in 1 BOTTLE; Type 0: Not a Combination			

1	Product			
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA211732	03/25/2020		

Labeler - A-S Medication Solutions (830016429)

Establishment				
Na me	Address	ID/FEI	Business Operations	
A-S Medication Solutions		830016429	RELABEL(50090-6888)	

Revised: 12/2023 A-S Medication Solutions