

**CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLU-  
acetaminophen, dextromethorphan hydrobromide, guaifenesin capsule, liquid  
filled  
Bayer HealthCare LLC.**

-----  
**Coricidin HBP Maximum Strength Cold, Cough & Flu Liquid Gels, UI1614532**

***Drug Facts***

**Active Ingredients**

***Active ingredients (in each capsule) Purposes***

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan hydrobromide 10 mg.....Cough suppressant  
Guaifenesin 200 mg.....Expectorant

**Purpose**

**Uses**

***Uses***

- temporarily relieves these symptoms due to a cold or flu: · minor aches and pains · headache · cough · sore throat
- helps loosen phlegm ( mucus) and thin bronchial secretions to make cough more productive
- temporarily reduces fever

**Warnings**

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe

allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

### **Ask a doctor before use if you have**

#### **Ask a doctor before use if you have**

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough is accompanied by excessive phlegm (mucus)

### **Ask a doctor or pharmacist**

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

#### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

· cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use.

### **Keep out of reach of children**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

#### ***Directions***

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

### **Other information**

#### ***Other information***

- store at 15°-25°C (59°-77°F)

### **Inactive ingredients**

***Inactive ingredients*** FD&C red No.40, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

### **Questions or comments**

**Questions or comments? 1-800-317-2165** (Mon-Fri 9AM -5PM EST)

### **Package label - 24 count carton**

Coricidin®

HBP

COLD RELIEF FOR PEOPLE WITH

**HIGH BLOOD PRESSURE †**

# MAXIMUM STRENGTH COLD, COUGH & FLU

Relieves

- Body aches & Pains
- Headache
- Fever
- Cough
- Sore Throat
- Chest Congestion
- 

†Decongestant Free

24 LIQUID GELS

(Liquid-Filled Capsules)

**Drug Facts**  
**Active ingredients (in each capsule)**  
Acetaminophen 325 mg Pain reliever/fever reducer  
Dextromethorphan hydrobromide 10 mg Cough suppressant  
Guaifenesin 200 mg Expectorant

**Purposes**  
■ temporarily relieves these symptoms due to a cold or flu:  
■ minor aches and pains ■ headache  
■ cough ■ sore throat  
■ helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive  
■ temporarily reduces fever

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:  
■ skin redness ■ blisters ■ rash ■ hives  
■ facial swelling ■ asthma (wheezing) ■ shock  
If a skin or general allergic reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients  
■ in children under 12 years of age

**Ask a doctor before use if you have**  
■ liver disease  
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema  
■ cough is accompanied by excessive phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Drug Facts (continued)**  
**Stop use and ask a doctor if**  
■ pain or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than the recommended dose  
■ adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.  
■ children under 12 years of age: do not use

**Other information**  
■ store at 15°-25°C (59°-77°F)

**Inactive ingredients** FD&C red No. 40, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, polyethylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or comments? 1-800-317-2105 (Mon-Fri 9AM - 5PM EST)**

**Drug Facts (continued)**  
**Stop use and ask a doctor if**  
■ pain or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Directions**  
■ do not take more than the recommended dose  
■ adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.  
■ children under 12 years of age: do not use

**Other information**  
■ store at 15°-25°C (59°-77°F)

**Inactive ingredients** FD&C red No. 40, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, polyethylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or comments? 1-800-317-2105 (Mon-Fri 9AM - 5PM EST)**

**Coricidin HBP**  
COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE!  
**MAXIMUM STRENGTH**  
COLD, COUGH & FLU 24 LIQUID GELS

**Coricidin HBP**  
COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE!  
**MAXIMUM STRENGTH**  
COLD, COUGH & FLU 24 LIQUID GELS

**Coricidin HBP**  
COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE!  
**MAXIMUM STRENGTH**  
COLD, COUGH & FLU 24 LIQUID GELS

**Relieves**  
• Body Aches & Pains  
• Headache  
• Fever  
• Cough  
• Sore Throat  
• Chest Congestion

†Decongestant Free

24 LIQUID GELS  
(Liquid Filled Capsules)

**LEARN MORE**  
American Heart Association  
Information Enclosed

**PARENTS:**  
Maximum Strength Per 4 Hour Dose  
Do not use if blister is cut or broken.  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**LOT/EXP**  
**LOT & EXP CODE AREA**

**Bayer**  
Bayer, the Bayer Cross, and Coricidin are registered trademarks of Bayer.  
Product of Spain  
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981  
87125661

**CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLU**  
acetaminophen, dextromethorphan hydrobromide, guaifenesin capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11523-0006
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>POTASSIUM ALUMINUM DISILICATE</b> (UNII: SRB14JRX6C)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	CHBPD
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11523-0006-1	2 in 1 CARTON	07/01/2020	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2020	

**Labeler** - Bayer HealthCare LLC. (112117283)

Revised: 11/2023

Bayer HealthCare LLC.