

**INON ACE- magnesium aluminosilicate, magnesium hydroxide solution**  
**Sato Pharmaceutical Co., Ltd.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**DRUG FACTS**

**Active ingredients (in each bottle)**

Magnesium Aluminosilicate 900 mg

Magnesium Hydroxide 250 mg

**Purpose**

Magnesium Aluminosilicate Antacid

Magnesium Hydroxide Antacid

**Uses** relieves these symptoms:

- heartburn
- sour stomach
- acid indigestion

**Warnings**

**Do not use** this product if you are taking a prescription antibiotic drug containing any form of tetracycline.

**Ask a doctor or pharmacist before use if you are** presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**When using this product**

- do not take more than 2 bottles (adults and children 12 years and over) in a 24-hour period or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.
- may have a laxative effect.

☐ **If pregnant or breast-feeding,** ☐ ask a health professional before use.

**Keep out of reach of children.**

**Directions**

- adults and children 12 years and over: Take 1 bottle (30mL) at a time 2 times daily, between meals, or as directed by a doctor.
- children under 12 years: Ask a doctor.
- shake well before use.

**Other information**

- each bottle contains: sodium 15 mg, magnesium 165 mg

**Inactive ingredients** alcohol, butylparaben, citric acid, flavors (cinnamon oil, fluid extract of ginger root, spearmint oil), hydroxypropylstarch, PEG-60 hydrogenated castor oil, propylparaben, sodium benzoate, sodium citrate, sorbitol, water.

**Drug Facts**

**Active ingredients (In each bottle) Purpose**  
Magnesium aluminometasilicate 900mg..... Antacid  
Magnesium hydroxide 250mg..... Antacid

**Uses** relieves these symptoms.  
■ heartburn ■ sour stomach ■ acid indigestion

**Warnings**  
Do not use this product if you are taking a prescription antibiotic drug containing any form of tetracycline.

**Ask a doctor or pharmacist before use if you are** presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**When using this product**  
Do not take more than 2 bottles (adults and children 12 years and over) in a 24-hour period or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.  
■ may have a laxative effect.

**If pregnant or breast-feeding, ask a health professional** before use. **Keep out of reach of children.**

**Directions**  
■ adults and children 12 years and over: Take 1 bottle (30ml) at a time 2 times daily, between meals, or as directed by a doctor.  
■ children under 12 years: Ask a doctor.  
■ shake well before use.

**Other information** ■ each bottle contains: sodium 15mg, magnesium 105mg  
■ tamper-evident Feature-The cap (printed "Sato") cannot be removed without breaking the ring crimped over the flange on the neck of the bottle. Do not use if ring is broken away from cap, missing, or damaged.

**Inactive ingredients** alcohol, butylparaben, citric acid, flavors (cinnamon oil, ginger fluid extract, mentha oil), hydroxypropyl corn starch, PEG-60 hydrogenated castor oil, propylparaben, sodium benzoate, sodium citrate, sorbitol, water.

Manufactured by  
**SATO PHARMACEUTICAL CO., LTD.**  
1-5-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN



**医薬品表示**

有効成分・分量  
1瓶(30 mL)中 配合目的  
メタケイ酸アルミン酸マグネシウム 900mg..... 制酸剤  
水酸化マグネシウム 250mg..... 制酸剤

**効能**  
下記症状の緩和  
■胸やけ ■胃酸過多

**注意**  
胃酸に強いコド、腸内細菌叢の減少の発生物質が腸を傷めている場合は、本剤を服用しないでください。

医師又は薬剤師に相談すること。制酸剤はある種の処方薬と相互作用を起こす場合があります。現在処方薬を服用している場合は医師又は薬剤師にご相談ください。

**本剤の服用に際して**  
■大人及び12歳以上の小児は、2週間以内、年に2瓶を超えて服用しないでください。また長期間の服用や過量服用は、副作用を発生(嘔吐)する危険性の上昇を伴うため、服用しないでください。

妊娠中または授乳中の方は、服用前に必ず医師又は薬剤師に相談してください。小児の手の届かないところに保管してください。

**用法・用量**  
■大人及び12歳以上の小児：1日1瓶(30ml)、1日2回食後に服用します。  
■12歳未満の小児：医師にご相談ください。  
■服用前に瓶をよく振ってください。

**他の情報** ■1瓶中：ナトリウム 15mg、マグネシウム 105mgを含有します。  
■タムパシ防止機能：キャップ("Sato")の文字(入り)は印刷すると、下部のリングが折れて外れます。もしリングがキャップから外れていたり、なを壊しているか、変形したりしている場合には、使用しないでください。

**他の成分** アルコール、ブチルパラベン、フェニルパラベン、香料(クミン油、シロウキョウ油、エッセンス、ハッカ油)、ヒドロキシプロピルコーンスターチ、ポリオキシエチレン硬脂酸セパン酸、プロピルパラベン、安息香酸ナトリウム、クエン酸ナトリウム、ソルビトール、精製水

Reports of serious side effects associated with use of this product can be sent to: 30094 S. Western Ave., Suite 240, Torrance, CA 90501

Magnesium aluminometasilicate: Antacid  
Magnesium hydroxide: Antacid

**INON ACE SOLUTION**  
イノンエス液

**sato**

NDC 49873-403-01

Magnesium aluminometasilicate: Antacid  
Magnesium hydroxide: Antacid

メタケイ酸アルミン酸マグネシウム:制酸剤  
水酸化マグネシウム:制酸剤

**INON ACE SOLUTION**  
イノンエス液


胸やけ、胃酸過多に  
For the relief of heartburn, sour stomach and acid indigestion.

1 FL OZ (30 mL) × 2 BOTTLES  
1 FL OZ (30 mL) × 2 瓶



EXP. DATE  
使用期限  
LOT  
製造番号

US-V



## INON ACE

magnesium aluminosilicate, magnesium hydroxide solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49873-403
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SILODRATE (UNII: 9T3UU8T0QK) (SILODRATE - UNII:9T3UU8T0QK)	SILODRATE	900 mg in 30 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	250 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
BUTYLPARABEN (UNII: 3QP1U3FV8)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
HYDROXYPROPYL CORN STARCH (5% SUBSTITUTION BY WEIGHT) (UNII: 9M44R3409A)	
POLYOXYL 60 HYDROGENATED CASTOR OIL (UNII: 02NG325BQG)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0K00R)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CINNAMON (CINNAMON OIL) , MINT (MENTHA OIL) , GINGER (GINGER, LIQUID EXTRACT)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-403-01	2 in 1 CARTON		
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	05/20/2000	

**Labeler** - Sato Pharmaceutical Co., Ltd. (690575642)

### Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-403) , label(49873-403) , pack(49873-403)

Revised: 12/2015

Sato Pharmaceutical Co., Ltd.