

BISACODYL- bisacodyl suppository
GERI-CARE PHARMACEUTICAL CORP

Bisacodyl Suppositories 443

Drug Facts

Active ingredient (in each suppository)

Bisacodyl USP, 10 mg

Purpose

Stimulant Laxative

Uses

- relieves occasional constipation
- this product generally produces a bowel movement in 15 minutes to 1 hour

Warnings

For rectal use only

Do not use

when abdominal pain, nausea, or vomiting are present.

Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period 2 weeks.

When using this product you may experience

- rectal burning
- mild cramps
- abdominal discomfort
- faintness

Stop use and ask a doctor if

- you have rectal bleeding or you fail to have bowel movement after using a laxative. These may be signs of a serious condition.
- you need to use a laxative product for more than one week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- tear off one suppository from the strip
- peel apart the wrapper
- continue peeling apart to almost the full length of the suppository (see illustration at the right)
- gently remove the suppository from the wrapper
- lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out

- retain it for about 15 to 20 minutes.
- **Do not exceed recommended dose**
- if suppository seems soft, place in refrigerator for a short time before use

adults and children 12 years and over	1 suppository once daily
children under 12 years	Ask a doctor

Other information

- store at temperatures below 25 C (77F)
- Made in India

Inactive ingredient hard fat

Questions or Comments? 1-800-540-3765



The image shows the front of a box for Bisacodyl Suppositories. The box has a green and yellow color scheme. At the top, the word "Suppositories" is written in a large, white, stylized font on a green background. Below this, the brand name "Bisacodyl" is written in a similar font, also in white on a green background. In the center, there is a logo for "HEALTH STAR" featuring a blue and yellow star with a red outline. Below the logo, the text "Fast Acting Stimulant Laxative" is written in white on a green background. At the bottom, the word "Suppositories" is written again in white on a green background. On the left side of the box, the text "50 Suppositories" and "10 mg each" is written in white on a green background. On the right side, there is a warning: "TAMPER EVIDENT: SUPPOSITORIES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED." written in white on a green background. The entire box is set against a white background.

Drug Facts

Active ingredient (in each suppository)

Bisacodyl USP 10 mg.....Stimulant laxative

Purpose

Uses ■ relieves occasional constipation ■ this product generally produces a bowel movement in 15 minutes to 1 hour



*This product is not manufactured or distributed by the owner of the registered trademark Dulcolax®

Distributed by: Qualitabs Inc.
Brooklyn, NY 11204

Questions or Comments: 1-800-540-3765

REV1119

Drug Facts (continued)

Warnings For rectal use only

Do not use when abdominal pain, nausea, or vomiting are present.

Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.

When using this product you may experience ■ rectal burning ■ mild cramps ■ abdominal discomfort ■ faintness


Stop use and ask a doctor if ■ you have rectal bleeding or you fail to have a bowel movement after using a laxative. These may be signs of a serious condition. ■ you need to use a laxative for more than one week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- tear off one suppository from the strip
- peel apart the wrapper
- continue peeling apart to almost the full length of the suppository (*see illustration at the right*)
- gently remove the suppository from the wrapper
- lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out
- retain it for 15 to 20 minutes ■ **Do not exceed recommended dose**
- if suppository seems soft, place in refrigerator for a short time before use



Adults and children 12 years and over	1 suppository once daily
Children under 12 years	Ask a doctor

Other information ■ store at temperatures below 25°C (77°F) ■ Made in India

Inactive ingredient hard fat

BISACODYL

bisacodyl suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-443
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg

Inactive Ingredients

Ingredient Name	Strength
FAT, HARD (UNII: 8334LX7S21)	

Product Characteristics

Color	white	Score	
Shape	BULLET	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-443-12	2 in 1 CARTON	11/01/2019	
1	NDC:57896-443-66	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:57896-443-05	10 in 1 CARTON	11/01/2019	
2	NDC:57896-443-55	5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:57896-443-01	20 in 1 CARTON	11/01/2019	
3	NDC:57896-443-55	5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	11/01/2019	

Labeler - GERI-CARE PHARMACEUTICAL CORP (611196254)**Registrant** - GERI-CARE PHARMACEUTICAL CORP (611196254)