

**ASPIRIN LOW DOSE- aspirin tablet, delayed release  
DOLGENCORP, LLC**

-----  
**Rexall 44-600A**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor.  
**Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - feel faint
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

### ***Inactive ingredients***

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

### ***Principal display panel***

***Since 1903***

**REXALL®**

Enteric Safety Coated

**81 mg Aspirin**

Pain Reliever (**NSAID**)

Low Dose Aspirin regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

- Easy open cap

**NOT FOR HOUSEHOLDS  
WITH YOUNG CHILDREN**

**300**

**Coated**

**Tablets**

**Actual Size**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

50844      REV0122C60017

**Visit us at: [Rexall.com](http://Rexall.com) or call 1-866-4-REXALL**

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE, GOODLETTSVILLE, TN 37072 USA

Since 1903 Rexall® is a registered trademark of Rexall Sundown, Inc.

**Drug Facts (continued)**

- if you have ever had an allergic reaction to this product or any of its ingredients
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - feel faint
  - have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients** corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**STOP PEELING**

Since 1903  
**Rexall**<sup>®</sup>

Enteric Safety Coated  
**81 mg Aspirin**  
Pain Reliever (NSAID)  
Low Dose Aspirin regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

Actual Size

300 Coated Tablets

Easy open cap

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*

**Purpose**  
Pain reliever

**Uses**  
for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

50844  
REXALL  
Visit us at: [Rexall.com](http://Rexall.com) or call 1-866-4-REXALL  
DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE, GOODLETTSVILLE, TN 37072 USA

Since 1903 Rexall<sup>®</sup> is a registered trademark of Rexall Sundown, Inc.

A0531

3 50844 60017 2

**PEEL HERE FOR MORE DRUG FACTS**

**Rexall 44-600A**

**ASPIRIN LOW DOSE**

aspirin tablet, delayed release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-563
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-563-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/01/2011	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2011	

**Labeler** - DOLGENCORP, LLC (068331990)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-563)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-563)

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		868734088	manufacture(55910-563) , pack(55910-563)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(55910-563)

Revised: 3/2024

DOLGENCORP, LLC