STOOL SOFTENER WITH STIMULANT LAXATIVE- docusate sodium and sennosides tablet P & L Development, LLC

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg Sennosides 8.6 mg

Purpose

Stool softener

Stimulant laxative

Uses

- relief occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless directed by a doctor
- if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

adults and children 12 years and older	take 2-4 tablets daily
children 6 to under 12 years of age	take 1-2 tablets daily
children 2 to under 6 years of age	take up to 1 tablet daily
children under 2	ask a doctor

Other information

• each tablet contains: calcium 30 mg

• each tablet contains: sodium 6 mg VERY LOW SODIUM

• store at 25°C (77°F); excursions permitted between 15-30°C(59-86°F)

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sillcon dioxide, talc, titanium dioxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

stool softener + stimulant laxative

docusate sodium 50 mg

(stool softener)

sennosides 8.6 mg

(stimulant laxative)

tablets

*Compare to the active ingredients in Colace® 2-IN-1


TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Avrio Health LP., distributor of Colace® 2-IN-1

Distributed by: **PL Developments**

200 Hicks Street, Westbury, NY 11590

Product Label



cellulose, polyethylene glycol, silicon dioxide, talc, titanium Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, each tablet contains: calcium 30 mg
each tablet contains: sodium 6 mg VERY LOW SODIUM hypromellose, magnesium stearate, microcrystalline medical help or contact a Poison Control Center right away Keep out of reach of children. In case of overdose, get Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. noticed a sudden change in bowel habits that **Drug Facts** (continued) Directions
■ take only by mouth. Doses may be taken as a single Ask a doctor before use if you have Call 1-877-753-3935 Monday-Friday 9AM-5PM EST Questions or comments? Other information before use. These may indicate a serious condition. over a period of 2 weeks store at 25°C (77°F); excursions permitted between daily dose, preferably in the evening, or in divided doses pregnant or breast-feeding, ask a health professional stomach pain ■ nausea children under 2 children 2 to under 6 12 years of age adults and children 12 years and over years of age children 6 to under 15-30°C (59-86°F) take up to 1 tablet daily ask a doctor take 1-2 tablets daily take 2-4 tablets daily ■ vomiting lasts

WELLNESS BASICS Stool Softener Laxative

STOOL SOFTENER WITH STIMULANT LAXATIVE

docusate sodium and sennosides tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-855	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	50 mg	
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	8.6 mg	

Ingredient Name CROSCARMELLOSE SODIUM (UNII: M280L1HH48) DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP) FD&C RED NO. 40 (UNII: WZB9127XOA) HYPROMELLOSES (UNII: 3NXW29V3WO) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3W)Q0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15EIX9V/2IP)	Inactive Ingredients				
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP) FD&C RED NO. 40 (UNII: WZB9127XOA) HYPROMELLOSES (UNII: 3NXW29V3WO) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	Ingredient Name	Strength			
FD&C RED NO. 40 (UNII: WZB9127XOA) HYPROMELLOSES (UNII: 3NXW29V3WO) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
HYPROMELLOSES (UNII: 3NXW29V3WO) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)				
MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	FD&C RED NO. 40 (UNII: WZB9127XOA)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	HYPROMELLOSES (UNII: 3NXW29V3WO)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	MAGNESIUM STEARATE (UNII: 70097M6I30)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4) TALC (UNII: 7SEV7J4R1U)	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
TALC (UNII: 7SEV7J4R1U)	POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
	SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
TITANIHM DIOYIDE (HNII: 15EIYOVZIP)	TALC (UNII: 7SEV7J4R1U)				
TTARION DIOXIDE (ONI. 1511X3V2)1)	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
ALUMINUM OXIDE (UNII: LMI2606933)	ALUMINUM OXIDE (UNII: LMI26O6933)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				

Product Characteristics				
Color	red	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	PSD21	
Contains				

Ш	Packaging					
	# Item Code	Package Description	Marketing Start Date	Marketing End Date		
	NDC:59726- 855-40	400 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2020			
	NDC:59726- 855-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2020			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	04/30/2020		

Labeler - P & L Development, LLC (800014821)

Revised: 4/2024 P & L Development, LLC