

# **CLOBETASOL PROPIONATE- clobetasol propionate ointment** **AvKARE**

-----  
**CLOBETASOL PROPIONATE OINTMENT USP, 0.05%**


**FOR TOPICAL DERMATOLOGIC USE ONLY.**

**NOT FOR OPHTHALMIC, ORAL OR INTRAVAGINAL USE.**

## **DESCRIPTION:**

Clobetasol Propionate Ointment USP contains the active compound clobetasol propionate, a synthetic corticosteroid, for topical dermatologic use. Clobetasol, an analog of prednisolone, has a high degree of glucocorticoid activity and a slight degree of mineralocorticoid activity.

Chemically, clobetasol propionate is (11 $\beta$ ,16 $\beta$ )-21-chloro-9-fluoro-11-hydroxy-16-methyl-17-(1-oxopropoxy)-pregna-1,4-diene-3,20-dione, and it has the following structural formula:



Clobetasol propionate USP has the empirical formula C<sub>25</sub>H<sub>32</sub>ClFO<sub>5</sub> and a molecular weight of 466.97. It is a white to almost white, crystalline powder, practically insoluble in water; slightly soluble in benzene and diethyl ether; sparingly soluble in ethanol; freely soluble in acetone, in dimethylsulfoxide, in chloroform, in methanol and in dioxane.

Clobetasol Propionate Ointment USP, 0.05% contains clobetasol propionate USP 0.5 mg/g in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

## **CLINICAL PHARMACOLOGY:**

Like other topical corticosteroids, clobetasol propionate has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A<sub>2</sub> inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent

mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A<sub>2</sub>.

**Pharmacokinetics:** The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle and the integrity of the epidermal barrier. Occlusive dressings with hydrocortisone for up to 24 hours has not been demonstrated to increase penetration; however, occlusion of hydrocortisone for 96 hours markedly enhances penetration. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin may increase percutaneous absorption.

Studies performed with clobetasol propionate ointment indicate that it is in the super-high range of potency as compared with other topical corticosteroids.

## **INDICATIONS AND USAGE:**

Clobetasol Propionate Ointment USP is a super-high potency corticosteroid formulation indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid responsive dermatoses. Treatment beyond 2 consecutive weeks is not recommended, and the total dosage should not exceed 50 g/week because of the potential for the drug to suppress the hypothalamic-pituitary-adrenal (HPA) axis. Use in pediatric patients under 12 years of age is not recommended.

As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of the diagnosis may be necessary.

## **CONTRAINDICATIONS:**

Clobetasol Propionate Ointment USP is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations.

## **PRECAUTIONS:**

### **General:**

Clobetasol Propionate Ointment USP should not be used in the treatment of rosacea or perioral dermatitis, and should not be used on the face, groin, or axillae.

Systemic absorption of topical corticosteroids can produce reversible HPA axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal from treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on therapy.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests. Patients receiving super-potent corticosteroids should not be treated for more than 2 weeks at a time and only small areas should be treated at any one time due to the

increased risk of HPA suppression.

Clobetasol Propionate Ointment USP produced HPA axis suppression when used at doses as low as 2 g/day for 1 week in patients with eczema.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Pediatric patients may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios (see **PRECAUTIONS: Pediatric Use**).

If irritation develops, Clobetasol Propionate Ointment USP should be discontinued and appropriate therapy instituted. Allergic contact dermatitis with corticosteroids is usually diagnosed by observing a *failure to heal* rather than noting a clinical exacerbation as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of Clobetasol Propionate Ointment USP should be discontinued until the infection has been adequately controlled.

### **Information for Patients:**

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged, otherwise covered or wrapped, so as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions to the physician.

### **Laboratory Tests:**

The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test

A.M. plasma cortisol test

Urinary free cortisol test.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility:**

Long-term animal studies have not been performed to evaluate the carcinogenic potential of clobetasol propionate.

Studies in the rat following subcutaneous administration at dosage levels up to 50 mcg/kg per day revealed that the females exhibited an increase in the number of

resorbed embryos and a decrease in the number of living fetuses at the highest dose.

Clobetasol propionate was non-mutagenic in three different test systems: the Ames test, the *Saccharomyces cerevisiae* gene conversion assay, and the *E. coli* B WP2 fluctuation test.

### **Pregnancy:**

**Teratogenic Effects:** Pregnancy Category C . Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application to laboratory animals.

Clobetasol propionate has not been tested for teratogenicity when applied topically; however, it is absorbed percutaneously, and when administered subcutaneously it was a significant teratogen in both the rabbit and mouse. Clobetasol propionate has greater teratogenic potential than steroids that are less potent.

Teratogenicity studies in mice using the subcutaneous route resulted in fetotoxicity at the highest dose tested (1 mg/kg) and teratogenicity at all dose levels tested down to 0.03 mg/kg. These doses are approximately 1.4 and 0.04 times, respectively, the human topical dose of Clobetasol Propionate Ointment USP. Abnormalities seen included cleft palate and skeletal abnormalities.

In rabbits, clobetasol propionate was teratogenic at doses of 3 and 10 mcg/kg. These doses are approximately 0.02 and 0.05 times, respectively, the human topical dose of Clobetasol Propionate Ointment USP. Abnormalities seen included cleft palate, cranioschisis, and other skeletal abnormalities.

There are no adequate and well-controlled studies of the teratogenic potential of clobetasol propionate in pregnant women. Clobetasol Propionate Ointment USP should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### **Nursing Mothers:**

Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when Clobetasol Propionate Ointment USP is administered to a nursing woman.

### **Pediatric Use:**

Safety and effectiveness of Clobetasol Propionate Ointment USP in pediatric patients have not been established. Use in children under 12 years of age is not recommended. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing's syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during or after withdrawal of treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children .

HPA axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include low plasma cortisol levels, and an absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

### **Geriatric Use:**

A limited number of patients at or above 65 years of age have been treated with clobetasol propionate ointment (n = 101) in US and non-US clinical trials. While the number of patients is too small to permit separate analysis of efficacy and safety, the adverse reactions reported in this population were similar to those reported by younger patients. Based on available data, no adjustment of dosage of Clobetasol Propionate Ointment USP in geriatric patients is warranted.

### **ADVERSE REACTIONS:**

In controlled clinical trials, the most frequent adverse events reported for Clobetasol Propionate Ointment USP were burning sensation, irritation, and itching in 0.5% of treated patients. Less frequent adverse reactions were stinging, cracking, erythema, folliculitis, numbness of fingers, skin atrophy, and telangiectasia.

Cushing's syndrome has been reported in infants and adults as a result of prolonged use of topical clobetasol propionate formulations.

The following additional local adverse reactions have been reported with topical corticosteroids, and they may occur more frequently with the use of occlusive dressings and higher potency corticosteroids. These reactions are listed in an approximately decreasing order of occurrence: dryness, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, irritation, striae, and miliaria.

### **OVERDOSAGE:**

Topically applied Clobetasol Propionate Ointment USP can be absorbed in sufficient amounts to produce systemic effects (see **PRECAUTIONS**).

### **DOSAGE AND ADMINISTRATION:**

Apply a thin layer of Clobetasol Propionate Ointment USP to the affected skin areas twice daily and rub in gently and completely. (See **INDICATIONS AND USAGE**.)

Clobetasol Propionate Ointment USP is a super-high potency topical corticosteroid; therefore, **treatment should be limited to 2 consecutive weeks, and amounts greater than 50 g/week should not be used.**

As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of diagnosis may be necessary.

Clobetasol Propionate Ointment USP should not be used with occlusive dressings.

## Geriatric Use:

In studies where geriatric patients (65 years of age or older, see **PRECAUTIONS**) have been treated with clobetasol propionate ointment, safety did not differ from that in younger patients; therefore, no dosage adjustment is recommended.

## HOW SUPPLIED:

Clobetasol Propionate Ointment USP, 0.05% is supplied in 15-g (NDC 42291-077-15), 30-g (NDC 42291-077-30), 45-g (NDC 42291-077-45), and 60-g (NDC 42291-077-60) tubes.

**Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° and 30°C (59° and 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.**

Manufactured For:

AvKARE

Pulaski, TN 38478

Product of Spain

04/2020 PI077-00

## PACKAGE LABEL- PRINCIPAL DISPLAY PANEL- 15 G CONTAINER

<p><b>EACH GRAM CONTAINS:</b> Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.</p> <p><b>USUAL DOSAGE:</b> See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.</p> <p><b>TO OPEN:</b> Use cap to puncture seal.</p> <p><b>IMPORTANT:</b> Do not use if seal has been punctured or is not visible. See crimp of tube for Lot Number and Expiration Date.</p> <p><b>Manufactured for:</b> AvKARE, Pulaski, TN 38478</p> <p><b>Product of Spain</b> Rev: 04/2020 077-15-00 569892-01</p>	 <p>7 07715 42291 3</p>
<p><b>AVKARE</b></p> <p><b>Clobetasol Propionate Ointment USP, 0.05%</b></p> <p><b>15 g</b></p> <p><b>R<sub>x</sub> only</b></p> <p>FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE. WARNING: Keep out of reach of children.</p>	<p><b>NDC 42291-077-15</b></p> <p>LOT:    EXP:</p>

**EACH GRAM CONTAINS:** Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:** See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.

**TO OPEN:** Use cap to puncture seal.

**IMPORTANT:** Do not use if seal has been punctured or is not visible. See crimp of tube for Lot Number and Expiration Date.

**Manufactured for:** AvKARE, Pulaski, TN 38478    **Product of Spain**    Rev: 04/2020    077-30-00    569893-01



**NDC 42291-077-30**

# Clobetasol Propionate Ointment USP, 0.05%

## 30 g

**R<sub>x</sub> only**

**FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE. WARNING: Keep out of reach of children.**

LOT:    EXP:

**EACH GRAM CONTAINS:** Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:** See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.

**TO OPEN:** Use cap to puncture seal.

**IMPORTANT:** Do not use if seal has been punctured or is not visible. See crimp of tube for Lot Number and Expiration Date.

**Manufactured for:** AvKARE, Pulaski, TN 38478

**Product of Spain** Rev: 04/2020 077-45-00

569894-01



**AVKARE.**

**NDC 42291-077-45**

# Clobetasol Propionate Ointment USP, 0.05%

**45 g**

**R<sub>x</sub> only**

**FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.**

LOT: EXP:



**EACH GRAM CONTAINS:** Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:** See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.

**TO OPEN:** Use cap to puncture seal.


**IMPORTANT:** Do not use if seal has been punctured or is not visible. See crimp of tube for Lot Number and Expiration Date.

**Manufactured for:** AvKARE, Pulaski, TN 38478

**Product of Spain**

Rev: 04/2020 077-60-00

569895-01



**NDC 42291-077-60**

# Clobetasol Propionate Ointment USP, 0.05%

## 60 g

**R<sub>x</sub> only**

**FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE. WARNING: Keep out of reach of children.**

LOT: EXP:

**PACKAGE LABEL- PRINCIPAL DISPLAY PANEL - 15 G CARTON**

NDC 42291-077-15



# Clobetasol Propionate Ointment USP, 0.05%

15 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

R<sub>x</sub> only

LOT:  
EXP:



**EACH GRAM CONTAINS:**

Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:**

See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.



Manufactured for:  
AvKARE, Pulaski, TN 38478  
Product of Spain  
Rev: 04/2020 077-15-00

NDC 42291-077-15



# Clobetasol Propionate Ointment USP, 0.05%

15 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

R<sub>x</sub> only

**IMPORTANT:** The opening of this product is covered by a metal tamper-resistant seal. If this seal has been punctured or is not visible, do not use and return product to the place of purchase.

**TO OPEN:** To puncture the seal, reverse the cap and place the puncture-top onto the tube. Push down firmly until seal is open.

**TO CLOSE:** Screw the cap back onto the tube.



56987401

NDC 42291-077-30



# Clobetasol Propionate Ointment USP, 0.05%

30 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

R<sub>x</sub> only

LOT:  
EXP:



**EACH GRAM CONTAINS:**

Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:**

See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.



Manufactured for:  
AvKARE, Pulaski, TN 38478  
Product of Spain  
Rev: 04/2020 077-30-00

NDC 42291-077-30



# Clobetasol Propionate Ointment USP, 0.05%

30 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

R<sub>x</sub> only

5698855

**IMPORTANT:** The opening of this product is covered by a metal tamper-resistant seal. If this seal has been punctured or is not visible, do not use and return product to the place of purchase.

**TO OPEN:** To puncture the seal, reverse the cap and place the puncture-top onto the tube. Push down firmly until seal is open.

**TO CLOSE:** Screw the cap back onto the tube.





NDC 42291-077-45

# Clobetasol Propionate Ointment USP, 0.05%

45 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

Rx only

LOT:  
EXP:



**EACH GRAM CONTAINS:**

Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:**

See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.



Manufactured for:  
AvKARE, Pulaski, TN 38478  
Product of Spain  
Rev: 04/2020 077-45-00



NDC 42291-077-45

# Clobetasol Propionate Ointment USP, 0.05%

45 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

Rx only



**IMPORTANT:** The opening of this product is covered by a metal tamper-resistant seal. If this seal has been punctured or is not visible, do not use and return product to the place of purchase.

**TO OPEN:** To puncture the seal, reverse the cap and place the puncture-top onto the tube. Push down firmly until seal is open.

**TO CLOSE:** Screw the cap back onto the tube.



1P-068895



NDC 42291-077-60

# Clobetasol Propionate Ointment USP, 0.05%

60 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

Rx only

LOT:  
EXP:



**EACH GRAM CONTAINS:**

Clobetasol Propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

**USUAL DOSAGE:**

See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Do not refrigerate.



Manufactured for:  
AvKARE, Pulaski, TN 38478  
Product of Spain  
Rev: 04/2020 077-60-00



NDC 42291-077-60

# Clobetasol Propionate Ointment USP, 0.05%

60 g

FOR DERMATOLOGIC USE ONLY. NOT FOR OPHTHALMIC USE.  
WARNING: Keep out of reach of children.

Rx only



**IMPORTANT:** The opening of this product is covered by a metal tamper-resistant seal. If this seal has been punctured or is not visible, do not use and return product to the place of purchase.

**TO OPEN:** To puncture the seal, reverse the cap and place the puncture-top onto the tube. Push down firmly until seal is open.

**TO CLOSE:** Screw the cap back onto the tube.



1P-068895

# CLOBETASOL PROPIONATE

clobetasol propionate ointment

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42291-077
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CLOBETASOL PROPIONATE</b> (UNII: 779619577M) (CLOBETASOL - UNII:ADN79D536H)	CLOBETASOL PROPIONATE	0.5 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITAN SESQUIOLEATE</b> (UNII: 0W8RRI5W5A)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-077-15	1 in 1 CARTON	06/18/2020	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:42291-077-30	1 in 1 CARTON	06/18/2020	
2		30 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:42291-077-45	1 in 1 CARTON	06/18/2020	
3		45 g in 1 TUBE; Type 0: Not a Combination Product		
4	NDC:42291-077-60	1 in 1 CARTON	06/18/2020	
4		60 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209701	06/18/2020	

**Labeler** - AvKARE (796560394)

